

Sodium bicarbonate ingestion and individual variability in time to peak pH

S. Andy Sparks¹, Emily Williams¹, Amy Robinson¹, Peter Miller¹, David Bentley², Craig
Bridge¹ and Lars R. Mc Naughton¹

1 Edge Hill University Exercise Nutrition Research Group, Department of Sport and
Physical Activity, Edge Hill University, Ormskirk UK

2 Flinders University, Adelaide, South Australia, Australia

Running Head: Variability in time to peak pH

Corresponding author:

Professor Lars Mc Naughton

Edge Hill University

Department of Sport and Physical Activity

St Helen's Road

Ormskirk L39 4QP

United Kingdom

Phone: +44 1704541967

Email: lars.mcnaughton@edgehill.ac.uk

Abstract

The aim of this study was to determine the individual variability in time to peak pH after the consumption of a $300\text{mg}\cdot\text{kg}^{-1}$ dose of sodium bicarbonate (NaHCO_3). Seventeen active males volunteered to participate in the study (mean \pm SD: age $21.38 \pm 1.5\text{y}$; mass $75.8 \pm 5.8\text{kg}$; height $176.8 \pm 7.6\text{cm}$). Participants reported to the laboratory where a resting capillary blood sample was taken aseptically from the fingertip. After this, $300\text{mg}\cdot\text{kg}^{-1}$ of NaHCO_3 in 400ml of water with 50ml of flavoured cordial was ingested. Participants then rested for 90 min during which repeated blood samples were procured at 10 minute intervals for 60 mins and then every 5 min until 90 min. Blood pH concentrations were measured using a blood gas analyser. Results suggested that time to peak pH (64.41 ± 18.78 min) was highly variable with a range of 10-85 min and a coefficient of variation of 29.16%. A bi-modal distribution occurred, at 65 and 75 min. In conclusion, researchers and athletes, when using NaHCO_3 as an ergogenic aid, should determine, in advance their time to peak pH to best utilise the added buffering capacity this substance allows.

Key Words: Performance, individual response, buffering, acidity

Introduction

Sodium bicarbonate ingestion is used a popular method of improving buffering against hydrogen ions induced by high intensity short duration exercise. There have been a number of review articles which have reaffirmed its effectiveness as an ergogenic aid when consumed prior to exercise performance lasting up to 10 minutes in duration ^{1, 2, 3}. A relatively recent meta-analysis of the effects of sodium bicarbonate ingestion on high intensity exercise performance suggested that the most effective pre-exercise doses should be between 0.3-0.5g/kg/BM, which is likely to improve mean power by $1.7 \pm 2.0\%$ ⁴ in appropriate exercise. Indeed it is this type of exercise that elicits the intracellular accumulation of hydrogen ions (H^+), which have been implicated as a cause of muscular fatigue⁵, typically occurring as a by-product of anaerobic glycolysis. This energy system is predominantly used in high intensity exercise, including repeated sprint activity (RSA), usually lasting less than a total of 10 minutes.

Although some evidence suggests that, at physiological temperatures, direct inhibition of force production by acidification is not as great as previously thought⁶, interventions that minimize intracellular H^+ accumulation may improve RSA. H^+ accumulation depends on both the production and removal of H^+ . The intra- and extracellular buffer systems act to reduce the build-up of free H^+ during high-intensity exercise and may therefore be important in maintaining repeated-sprint performance. Indeed, Bishop et al.⁷ have reported a significant relationship between RSA and both change in blood pH and *in vivo* muscle buffer capacity⁸. The intracellular accumulation of H^+ also depends on the extracellular H^+ concentration. H^+ efflux out of the muscle cell has been reported to be inhibited by extracellular acidosis⁹ and enhanced by a greater extracellular buffer concentration¹⁰. It is therefore frequently hypothesized that increases in the extracellular buffer concentration, via the ingestion of an alkaline solution such as sodium bicarbonate

(NaHCO₃), may improve H⁺ efflux out of the muscle cell and improve repeated-sprint performance¹¹.

It is now however generally accepted that the increased H⁺ production causes competition on the ionisable binding sites of the actin / myosin complex, as well as sarcoplasmic reticulum dysfunction with regard to Ca²⁺ release and uptake^{12, 13, 14}. With respect to both of these models of skeletal muscle fatigue, attenuating the increase in muscle (and subsequently blood) acidosis should help delay the onset of fatigue during repeated bouts of high intensity exercise, thereby helping to minimise the decline in power output inevitable during this type of exercise. Although not conclusive, it appears that increasing the blood buffering potential via NaHCO₃ ingestion either creates an electro-chemical gradient between the intra- and extracellular milieu, thus allowing for greater facilitation of proton removal from inside the cell; or sustains Ca²⁺ release and re-sequestering in the sarcoplasmic reticulum by increasing the strong ion difference¹⁵. Sustaining these mechanisms may prolong skeletal muscle function and perhaps maintain exercise performance, but the degree of efficacy in enhancing physical performance remains equivocal¹.

As a result of the mechanisms by which this supplement may act to delay fatigue, many laboratory investigations have used a variety of relevant exercise models including running^{16, 17, 18}, cycling^{11, 19, 20, 21}, boxing and swimming^{22, 23} in order to assess its effectiveness. Indeed, such is the wealth of published studies on sodium bicarbonate, that recently, researchers have started to focus on its co-ingestion with other active ingredients such as caffeine^{24, 25} and β-alanine^{26, 27, 28} in order to assess the potential additive effects in order to provide further performance enhancements via the activation of different ergogenic mechanisms simultaneously.

Christensen et al.²⁹ performed a study using elite rowers, which is one of the few investigations that have not demonstrated an ergogenic effect of sodium bicarbonate. Their protocol required participants to complete a 6 min time trial task following a dose of 3 g.kg⁻¹ ingested 75 min prior to the start of the performance test. An additional condition of the study also used this same dose of sodium bicarbonate co-ingested with a 3 mg.kg⁻¹ dose of caffeine. In the caffeine trials they observed a significant improvement in performance which was not observed in either the co-ingestion trial or with a single dose of sodium bicarbonate. Interestingly, there is some variety in the timing of pre-exercise administration in the literature which typically ranges from 60-90 min^{29, 30, 31, 32}. In some cases a multiple acute dose has been used starting at 90 mins and continuing until 50 min pre-exercise³³ or more chronic supplementation across several days³⁴. This range of pre-exercise ingestion times are likely to influence the effectiveness of the supplement and therefore the magnitude of the potential performance benefits which are reported. Presently there is no standardised pre-exercise ingestion time which has been determined as most effective, and there are also some suggestions that training status, diet and activity may affect buffering capacity. We hypothesize that these factors lead to considerable inter-individual variation in the time at which optimal buffering may occur following the ingestion of supplements designed to alter the pH of the blood. Therefore the aim of this experiment was to determine the variability in individual responses to a single bolus of sodium bicarbonate.

Methods

Participants

Seventeen male active team and individual sports participants (mean \pm SD: age 21.38 \pm 1.5y; mass 75.8 \pm 5.8kg; height 176.8 \pm 7.6cm) volunteered to take part in the study. All participants were familiar with high-intensity exercise and on took part in a minimum of

two hours of intermittent team or individual sporting activity per week. All participants were informed of both the benefits and the potential side effects associated with the study (both verbally and in writing), before they provided written informed consent and then underwent screening. The study was approved by the institutional Departmental Ethics Committee. Following screening

Procedures

The participants attended the laboratory once in order to obtain basic anthropometric measurements and to determine each individual's resting blood pH responsiveness to NaHCO₃ ingestion. Following the screening and anthropometric data collection participants ingested 300 mg·kg⁻¹ (BM) of NaHCO₃ taken in 400 ml of water with 50 ml of flavoured cordial (Robinsons Fruit Squash, UK). This method has previously been used by Price et al.³⁵ as it has been shown to improve drink palatability³⁶. Participants were asked to refrain from maximal exercise, to maintain a typical diet and avoid consuming alcohol and beverages other than water for the 24 hour period prior to their laboratory trial in order to minimise disturbances to normal acid-base status^{36, 37, 38}.

At the visit, participants reported to the laboratory where a 300 µl resting capillary blood sample was taken aseptically from the fingertip. The participants then consumed 300 mg·kg⁻¹ of NaHCO₃ in 400ml of water with 50ml of flavoured cordial within a five minute period. This dose has previously been found to improve individual anaerobic performance^{19, 39, 40} as well as repeated sprint performance⁴¹ in men and women^{19, 21, 39}. Participants then rested quietly for a 90 min period following the completion of ingestion. During this time additional capillary blood samples were procured at 10 minute intervals for the first 60 min and then at 5 min intervals until 90 min. Blood pH concentrations were measured using a blood gas analyser (Radiometer ABL800, Denmark).

Statistical Analysis

All data were assessed for normality using standard graphical methods prior to analysis⁴². Blood pH responses over the post ingestion period were assessed using repeated measures ANOVA. Post hoc pair-wise comparisons were made using a Bonferroni adjustment and statistical significance was assumed as $p < 0.05$. Calculations of effect sizes were done using partial eta squared (ηp^2) for ANOVA. The conventional interpretations of Cohen⁴³ were used to evaluate effect sizes where < 0.20 = trivial, $0.20-0.49$ = small, $0.50-0.79$ = moderate, and large ≥ 0.80 = large. All data are presented as mean \pm SD and were analysed using SPSS v22 for Windows (SPSS Inc., Chicago, IL, USA).

Results

The ingestion of the sodium bicarbonate bolus had a significant effect on pH ($f = 16.08$, $p < 0.001$, $\eta p^2 = 0.50$). Indeed the post ingestion pH values were all significantly higher than the pre-ingestion sample ($p < 0.05$). Most notably there was a significant increase in pH at the 10 ($p = 0.007$) and 20 min ($p < 0.001$) sample points compared to the pre ingestion values (Figure 1A). There was a further increase in pH after 40 min compared to the 20 min value ($p = 0.01$) after which pH did not significantly change until a decrease occurred between 75-80 min ($p = 0.03$). There were further significant decreases in pH between 75-85 min ($p = 0.006$) and 75-90 min ($p = 0.018$). Mean time to peak pH was 64.41 ± 18.78 min with a coefficient of variation of 29.16%. Furthermore between subject effects analysis revealed that there was significant variation in the pH responses ($f = 5830237.7$, $p < 0.001$, $\eta p^2 = 1.00$). The times to peak pH to determine the optimum loading period strategy, are shown in Table 1 with the range of times spread between

10-85 min (Figure 1B). Time to peak pH frequency was bi-modally distributed between 65 and 75 min.

The correlation between time to peak pH and time to bicarbonate peak time was $r=0.95$, ($p=0.001$). Peak pH achieved was not correlated to weight, with a low correlation ($r=0.07$, $p=0.79$) and neither was weight correlated to change in minimum-maximum pH achieved, $r=0.124$, $p=0.64$).

Discussion and Conclusion

The results of this study suggest that after ingestion of a bolus of $300\text{mg}\cdot\text{kg}^{-1}$ body mass of sodium bicarbonate, the time to reach peak pH is variable, with a range of 10-85 min. This suggests that when used as an ergogenic aid to improve sprint performance, in studies that have either used 60 min after ingestion^{23, 35} or 90 min post ingestion^{20, 21, 44}, the time lag is probably too short (60 min) or too long respectively. This is supported by the fact that the mean time to peak pH, across all participants is 65.0 ± 18.4 min, confirming that an exercise time of 60-90 min post ingestion is either too short, or too long respectively. Hence, this would then suggest that these participants are not making the most of the possible ergogenic, buffering capacity allowed by the ingestion of NaHCO_3 .

It is not possible to predict pH on the basis of a participant's body mass as the correlation between body mass and peak pH was low. Neither is it possible to predict maximum pH values on the basis of body weight. As has been seen previously^{20, 21, 44}, responses to the same dose of are varied with respect to changes in pH even when resting pH values are almost identical.

In conclusion, researchers, athletes and coaches should endeavour to undertake testing to ensure that if sodium bicarbonate is being used as an ergogenic aid, that their time to peak pH is known so that performance can be maximised at the time when peak pH is

achieved. This will require analysis with blood gas analysers which can be found in hospitals and at some academic institutions.

References

1. Mc Naughton LR, Siegler J, and Midgley A. Ergogenic Effects of Sodium Bicarbonate. *Curr Sports Med Rep.* 2008; 7: 230-236.
2. Peart DJ, Siegler JC, Vince RV. (2012). Practical recommendations for coaches and athletes: a meta-analysis of sodium bicarbonate use for athletic performance. *J Strength Cond Res* 26(7):1975-83. doi: 10.1519/JSC.0b013e3182576f3d
3. Burke LM, Pyne DB. Bicarbonate Loading to Enhance Training and Competitive Performance. *Int J Sports Physiol Perform.* 2007; 2: 93-96.
4. Carr AJ, Slater GJ, Gore CJ, Dawson B, and Burke LM (2011). Effect of Sodium Bicarbonate on [HCO₃⁻], pH and Gastrointestinal Symptoms. *Int J Sport Nutr Exerc Metab.* 2011; 21: 189-194.
5. Spriet LL, Matsos CG, Peters SJ, Heigenhauser GJ, Jones NL. Effects of acidosis on rat muscle metabolism and performance during heavy exercise. *Am J Physiol.* 1985; 24: C337-C347.
6. Westerblad H, Bruton J, Lannergren J. The effect of intracellular pH on contractile function of intact, single fibres of mouse declines with increasing temperature. *J. Physiol.* 1997; 500:193–204.
7. Bishop D, Lawrence M, Spencer M. Predictors of repeated sprint ability in elite female hockey players. *J. Sci. Med. Sport.* 2003; 6:199 –209.
8. Edge J, Bishop D, Goodman C, Davis C, Dawson B, Muscle buffer capacity and aerobic fitness predict repeated-sprint ability (RSA) in females. *Eur Congress Sport Science.* 2002; 100.
9. Hirche HJ, Hombach V, Langohr HD, Wacker U, Busse J. Lactic acid permeation rate in working gastrocnemii of dogs during metabolic alkalosis and acidosis. *Pflugers Arch.* 1975; 356: 209–222, 1975.

10. Mainwood GW, Worseley-Brown P. The effect of extracellular pH and buffer concentration on the efflux of lactate from frog sartorius muscle. *J. Physiol.* 1975; 250:1–22.
11. Miller P, Robinson A, Sparks SA, Bridge CA, Bentley D, and Mc Naughton LR. (2015). The effects of novel ingestion of sodium bicarbonate on repeated sprint ability. *J Strength and Cond Res* (In Press).
12. Allen DG, Lamb GD, Westerblad H. Impaired calcium release during fatigue. *J Appl Physiol.* 2008; 104:296 – 305
13. Fitts RH. The cross-bridge cycle and skeletal muscle fatigue. *J Appl Physiol.* 2008; 104: 551 – 558
14. Stephenson DG, Lamb GD, Stephenson GM. Events of the excitation-contraction-relaxation (E-C-R) cycle in fast- and slow-twitch mammalian muscle fibres relevant to muscle fatigue. *Acta Physiol Scand.* 1998; 162:229 – 245
15. Kemp G, Boning D, Beneke R, Maassen N. Explaining pH change in exercising muscle: lactic acid, proton consumption, and buffering vs. strong ion difference. *Am J Physiol.* 2006; 291: R235 – R237
16. Bird SR, Wiles J, Robbins J. The effect of sodium bicarbonate ingestion on 1500-m racing time, *J Sports Sci.* 1995, 13: 399-403
17. Goldfinch J, Mc Naughton LR, Davies P. Induced metabolic alkalosis and its effects on 400 m racing time. *Eur J Appl Physiol Occ Physiol.* 1988; 57: 45-48.
18. Tiryaki GR, Atterbom HA. The effects of sodium bicarbonate and sodium citrate on 600 m running time of trained females. *Jour Sports Med Phys Fit.* 1995, 35:194-198.
19. Mc Naughton LR, and Cedaro R. The effect of sodium bicarbonate on rowing ergometer performance in elite rowers. *Aust J Sci Med Sport.* 1991; 23: 66-69.

20. Mc Naughton LR. Sodium bicarbonate ingestion and its effects on anaerobic exercise of various duration. *J Sports Sci.* 1992; 10: 425-435.
21. Mc Naughton, L.R. Bicarbonate ingestion: effects of dosage on 60s cycle ergometry. *J Sports Sci.* 1992; 10: 415-423
22. Siegler JC, Gleadall-Siddall DO. Sodium Bicarbonate Ingestion and Repeated Swim Sprint Performance. *J Strength Cond Res.* 2010; 24:3015-3111.
23. Gao J, Costill DL, Horswill CA, Park, SH. Sodium bicarbonate ingestion improves performance in interval swimming. *Eur J Appl Physiol Occ Physiol.* 1988, 58: 171-174.
24. Kilding AE, Overton C, Gleave J. (2012) Effects of caffeine, sodium bicarbonate, and their combined ingestion on high-intensity cycling performance. *Int J Sport Nutr Exerc Metab.* 22(3):175-83.
25. Pruscino CL, Ross ML, Gregory JR, Savage B, Flanagan TR. (2008). Effects of sodium bicarbonate, caffeine, and their combination on repeated 200-m freestyle performance. *Int J Sport Nutr Exerc Metab.* 18(2):116-30.
26. Danaher J, Gerber T, Wellard RM, Stathis CG. (2014). The effect of β -alanine and NaHCO₃ co-ingestion on buffering capacity and exercise performance with high-intensity exercise in healthy males. *Eur J Appl Physiol.* 114(8):1715-24. doi: 10.1007/s00421-014-2895-9.
27. Saunders B, Sale C, Harris RC, Sunderland C. (2014b). Effect of sodium bicarbonate and Beta-alanine on repeated sprints during intermittent exercise performed in hypoxia. *Int J Sport Nutr Exerc Metab.* 24(2):196-205. doi: 10.1123/ijsnem.2013-0102.
28. Saunders B, Sale C, Harris RC, Sunderland C. (2014a). Sodium bicarbonate and high-intensity-cycling capacity: variability in responses. *Int J Sports Physiol Perform.* 2014; 9:627-632.

29. Christensen PM, Petersen MH, Friis SN, Bangsbo J. (2014). Caffeine, but not bicarbonate, improves 6 min maximal performance in elite rowers. *Appl Physiol Nutr Metab.* 39(9):1058-63. doi: 10.1139/apnm-2013-0577.
30. Egger F, Meyer T, Such U, Hecksteden A. (2014). Effects of Sodium Bicarbonate on High-Intensity Endurance Performance in Cyclists: A Double-Blind, Randomized Cross-Over Trial. *PLoS One.* 2014 Dec 10;9(12):e114729. doi: 10.1371/journal.pone.0114729.
31. Higgins MF, James RS, Price MJ. The effects of sodium bicarbonate (NaHCO₃) ingestion on high intensity cycling capacity. *J Sports Sci.* 2013; 31: 972-981.
32. Marriott M, Krstrup P, Mohr M. (2015). Ergogenic effects of caffeine and sodium bicarbonate supplementation on intermittent exercise performance preceded by intense arm cranking exercise. *J Int Soc Sports Nutr.* Feb 27;12:13. doi: 10.1186/s12970-015-0075-x.
33. Krstrup P, Ermidis G, Mohr M. (2015). Sodium bicarbonate intake improves high-intensity intermittent exercise performance in trained young men. *J Int Soc Sports Nutr.* 2015 Jun 4;12:25. doi: 10.1186/s12970-015-0087-6.
34. Mueller SM¹, Gehrig SM, Frese S, Wagner CA, Boutellier U, Toigo M. (2013). Multiday acute sodium bicarbonate intake improves endurance capacity and reduces acidosis in men. *J Int Soc Sports Nutr.* 10(1):16. doi: 10.1186/1550-2783-10-16.
35. Price M, Moss P, Rance S (2003). Effects of Sodium Bicarbonate Ingestion on Prolonged Intermittent Exercise. *J Med Sci Sports Exerc.* 2003; 35: 1303-1308.
36. Lavender G, Bird SR. Effect of sodium bicarbonate ingestion upon repeated sprints. *Br J Sports Med.* 1989; 23: 1-6.
37. Mc Naughton, LR, Siegler JC, Keatley S, Hillman A. The effects of sodium bicarbonate ingestion on maximal tethered treadmill running, *Gazzetta Medica Italiana - Archivio per le Scienze Mediche.* 2011; 170: 33-39.

38. Bishop D, Spencer M. Determinants of repeated-sprint ability in well-trained team-sport athletes and endurance trained athletes. *J Sports Med and Phys Fit.* 2004; 44: 1–7.
39. Mc Naughton LR, Ford S, Newbold C. The effects of sodium bicarbonate ingestion on high intensity exercise in moderately trained women. *J Str Cond Res.* 1997; 11: 98-102.
40. Mc Naughton LR, Curtin R, Perry D, Turner B, Showell C. Bicarbonate loading and the effects on anaerobic work and power output during cycle ergometer performance. *J Sports Sci.* 1991; 9: 151-160.
41. Gaitanos GC, Nevill ME, Brooks S, Williams C. Repeated bouts of sprint running after induced alkalosis. *J Sports Sci.* 1990; 9: 335-370.
42. Grafen A, Hails R (2002). *Modern Statistics for the Life Sciences.* Oxford University Press, Oxford, UK.
43. Cohen J (1988). *Statistical power analysis for the behavioural sciences.* 2nd Edition. Erlbaum, Hillsdale, New Jersey, USA.
44. McNaughton L, Dalton B., Palmer G. Sodium bicarbonate can be used as an ergogenic aid in high-intensity, competitive cycle ergometry of 1 h duration, *Eur J Appl Physiol Occ Physiol.* 1999, 80: 64-64

Table 1. Individual responses to ingestion of 300mg·kg⁻¹ sodium bicarbonate

Participant	Time (mins)												
	0	10	20	30	40	50	60	65	70	75	80	85	90
1	7.409	7.433	7.441	7.465	7.485	7.49	7.495	7.495	7.489	7.496	7.499	7.528	7.520
2	7.410	7.431	7.453	7.476	7.485	7.473	7.467	7.483	7.506	7.517	7.512	7.508	7.498
3	7.390	7.429	7.452	7.459	7.475	7.475	7.497	7.485	7.491	7.518	7.482	7.513	7.489
4	7.429	7.472	7.462	7.443	7.472	7.447	7.495	7.517	7.509	7.501	7.499	7.478	7.476
5	7.431	7.45	7.457	7.477	7.468	7.47	7.472	7.484	7.484	7.468	7.465	7.463	7.448
6	7.388	7.416	7.448	7.429	7.466	7.468	7.457	7.460	7.465	7.477	7.484	7.450	7.450
7	7.414	7.482	7.480	7.474	7.473	7.468	7.461	7.459	7.454	7.451	7.447	7.445	7.442
8	7.411	7.454	7.457	7.460	7.481	7.477	7.494	7.518	7.494	7.513	7.495	7.503	7.498
9	7.439	7.459	7.471	7.486	7.468	7.456	7.476	7.465	7.472	7.471	7.457	7.465	7.469
10	7.454	7.474	7.494	7.488	7.508	7.48	7.500	7.501	7.496	7.483	7.486	7.473	7.472
11	7.42	7.434	7.454	7.428	7.439	7.469	7.453	7.451	7.471	7.449	7.455	7.44	7.471
12	7.468	7.452	7.516	7.521	7.507	7.516	7.510	7.459	7.462	7.466	7.443	7.445	7.44
13	7.417	7.423	7.441	7.445	7.45	7.44	7.457	7.465	7.462	7.463	7.451	7.446	7.441
14	7.447	7.407	7.448	7.482	7.457	7.486	7.48	7.477	7.486	7.465	7.485	7.461	7.49
15	7.403	7.406	7.456	7.438	7.46	7.464	7.444	7.462	7.481	7.487	7.448	7.452	7.441
16	7.431	7.444	7.461	7.447	7.46	7.488	7.477	7.497	7.487	7.509	7.474	7.49	7.487
17	7.419	7.426	7.431	7.446	7.454	7.467	7.464	7.483	7.477	7.49	7.48	7.481	7.477
Mean	7.422	7.441	7.460	7.463	7.471	7.473	7.476	7.480	7.482	7.484	7.474	7.473	7.471
SD	0.021	0.023	0.021	0.025	0.018	0.017	0.019	0.021	0.016	0.023	0.021	0.027	0.024

Note: Peak pH is illustrated in bold font.

Figure 1. Mean (\pm SD) changes in pH following sodium bicarbonate ingestion (A) and individual participant time to peak pH frequency (B). (*) Denotes a significant increase in pH from the previous time point ($p < 0.01$). (●) Denotes a significant increase in pH from the 20 min sample $p \leq 0.01$. (Δ) Denotes a significant decrease in pH from the 75 min sample ($p < 0.05$).