



LEEDS
BECKETT
UNIVERSITY

Citation:

Phibbs, P and Jones, B and Roe, G and Read, D and Darrall-Jones, J and Weakley, J and Till, K (2017) We know they train, but what do they do? Implications for coaches working with adolescent rugby union players. *International Journal of Sports Science and Coaching*, 12 (2). pp. 175-182. ISSN 1747-9541 DOI: <https://doi.org/10.1177/1747954117694734>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3177/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

**We know they train, but what do they do? Implications for coaches working
with adolescent rugby union players**

Running Title: Session Training Loads in Adolescent Rugby Union

Padraic J Phibbs^{1,2}, Ben Jones^{1,2}, Gregory AB Roe^{1,2}, Dale B Read^{1,2}, Joshua
Darrall-Jones^{1,2}, Jonathon JS Weakley^{1,2} and Kevin Till^{1,2}

¹Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, UK

²Yorkshire Carnegie Rugby Club, Leeds, UK

Corresponding Author:

Padraic J. Phibbs, Institute for Sport, Physical Activity and Leisure,
G07 Cavendish, Headingley Campus, Leeds Beckett University, LS63QS, UK.

Email: P.Phibbs@leedsbeckett.ac.uk

We know they train, but what do they do? Implications for coaches working with adolescent rugby union players

Abstract

Limited information is available regarding the training loads (TLs) of adolescent rugby union players. One-hundred and seventy male players (age 16.1 ± 1.0 years) were recruited from ten teams representing two age categories (under-16 and under-18) and three playing standards (school, club and academy). Global positioning systems, accelerometers, heart rate and session-rating of perceived exertion (s-RPE) methods were used to quantify mean session TLs. Session demands differed between age categories and playing standards. Under-18 academy players were exposed to the highest session TLs in terms of s-RPE (236 ± 42 AU), total distance (4176 ± 433 m), high speed running (1270 ± 288 m) and PlayerLoad™ (424 ± 56 AU). Schools players had the lowest session TLs in both respective age categories. Training loads and intensities increased with age and playing standard. Individual monitoring of TL is key to enable coaches to maximise player development and minimise injury risk.

Keywords

Training load, youth, football, demands

Introduction

Rugby union is a collision team sport which elicits a variety of physiological responses during match play, due to the repeated high-intensity effort and collision nature of the sport.¹ The demands of match play have been frequently investigated,²⁻⁴ yet little information is available on the training loads which are a major factor in the development of rugby players.⁵ Despite England having the highest number of participants in rugby union,⁶ to date there is limited knowledge of the physical demands of English adolescent rugby union training. A recent study found that junior rugby union players were heavier (20%), stronger (50%), and faster (7%) than their counterparts 13 years previous.⁷ Although the authors suggested that adaptation to the changing demands of the game and enhanced training methodologies have resulted in these physical changes, little is known about either suggestion in this population. Recently, training demands have been shown to relate to injury,⁸⁻¹⁰ performance,¹¹ and physical development,⁵ thus this is imperative information for the coach to make evidence-based decisions on how they structure training sessions for developing athletes.

Training load (TL) is the product of training volume and intensity, and can be categorised as either internal or external.¹² The internal load is the relative physiological and psychological stress imposed on an individual by an activity, whereas the external load is the total work completed by the individual independent of internal factors.¹² Quantification of TL is an emerging research area and common practice in elite and professional sport, but limited information has been published in adolescent team sport athletes due to the protection of the data by sporting clubs and organisations.¹³ For optimal athletic performance a balance of training stimuli and recovery processes are required, which can be further complicated during adolescence due to the associated physiological and psychological changes.¹⁴

Monitoring TL in adolescent athletes is critical to ensure that positive training effects are maximised and negative consequences such as injury, illness and/or overtraining are avoided.⁵

To date, the quantification of TL in adolescent rugby union players is limited to a single research group in Australia.¹⁵⁻¹⁷ The research group found that although weekly training volumes were high, individual session intensities were significantly lower than those observed in competitive match play.¹⁷ A major limitation of these studies is that there were no within-group comparisons, with respect to age categories, in a cohort of 14-18 year olds. Furthermore, previous research in adolescent soccer players identified a significant increase in both training volume and intensity between the ages of 14 and 18 years.¹⁸ The differences in training frequency, volume and intensity between age groups in junior rugby union players are likely to differ but are yet to be determined.

Adolescent rugby union players may participate with teams in multiple age categories and playing standards simultaneously. As both excessive and insufficient TL during adolescence may impede optimal athletic development, understanding the specific TLs undertaken by adolescent rugby union players is clearly important for their athletic performance, injury prevention, playing progression and general wellbeing.^{14,16,19} A greater understanding of the demands of individual training sessions by coaches will help to optimise development, performance, and progression, whilst simultaneously reducing the participants' likelihood of exposure to negative training effects.⁵ Therefore the aim of the present study was to compare in-season field-based session TLs of adolescent rugby union players by age category and playing standard.

Methods

Participants

Adolescent rugby union players (n=170) from ten rugby union teams across three playing standards (independent school, amateur club and professional regional academy) at under-16 and under-18 age groups were recruited for this prospective study. Players were categorised into six independent groups; U16 schools (U16-S; n=31), U18 schools (U18-S; n=39), U16 clubs (U16-C; n=36), U18 clubs (U18-C; n=30), U16 regional academy (U16-A; n=18) and U18 regional academy (U18-A; n=16). Players may participate with teams at multiple playing standards or age categories simultaneously, thus for the purpose of this study participants were grouped into the highest age category or playing standard that they participated in at the time of data collection. Table 1 shows the participant characteristics (i.e. age, stature and body mass) and weekly training frequencies of each group. All players and parents provided informed written consent prior to participation. Ethics approval was granted by the Leeds Beckett University research ethics committee.

****Insert Table 1 Near Here****

Design of Study

In a prospective cohort design, each team was monitored during one complete in-season training week to quantify mean session TLs. Training practices were not altered or interfered with by the researchers at any time. All data were collected mid-season for each respective squad (between October 2014 and January 2015) to control for potential differences in TL due to the stage of season. The week was described as a “typical” training week (i.e. training frequency and intended intensity) by the coaches who were leading the sessions. Each training week was selected to

provide the most representative microcycle for the respective teams in-season phase, in preparation for a single home competitive fixture.

During all training sessions, players wore a 10 Hz global positioning system (GPS) device (Optimeye S5, Catapult Innovations, Victoria, Australia) positioned on the upper back between the scapulae in a tight fitting custom-made vest. The mean \pm standard deviation (*SD*) number of satellites connected was 14.4 ± 0.3 and horizontal dilution of precision was 0.82 ± 0.14 during data collection. Heart rate (HR) was captured using a portable HR monitor (T31c, Polar Electro, Kempele, Finland), which was fitted around the torso, level with the xiphoid process. All players wore the same GPS and HR monitoring devices for each session and the reliability and validity of the devices used in this study have been previously reported.²⁰⁻²² Session-rating of perceived exertion (s-RPE) was also recorded post-training for all players.²³

Procedures

Perceptual intensity of training was quantified using a modified Borg Category Ratio-10 RPE scale.²³ All players were familiarised with the scale prior to commencement of the study. Approximately 30 minutes post-training, participants recorded their RPE, to minimise bias from the most recent phase of exercise. Recordings were taken non-verbally with each participant on their own and also blinded from previous scores to control for external influences. The RPE score was then multiplied by session duration (min) to provide a s-RPE value. Physiological intensity was quantified using mean exercise HR (HR_{mean}) to provide an objective comparison for internal TL.

External TL was monitored using GPS and tri-axial accelerometer measures (total distance [TD], high speed running [HSR] distance, and PlayerLoad™ [PL]). The

threshold for HSR was defined as distance covered $>12 \text{ km}\cdot\text{h}^{-1}$ due to previously suggested population-specific movement classifications.¹⁷ Tri-axial accelerometers within the GPS device provides a measure of combined anteroposterior, mediolateral and vertical accelerations (i.e. PL) to account for the additional non-locomotor activity demands of rugby union training. Accelerometer and GPS variables standardised for time were calculated to quantify external intensity (relative distance [$\text{m}\cdot\text{min}^{-1}$], relative HSR distance [$\text{HSR}\cdot\text{min}^{-1}$], and relative PL [$\text{PL}\cdot\text{min}^{-1}$]). Relative measures were calculated by dividing the total value by the duration of the session (min). Movement demands of training were also quantified using population-specific absolute velocity bands categorised as stationary ($0\text{-}1 \text{ km}\cdot\text{h}^{-1}$), low ($1\text{-}7 \text{ km}\cdot\text{h}^{-1}$), moderate ($7\text{-}12 \text{ km}\cdot\text{h}^{-1}$), high ($12\text{-}21 \text{ km}\cdot\text{h}^{-1}$) and very high ($>21 \text{ km}\cdot\text{h}^{-1}$) speed running, as per previous adolescent rugby research.¹⁷

Following each session, all GPS and HR data recorded were downloaded to the manufacturer's software (Sprint 5.1.4, Catapult Innovations, Victoria, Australia). Once downloaded, all data were cropped so that only on-field activity for the recorded session time were included. Data were then exported to a customised Excel spreadsheet (Microsoft, Redmond, USA) for calculation of the selected variables.

Statistical Analyses

Mean data were calculated for each participant from their respective weekly sessions to control for multiple and uneven observations.²⁴ After verification of normality, to assess the magnitude of between-group differences, Cohen's *d* effect sizes (*ES*) were calculated with threshold values set at <0.2 (*trivial*), $0.2\text{-}0.6$ (*small*), $0.6\text{-}1.2$ (*moderate*), $1.2\text{-}2.0$ (*large*) and ≥ 2.0 (*very large*).^{25,26} Magnitude based-inferences were used to assess for practical significance.²⁶ The threshold for a change

to be considered practically important (the smallest worthwhile change; SWC) was set at 0.2 x between subject standard deviation (*SD*), based on Cohen's *d ES* principle. The probability that the magnitude of change was greater than the SWC was rated as 25-75%, *possibly*; 75-95%, *likely*; 95-99.5%, *very likely*; >99.5%, *most likely*.²⁶ Where the 90% Confidence Interval (CI) crossed both the upper and lower boundaries of the SWC ($ES \pm 0.2$), the magnitude of change was described as unclear.²⁶ All normally distributed data are presented as mean \pm *SD*. Non-normally distributed data are presented as descriptive statistics only (i.e. median and interquartile range), and no further analyses were performed.

Results

Table 2 presents the training load and intensity differences between age categories and Table 3 presents the differences between playing standards.

****Insert Table 2 Near Here****

****Insert Table 3 Near Here****

Within the school standard, U18-S were *likely* greater than U16-S for TD (*small*), *very likely* greater for training duration (*moderate*) and RPE (*moderate*), and *most likely* greater for s-RPE (*moderate*). However, U18-S were *likely* lower than U16-S for relative PL (*small*), *very likely* lower for relative HSR (*moderate*), and *most likely* lower for HR_{mean} (*large*).

Within club standard, U18-C were *very likely* greater than U16-C for training duration (*moderate*), but *likely* lower than U16-C for RPE (*small*), and *most likely* lower for total and relative HSR (*large to very large*).

Within the academy standard, U18-A were *likely* greater than U16-A for s-RPE (*small*), and *most likely* greater for training duration (*very large*), total and relative

distance (*large to very large*), total and relative HSR (*large to very large*) and total PL (*large*). Although, U18-A were *likely* lower than U16-A for HR_{mean} (*moderate*) and RPE (*moderate*).

Within the U16 age category, U16-C were *likely* greater than U16-S for total HSR (*moderate*) and *most likely* greater for training duration (*large*), s-RPE (*large*), TD (*large*), total PL (*large*) and RPE (*large*). U16-A were *likely* greater than U16-S for HR_{mean} (*moderate*), TD (*small*), and relative distance (*small*) and *most likely* greater for s-RPE (*large*), total PL (*moderate*), RPE (*very large*) and relative PL (*large*). However, total and relative HSR were both *likely* lower (*moderate*) in the U16-A compared to the U16-S. The U16-A group were *likely* greater than U16-C for HR_{mean} (*small*) and relative distance (*small*), *very likely* greater for RPE (*moderate*), and *most likely* greater for relative PL (*large*). Although, U16-A were *likely* lower than U16-C for total PL (*moderate*), *very likely* lower for relative HSR (*moderate*) and *most likely* lower for training duration (*very large*), TD (*large*), and total HSR (*large*).

Within the U18 age category, U18-C were *likely* greater than U18-S for RPE (*small*), and *most likely* greater for training duration (*large*), s-RPE (*moderate*), HR_{mean}, TD, and total PL (all *large*). However, U18-C were *likely* lower than U18-S for total HSR (*small*) and *most likely* lower for relative HSR (*large*). U18-A were *likely* greater than U18-S for training duration (*moderate*) and *most likely* greater for all other measured variables (*large to very large*). U18-A were *likely* greater than U18-C for TD (*moderate ES*), *very likely* greater for total PL (*moderate*) and RPE (*moderate*), and *most likely* greater for total and relative HSR, relative distance and relative PL, with *most likely* lower training durations (all *very large*).

Table 4 presents the movement demands of training for each group. All six groups covered the highest distance at low speed and the lowest distance at very high

speed running (VHSR). Median percentage time spent stationary was highest for U16-S and U18-A groups with the lowest observed in U18-C. Median percentage time spent HSR was highest for U18-A and lowest for U18-C, with median percentage time spent VHSR was 0% for all groups. Median total distance covered at VHSR was also highest for U18-A and lowest for U18-C.

****Insert Table 4 Near Here****

Discussion

This is the first study to quantify and compare the physical outputs of adolescent rugby union players during training by both playing standard and age category. The U18-A group (i.e. highest age and standard) had the greatest s-RPE, TD, HSR and PL while the U16-S group (i.e. lowest age and standard) had the lowest s-RPE, TD and PL of all groups. There was an increase in session durations (i.e. volume) between U16 and U18 age categories in all playing standards. When standardised for time, training intensities (i.e. RPE, $\text{m}\cdot\text{min}^{-1}$ and $\text{PL}\cdot\text{min}^{-1}$) were highest in academy groups and lowest in school groups for both respective age categories. The findings of this study suggest that the demands of field-based training sessions increase with age and playing standard in adolescent rugby union players.

Overall, subjective and objective measures of session TL were highest in the U18-A group, which is the highest playing standard and therefore may be expected to elicit the highest training demands. The U16-S and U18-S groups had the lowest subjective and objective measures of session TL in each respective age category. However, when standardised for time, objective measures of session intensity (i.e. $\text{m}\cdot\text{min}^{-1}$ and $\text{PL}\cdot\text{min}^{-1}$) were similar between school and club groups at both age categories suggesting that the differences in TL are predominantly due to increased

session durations in the club groups. Interestingly, the subjective measure of intensity (i.e. RPE) did not follow the trend of the objective measures, but as RPE is an internal load measure it is influenced by individual psychological and physiological characteristics. At the academy standard there was an increase in session durations between U16 and U18 age categories but also an increase in the objective measures of intensity. Club groups had the lowest weekly training frequencies in both age categories, which may explain the perceived requirement for these groups to have longer session durations. Coaches should aim to prepare players for the specific demands of match-play and not simply increase training volumes due to limited training availability. The intensity of the session may be more important for both physical development and injury prevention,^{10,17} thus coaches should look to prescribe specific sessions to elicit these positive outcomes. High training volumes may become problematic for players participating with multiple teams simultaneously resulting in excessive weekly workloads.⁵

Training session durations increased with age in all three playing standards. This increase in training volume in association with age has also been observed in other adolescent team sports.¹⁸ Optimal TL has been suggested to maximise athletic development whilst minimizing risk of illness and injury when structuring age appropriate training.²⁷ Adolescent athletes have been suggested to be at the greatest risk of microtrauma during times of peak physical growth.²⁸ This may partially explain the lesser training volumes in the U16 age category in the present study. Fourteen year old team sport athletes are at a different stage of athletic development to 18 year olds, where there would be a greater emphasis on competition and competition-specific training for the older age category.²⁹ The findings of this study suggest that

total TLs were higher in U18 players due to their longer field-based session durations and increased weekly training frequencies.

Training durations in the present study were generally lower than previously reported in adolescent rugby union training (range; 58-93 min),¹⁵ while unfortunately the use of an alternative scale to measure s-RPE in previous research prevents direct comparisons for subjective TLs. However, HR_{mean} measures reported in this study are similar to those previously reported (range; 136-141 b·min⁻¹) suggesting a similar mean physiological intensity of training.¹⁵ The total distances covered during training sessions in the current study are also comparable to previously reported values (range; 2208-3576 m).¹⁵ However, compared to the previously reported training data (range; 38-50 m·min⁻¹), the relative distances covered in the current study are greater for all six groups.¹⁵ These findings suggest that although session durations were lower in the present study, session intensities were higher than previously reported during training in a similar population resulting in similar external and internal loads.

Of note, the median percent time spent undertaking HSR and VHSR during training was 3-7% and 0%, respectively. Interestingly, at school and club standards the U16 players were exposed to greater HSR demands than the U18 players, however this finding is reversed in the academy standard. This may be attributed to the academy training programmes approach to improve high intensity running ability alongside increases in body mass. The use of 12 km·h⁻¹ as a threshold for HSR may be considered conservative, however it was selected as a population specific absolute value based on the findings of previous adolescent rugby research.¹⁷ The individual design of training sessions by the coaches will likely influence the HSR demands of field-based sessions, as evident in the current study with HSR distances substantially different between groups. The selection of various training drills and games in

sessions allows manipulation of HSR exposure in training by altering pitch dimensions and player numbers.

Similar VHSR values have been reported in a comparison of training and match movement demands in a similar population, where the median time spent above the same threshold was 0% for training and 1.3% for matches.¹⁷ It should be further noted that the previous research investigating the differences in running demands of training in comparison to match play in adolescent rugby union employed two different analysis systems for each condition (i.e. GPS for training and computer-based tracking software for matches), due to restrictions on the use of microtechnology during match play at the time. Future studies are also required to examine the differences in HSR relative to the individual's maximum sprinting speed (V_{max}) to further explore the relative running demands in adolescent rugby union players. The absolute values in the current study may overestimate HSR and VHSR demands for faster players and underestimate demands for slower players.³⁰

The U18-A group covered the greatest HSR distance, although without relative speed comparisons it is difficult to distinguish whether this is due to enhanced session design or simply because they are faster athletes. However, adolescent rugby union players across all age categories and playing standards do not appear to be exposed to adequate VHSR in training that have been previously reported in match play.¹⁷ A recent study showed that increases in momentum were greater than increases in V_{max} with age for regional academy rugby union players, with greater increases in body mass than speed.³¹ The development of V_{max} may therefore be inhibited due to the lack of exposure to VHSR in training for young rugby union players. Exposure to VHSR may be easily incorporated into a field-based session following an adequate warm up with the inclusion of one or two 30-40 m maximal sprint efforts. However,

caution must be taken by coaches not to overprescribe VHSR as excessive distances at these velocities have also been linked to injury risk.³²

As this study is the first to report measurements of PL in junior rugby union training no direct comparisons can be made for this variable. The inclusion of PL as a measure of global external load may provide valuable information on the additional non-locomotor demands of rugby union training when used in combination with movement demand variables.^{21,33} As total external load in rugby union consists of much more than simply movement-based demands, the use of PL and its derivatives provide an insight of the global external load of training inclusive of additional rugby-specific activities such as jumping, tackles, mauls, rucks, and scrums.^{1,4,22} Due to the limited involvement of physical contact and collisions in training compared to match play, low speed activity (LSA) or HSR demands may be increased to compensate for the difference in physiological intensity between training and matches to produce similar loads.

In conclusion, the current study provides important information for coaches working in adolescent rugby union in relation to the TLs of in-season field-based training sessions, specific to each respective age category and playing standard. School and amateur club coaches may want to adopt training practices similar to academy sessions, by focussing on intensity rather than volume, to maximise player development. However, coaches must remain cognisant that if session intensities are increased, training volumes may need to be decreased to avoid excessive accumulated TLs. Due to the limited exposure to full collision-based activity in training and its resultant effect on the reduced physiological intensity of sessions compared to match-play, coaches should consider strategies and behaviours to compensate for this deficit and maximise player involvement, where increasing LSA or HSR may be

beneficial. Including small exposures to VHSR during training may assist to improve sprint performance. Finally, as many adolescent rugby players participate with multiple teams concurrently, TLs should be monitored for players who may be undertaking additional training with teams away from the coaches' supervised environment to help inform appropriate prescription of training to maximise performance and protect the player from injury.

Acknowledgements

The authors would like to thank all of the coaches, parents and players who were involved in the project. This research was part funded by Leeds Rugby as part of the Carnegie Adolescent Rugby Research (CARR) project.

References

1. Duthie G, Pyne D, and Hooper S. Applied physiology and game analysis of rugby union. *Sports Med* 2003; 33: 973-991.
2. Quarrie, KL, Hopkins WG, Anthony MJ, et al. Positional demands of international rugby union: evaluation of player actions and movements. *J Sci Med Sport* 2013; 16: 353-359.
3. Cahill N, Lamb K, Worsfold P, et al. The movement characteristics of English Premiership rugby union players. *J Sport Sci* 2013; 31: 229-237.
4. Roberts SP, Trewartha G, Higgitt RJ, et al. The physical demands of elite English rugby union. *J Sport Sci* 2008; 26: 825-833.
5. Gabbett TJ, Whyte DG, Hartwig TB, et al. The relationship between workloads, physical performance, injury and illness in adolescent male football players. *Sports Med* 2014; 44: 989-1003.

6. Freitag A, Kirkwood G and Pollock AM. Rugby injury surveillance and prevention programmes: are they effective? *British Medical Journal* 2015; 350: h1587.
7. Lombard WP, Durandt JJ, Masimla H, et al. Changes in body size and physical characteristics of South African under-20 rugby union players over a 13-year period. *J Strength Cond Res* 2015; 29: 980-988.
8. Rogalski B, Dawson B, Heasman J, et al. Training and game loads and injury risk in elite Australian footballers. *J Sci Med Sport* 2012; 16: 499-503.
9. Cross MJ, Williams S, Trewartha G, et al. The Influence of In-Season Training Loads on Injury Risk in Professional Rugby Union. *Int J Sports Physiol Perform*. Epub ahead of print 26 Aug 2015. DOI: 10.1123/ijsp.2015-0187.
10. Gabbett TJ. The training-injury paradox; should athletes be training smarter and harder? *Br J Sports Med* 2016; 50: 273-280.
11. Aughey RJ, Elias GP, Esmaeili A, et al. Does the recent internal load and strain on players affect match outcome in elite Australian football? *J Sci Med Sport* 2016; 19: 182-186.
12. Borresen J and Lambert MI. The quantification of training load, the training response and the effect on performance. *Sports Med* 2009; 39: 779-795.
13. Halson SL. Monitoring training load to understand fatigue in athletes. *Sports Med* 2014; 44 Suppl 2: S139-S147.
14. Bergeron MF, Mountjoy M, Armstrong N, et al. International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med* 2015; 49: 843-851.

15. Hartwig TB, Naughton G, and Searl J. Defining the volume and intensity of sport participation in adolescent rugby union players. *Int J Sports Physiol Perform* 2008; 3: 94-106.
16. Hartwig TB, Naughton G, and Searl J. Load, stress, and recovery in adolescent rugby union players during a competitive season. *J Sport Sci* 2009; 27: 1087-1094.
17. Hartwig TB, Naughton G, and Searl J. Motion analyses of adolescent rugby union players: a comparison of training and game demands. *J Strength Cond Res* 2011; 25: 966-972.
18. Wrigley R, Drust B, Stratton G, et al. Quantification of the typical weekly in-season training load in elite junior soccer players. *J Sport Sci* 2012; 30: 1573-1580.
19. Smith DJ. A framework for understanding the training process leading to elite performance. *Sports Med* 2003; 33: 1103-1126.
20. Varley MC, Fairweather IH, and Aughey RJ, Validity and reliability of GPS for measuring instantaneous velocity during acceleration, deceleration, and constant motion. *J Sport Sci* 2012; 30: 121-127.
21. Boyd LJ, Ball K, and Aughey RJ, Quantifying external load in Australian football matches and training using accelerometers. *Int J Sports Physiol Perform* 2013; 8: 44-51.
22. Gabbett TJ. Relationship between Accelerometer Load, Collisions, and Repeated High-Intensity Effort Activity in Rugby League Players. *J Strength Cond Res* 2015; 29: 3424-3431.
23. Foster C, Florhaug JA, Franklin J, et al. A new approach to monitoring exercise training. *J Strength Cond Res* 2001; 15: 109-115.

24. Wilkinson M, and Akenhead, R. Violation of statistical assumptions in a recent publication? *Int J Sport Med* 2013; 34: 281.
25. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd Edition. NJ: Hillsdale, 1988.
26. Hopkins WG, Marshall, SW, Batterham, AM, et al. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 2009; 41: 3-13.
27. Naughton G, Farpour-Lambert NJ, Carlson J, et al. Physiological issues surrounding the performance of adolescent athletes. *Sports Med* 2000; 30: 309-325.
28. van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ, et al. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sport Med* 2014; 35: 351-355.
29. Balyi I, and Hamilton A. Long-term athlete development: Trainability in childhood and adolescence. *Olympic Coach* 2004; 16: 4-9.
30. Gabbett TJ. The use of relative speed zones increases the high-speed running performed in team sport match-play. *J Strength Cond Res* 2015; 29: 3353-3359.
31. Darrall-Jones JD, Jones B, and Till K. Anthropometric and Physical Profiles of English Academy Rugby Union Players. *J Strength Cond Res* 2015; 29: 2086-2096.
32. Gabbett TJ, and Ullah S. Relationship between running loads and soft-tissue injury in elite team sport athletes. *J Strength Cond Res* 2012; 26: 953-960.

33. Scott BR, Lockie RG, Knight TJ, et al. A comparison of methods to quantify the in-season training load of professional soccer players. *Int J Sports Physiol Perform* 2013; 8: 195-202.

Table 1. Participant Characteristics.

	U16 Schools (n=31)	U16 Club (n=36)	U16 Academy (n=18)	U18 Schools (n=39)	U18 Club (n=30)	U18 Academy (n=16)
Age (years)	15.3 ± 0.4	15.3 ± 0.5	15.3 ± 0.7	16.9 ± 0.6	16.4 ± 0.5	17.1 ± 0.7
Stature (cm)	174.1 ± 7.0	177.9 ± 6.0	177.4 ± 5.2	180.2 ± 6.2	178.8 ± 6.2	182.7 ± 7.2
Mass (kg)	70.7 ± 10.1	74.2 ± 11.0	79.8 ± 13.7	79.2 ± 11.8	80.4 ± 12.0	87.7 ± 10.4
Weekly Training Frequency	2 ± 0	1 ± 0	1 ± 0	3 ± 0	1 ± 0	2 ± 0

Data presented as mean ± SD.

Table 2. Training load and intensity differences between age categories.

	School Standard		Club Standard		Academy Standard	
	U16-S (n=31)	U18-S (n=39)	U16-C (n=36)	U18-C (n=30)	U16-A (n=18)	U18-A (n=16)
Duration (min)	50.1 ± 6.6 0.7 [0.3, 1.1]; Very Likely ↑	56.8 ± 11.9	63.9 ± 9.7 0.7 [0.3, 1.1]; Very Likely ↑	70.3 ± 8.8	48.3 ± 5.1 3.5 [2.9, 4.0]; Most Likely ↑	62.0 ± 0.0
s-RPE (AU)	123 ± 39 1.00 [0.6, 1.3]; Most Likely ↑	168 ± 55	231 ± 73 0.0 [-0.4, 0.4]; Unclear	230 ± 67	211 ± 50 0.5 [-0.1, 1.1]; Likely ↑	236 ± 42
Mean HR (b·min⁻¹)	145 ± 8 -1.2 [-0.8, -1.6]; Most Likely ↓	134 ± 9	145 ± 11 0.3 [-0.1, 0.7]; Possibly ↑	148 ± 14	151 ± 12 -0.6 [-1.1, 0.0]; Likely ↓	146 ± 7
Total Distance (m)	2672 ± 456 0.5 [0.1, 0.9]; Likely ↑	2925 ± 467	3619 ± 664 0.4 [-0.1, 0.8]; Possibly ↑	3845 ± 577	2903 ± 434 2.9 [2.3, 3.4]; Most Likely ↑	4176 ± 433
Total HSR Distance (m)	751 ± 242 -0.3 [-0.7, 0.1]; Possibly ↓	678 ± 179	955 ± 256 -1.4 [-1.8, -1.0]; Most Likely ↓	597 ± 246	590 ± 219 2.6 [2.0, 3.2]; Most Likely ↑	1270 ± 288
Total PL (AU)	262 ± 41 0.2 [-0.2, 0.6]; Unclear	270 ± 42	354 ± 74 0.2 [-0.2, 0.6]; Possibly ↑	371 ± 75	316 ± 53 1.9 [1.4, 2.5]; Most Likely ↑	424 ± 56
RPE (AU)	2.5 ± 0.6 0.7 [0.3, 1.1]; Very Likely ↑	2.9 ± 0.5	3.6 ± 0.9 -0.4 [-0.8, 0.0]; Likely ↓	3.2 ± 0.7	4.3 ± 0.8 -0.7 [-1.3, -0.1]; Likely ↓	3.8 ± 0.7
Relative Distance (m·min⁻¹)	54.9 ± 12.3 0.0 [-0.4, 0.4]; Unclear	54.5 ± 10.4	56.8 ± 7.4 -0.3 [-0.7, 0.2]; Possibly ↓	54.9 ± 7.5	59.9 ± 5.7 1.2 [0.7, 1.8]; Most Likely ↑	68.1 ± 7.3
Relative HSR (m·min⁻¹)	15.4 ± 5.8 -0.6 [-1.0, -0.2]; Very Likely ↓	12.4 ± 3.6	15.0 ± 3.5 -2.0 [-2.4, -1.6]; Most Likely ↓	8.3 ± 3.1	12.1 ± 4.2 1.9 [1.3, 2.5]; Most Likely ↑	20.7 ± 4.7
Relative PL (AU·min⁻¹)	5.3 ± 1.0 -0.4 [-0.8, 0.0]; Likely ↓	5.0 ± 1.0	5.5 ± 0.8 -0.3 [-0.7, 0.1]; Possibly ↓	5.3 ± 0.9	6.5 ± 0.8 0.4 [-0.2, 1.0]; Possibly ↑	6.9 ± 0.9

Data presented as mean ± SD, Cohen's *d* effect size [90% confidence intervals]; magnitude-based inference.

Table 3. Training load and intensity differences between playing standards.

	U16-S (n=31)	U16-C (n=36)	U16-A (n=18)	U16-S vs U16-C	U16-S vs U16-A	U16-C vs U16-A
Duration (min)	50.1 ± 6.6	63.9 ± 9.7	48.3 ± 5.1	1.7 [1.3, 2.1]; Most Likely ↑	-0.3 [-0.7, 0.2]; Unclear	-2.0 [-2.4, -1.6]; Most Likely ↓
s-RPE (AU)	123 ± 39	231 ± 73	211 ± 50	1.8 [1.4, 2.2]; Most Likely ↑	1.9 [1.4, 2.4]; Most Likely ↑	-0.3 [-0.7, 0.1]; Possibly ↓
Mean HR (b·min⁻¹)	145 ± 8	145 ± 11	151 ± 12	0.1 [-0.3, 0.5]; Unclear	0.6 [0.1, 1.2]; Likely ↑	0.5 [0.0, 1.0]; Likely ↑
Total Distance (m)	2672 ± 456	3619 ± 664	2903 ± 434	1.6 [1.2, 2.0]; Most Likely ↑	0.5 [0.0, 1.0]; Likely ↑	-1.2 [-1.7, -0.8]; Most Likely ↓
Total HSR Distance (m)	751 ± 242	955 ± 256	590 ± 219	0.8 [0.4, 1.2]; Very Likely ↑	-0.7 [-1.2, -0.2]; Likely ↓	-1.5 [-2.0, -1.0]; Most Likely ↓
Total PL (AU)	262 ± 41	354 ± 74	316 ± 53	1.5 [1.1, 1.9]; Most Likely ↑	1.1 [0.6, 1.6]; Most Likely ↑	-0.6 [-1.0, -0.1]; Likely ↓
RPE (AU)	2.5 ± 0.6	3.6 ± 0.9	4.3 ± 0.8	1.4 [1.0, 1.8]; Most Likely ↑	2.5 [2.0, 3.0]; Most Likely ↑	0.8 [0.4, 1.3]; Very Likely ↑
Relative Distance (m·min⁻¹)	54.9 ± 12.3	56.8 ± 7.4	59.9 ± 5.7	0.2 [-0.2, 0.6]; Unclear	0.5 [0.1, 1.0]; Likely ↑	0.5 [0.0, 0.9]; Likely ↑
Relative HSR (m·min⁻¹)	15.4 ± 5.8	15.0 ± 3.5	12.1 ± 4.2	-0.1 [-0.5, 0.3]; Unclear	-0.6 [-1.1, -0.2]; Likely ↓	-0.7 [-1.2, -0.2]; Very Likely ↓
Relative PL (AU·min⁻¹)	5.3 ± 1.0	5.5 ± 0.8	6.5 ± 0.8	0.2 [-0.2, 0.6]; Possibly ↑	1.3 [0.8, 1.8]; Most Likely ↑	1.2 [0.7, 1.7]; Most Likely ↑
	U18-S (n=39)	U18-C (n=30)	U18-A (n=16)	U18-S vs U18-C	U18-S vs U18-A	U18-C vs U18-A
Duration (min)	56.8 ± 11.9	70.3 ± 8.8	62.0 ± 0.0	1.3 [0.9, 1.7]; Most Likely ↑	0.6 [0.2, 0.9]; Likely ↑	-1.4 [-1.8, -1.0]; Most Likely ↓
s-RPE (AU)	168 ± 55	230 ± 67	236 ± 42	1.0 [0.6, 1.4]; Most Likely ↑	1.3 [0.9, 1.8]; Most Likely ↑	0.1 [-0.4, 0.6]; Unclear
Mean HR (b·min⁻¹)	134 ± 9	148 ± 14	146 ± 7	1.2 [0.8, 1.6]; Most Likely ↑	1.4 [0.9, 1.9]; Most Likely ↑	-0.3 [-0.7, 0.2]; Possibly ↓
Total Distance (m)	2925 ± 467	3845 ± 577	4176 ± 433	1.7 [1.3, 2.1]; Most Likely ↑	2.7 [2.2, 3.2]; Most Likely ↑	0.6 [0.2, 1.1]; Likely ↑
Total HSR Distance (m)	678 ± 179	597 ± 246	1270 ± 288	-0.4 [-0.8, 0.0]; Likely ↓	2.4 [1.9, 3.0]; Most Likely ↑	2.5 [1.9, 3.0]; Most Likely ↑
Total PL (AU)	270 ± 42	371 ± 75	424 ± 56	1.6 [1.2, 2.0]; Most Likely ↑	3.0 [2.5, 3.6]; Most Likely ↑	0.8 [0.3, 1.3]; Very Likely ↑
RPE (AU)	2.9 ± 0.5	3.2 ± 0.7	3.8 ± 0.7	0.5 [0.1, 1.0]; Likely ↑	1.4 [0.9, 2.0]; Most Likely ↑	0.8 [0.3, 1.3]; Very Likely ↑
Relative Distance (m·min⁻¹)	54.5 ± 10.4	54.9 ± 7.5	68.1 ± 7.3	0.0 [-0.4, 0.4]; Unclear	1.5 [1.0, 2.0]; Most Likely ↑	1.8 [1.3, 2.3]; Most Likely ↑
Relative HSR (m·min⁻¹)	12.4 ± 3.6	8.3 ± 3.1	20.7 ± 4.7	-1.2 [-1.6, -0.8]; Most Likely ↓	2.0 [1.4, 2.5]; Most Likely ↑	3.0 [2.5, 3.6]; Most Likely ↑
Relative PL (AU·min⁻¹)	5.0 ± 1.0	5.3 ± 0.9	6.9 ± 0.9	0.3 [-0.1, 0.7]; Possibly ↑	2.0 [1.5, 2.5]; Most Likely ↑	1.7 [1.2, 2.2]; Most Likely ↑

Data presented as mean ± SD, Cohen's *d* effect size [90% confidence intervals]; magnitude-based inference.

Table 4. Total distance and percent time at various movement velocity bands during training.

	U16 Schools (n=31)	U16 Club (n=36)	U16 Academy (n=18)	U18 Schools (n=39)	U18 Club (n=30)	U18 Academy (n=16)
Low Speed Total Distance (0-7 km·h⁻¹)	1278 (301)	1915 (719)	1412 (172)	1499 (418)	2243 (298)	1527 (213)
Moderate Speed Total Distance (7-12 km·h⁻¹)	588 (204)	775 (308)	924 (187)	736 (253)	908 (227)	1385 (175)
High Speed Total Distance (12-21 km·h⁻¹)	692 (320)	872 (301)	579 (297)	601 (233)	579 (360)	1164 (398)
Very High Speed Total Distance (>21 km·h⁻¹)	32 (59)	83 (208)	36 (57)	51 (63)	24 (59)	140 (137)
Percent Time Stationary (0-1 km·h⁻¹)	37 (10)	31 (10)	33 (5)	32 (15)	24 (15)	37 (4)
Percent Time at Low Speed (1-7 km·h⁻¹)	47 (7)	55 (8)	49 (5)	47 (9)	65 (16)	41 (4)
Percent Time at Moderate Speed (7-12 km·h⁻¹)	11 (3)	7 (3)	11 (2)	9 (4)	8 (2)	14 (2)
Percent Time at High Speed (12-21 km·h⁻¹)	4 (2)	5 (2)	4 (2)	4 (2)	3 (1)	7 (3)
Percent Time at Very High Speed (>21 km·h⁻¹)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (1)

Data presented as median (interquartile range)