



LEEDS
BECKETT
UNIVERSITY

Citation:

Potter, CA (2017) Father experiences of sleeping problems in children with autism. *Advances in Autism*, 3 (2). pp. 100-111. ISSN 2056-3868 DOI: <https://doi.org/10.1108/AIA-06-2016-0017>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3397/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Father experiences of sleeping problems in children with autism

Journal:	<i>Advances in Autism</i>
Manuscript ID	AIA-06-2016-0017.R1
Manuscript Type:	Research Paper
Keywords:	children, fathers, sleeping problems, Health, employment, couple relationship

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Father experiences of sleeping problems in children with autism

Advances in Autism

Abstract

Purpose: This study aimed to explore the role of fathers in the management of sleeping problems in children with autism and their perspectives of the impact of these difficulties on family life. **Methodology:** Semi-structured interviews with 25 UK based fathers of children with autism were undertaken. **Findings:** Two thirds of fathers reported that their children experienced severe sleeping problems in the areas of bed-time resistance, sleep onset and night-time waking. Fathers were significantly involved in the management of these difficulties and reported a range of associated deleterious impacts on the family, including significant negative effects on paternal and maternal health, father employment, couple relationship and sibling experiences.

Research limitations: The interview sample cannot be said to be representative of all fathers of children with autism since the backgrounds of those taking part were relatively homogeneous in respect of ethnicity, marital status and level of education.

Practical implications: Improvements in effective, family-centred provision are urgently needed which employ a co-parenting, gender-differentiated methodology.

Social implications: Given the severity and frequency of difficulties, sleeping problems in children with autism should be viewed as a significant public health concern. **Originality/value:** This is one of the first studies, qualitative or quantitative, to explore the role and perspectives of fathers of children with autism in the important area of sleep management.

Keywords: autism; children; fathers; sleeping problems; health; couple relationships; siblings

Introduction

Amount and quality of sleep is known to be strongly associated with general well-being, with impact on cognitive development, levels of stress and immune responses (Gallagher, Phillips, & Carroll, 2010). Bruni, (2010) noted that ensuring good sleep in childhood should be viewed as a preventative measure designed to ensure a healthy adult life. It has been estimated that between 40% - 80% of children with autism experience problems relating to sleep, believed to be twice the rate found in typically developing children (Polimeni, Richdale and Francis, 2005). Difficulties include bed-time resistance, sleep onset, night waking, sleep anxiety and low sleep efficiency (Kotagal & Broomall, 2012). Sleep problems in children with autism were not reported to be associated with gender, age, IQ, parental occupation or learning ability, but were linked to the severity of autism symptoms (Mayes, Calhoun, Murray, & Zahid, 2011). Chronological changes in the nature of sleeping problems have been reported, with younger children on the autism spectrum experiencing greater bed-time resistance, parasomnias, night wakings and sleep anxiety while in adolescence problems related to falling asleep, shorter periods of sleep, and tiredness during the day (Goldman, Richdale, Clemons, & Malow, 2012). A number of elements are believed to be causally associated with sleep dysfunction in autism, including genetic, environmental and neurological factors (Devnani & Hegde, 2015).

Sleep problems result in a range of negative consequences for individuals with autism, such as reduced social ability compared to individuals with autism without sleep problems (Lambert et al., 2016), greater communication problems and stricter adherence to rigid routines (Schreck, Mulick, and Smith, 2004). Poor sleep patterns have also been linked to poorer health related quality of life (Delahaye et al., 2014),

1
2
3 increased behaviour problems in both children (Goldman et al., 2011., Hirata et al.,
4
5 2016) and adults with ASD and intellectual impairment (Matson, Ancona, & Wilkins,
6
7 2008).
8
9

10
11
12 The impact of sleep problems on families is pronounced. Parents of typically
13
14 developing children have been found to be at increased risk of stress, problems with
15
16 their own sleep and negative interactions with their children (Polimeni, Richdale and
17
18 Francis 2005). In children with autism, poor sleep has been found to predict levels of
19
20 maternal stress (Herrmann, 2015; and fatigue (Giallo, Wood, Jellett, & Porter, 2013);
21
22 parental sleeping patterns (Meltzer, 2008) and maternal and paternal symptoms of
23
24 depression (Meltzer, 2011). Findings from the online survey of 306 fathers,
25
26 associated with this study (XXX) found that 31% of fathers experienced a little stress
27
28 associated with children's sleeping problems, 27% a moderate amount and 24% "a
29
30 great deal of stress".
31
32
33

34
35 Treatments for sleeping problems in children with autism are reported as limited
36
37 (Tilford et al., 2015), consisting of behavioural approaches where parents are
38
39 educated in the use of sleep hygiene strategies, such as consistent bed-time
40
41 routines (Turner & Johnson, 2013); pharmacological interventions, especially the use
42
43 of melatonin (Goldman et al., 2014) and complementary therapies such as massage
44
45 or the use of weighted blankets (Devnani and Hegde, 2015). The use of sensory
46
47 strategies such as optimising noise and light levels have been reported as having
48
49 some positive impact (Jan, 2008), as have the use of aerobic exercise and motor
50
51 skills (Brand, Jossen, Holsboer-Trachsler, Pühse, & Gerber, 2015).
52
53

54
55 Sleeping difficulties in children with autism are both under-recognised and under-
56
57 treated (Tilford et al., 2015), with qualitative studies exploring the impact of such
58
59
60

1
2
3 problems on families extremely limited. Russell & McCloskey, (2016) suggested that
4
5 such a methodological approach can serve to illuminate omissions in practice and
6
7 support the development of service models which are capable of addressing
8
9 articulated family needs. Paternal perspectives in the field of autism are especially
10
11 rare, with most data collected from mothers (Donaldson et al., 2011). A meta-
12
13 analysis of 404 studies in the field reported that mothers outnumbered fathers by 8 to
14
15 1 as research participants (Braunstein, Peniston, Perelman, & Cassano, 2013).
16
17 Proposed explanations for father exclusion include on-going presumption of maternal
18
19 care (Flippin & Crais, 2011), difficulty in recruiting fathers to research studies
20
21 (Johnson & Simpson, 2013) and a more general deficit model view of father
22
23 competence in the nurturing of children (Brotherson, Dollahite, & Hawkins, 2005). In
24
25 relation to sleep, a comprehensive literature search could discover no qualitative
26
27 studies which focused on paternal experiences of the sleeping problems of their
28
29 children with autism.
30
31
32
33
34
35

36 Braunstein et al., (2013) argued that failure to include fathers of children with autism
37
38 undermines applied research efforts, since paternal treatment and support needs
39
40 have been found to differ from those of mothers in various ways. For example, in a
41
42 study of parental perception of unmet need, Hartley et al., (2015) found that although
43
44 there was significant agreement between fathers and mothers, fathers were more
45
46 likely than mothers to identify parental self care as an area where support was
47
48 lacking. In addition, Papageorgiou & Kalyva, (2010) reported that fathers are much
49
50 more likely than mothers not to disclose their child's autism to colleagues at work,
51
52 and as a result reported receiving less support and understanding overall. It is such
53
54
55
56
57
58
59
60

1
2
3 omissions, regarding qualitative approaches and paternal perspectives in the critical
4
5 area of sleep which the current study seeks to address.
6
7

8
9 Findings presented here are drawn from a wider study of fathers of children with
10
11 autism (XXX). Research questions relevant to this article, were:
12
13

14
15
16 To what extent do children experience problems with sleep?
17

18 What is the nature of any sleeping problems?
19

20 What is the impact of children's sleeping problems on families?
21
22
23
24

25 **Methods**

26 *Research design*

27
28
29 This paper utilises data from a two year mixed methods study. The first phase
30
31 consisted of an online cross sectional survey of fathers of children with autism,
32
33 autism spectrum disorder or Asperger syndrome (hereafter referred to as ASD) and
34
35 during the second phase, 25 semi-structured interviews were undertaken to explore
36
37 survey themes in greater depth.
38
39
40
41
42

43 *Participants and procedure*

44 **Sampling**

45
46
47 A convenience sample of 306 fathers was recruited to an online survey of
48
49 fathers/male carers (namely biological, adoptive, foster or step-fathers) of children
50
51 with ASD, aged up to 19 years and living in the UK. Fathers were recruited through
52
53 major UK autism charity websites, e-mail invitations sent out to over 80 local autistic
54
55
56
57
58
59
60

1
2
3 societies, an advert in a national autism publication for parents and flyers to autism
4
5 specific schools.
6

7
8 Fathers were asked if they would agree to be interviewed within the survey and 185
9
10 fathers did so. A maximum of 25 interviews was specified in the research design in
11
12 the light of time and resources available. Fathers were chosen for interview
13
14 purposively with a view to exploring paternal experiences which might be affected by
15
16 age, socio-economic, ethnic and educational backgrounds, as well as differences in
17
18 children's age and ability (see Table 1). Not all of the 185 men who agreed to be
19
20 interviewed in the survey, remained willing or able to take part and therefore further
21
22 purposive sampling was undertaken until 25 interviews had been arranged.
23
24

25 26 **Procedure**

27
28 Interviews lasted 1-2 hours and occurred at a place and time convenient for fathers.
29
30 Most fathers were interviewed in a neutral venue, during the day and near their place
31
32 of work. Fathers were asked to talk about their involvement in caring for their
33
34 child/ren with autism, parenting challenges and ways of coping (see XXX,
35
36 forthcoming) and pertinent here, to discuss sleeping patterns, any associated
37
38 difficulties and the impact of sleeping problems on family members.
39
40
41
42

43 44 **Participants**

45
46 The majority of fathers completing the survey were white (95%), aged over 40 (80%)
47
48 and married (79%). Most worked full-time 62%) and just over half (52%) were
49
50 educated to degree level or above.
51
52

53
54 Characteristics of the survey sample appear in Table 1.
55
56
57
58
59
60

1
2
3 INSERT TABLE 1 ABOUT HERE
4
5
6

7 Most fathers interviewed were white (n=24), aged 40 or over (n=19) and married
8 (n=23). Seventeen men were educated to degree level or above. Most children were
9 boys (n=20) and 15 were aged 10 years or below. Eight children had a diagnosis of
10 autism, 12, ASD and five, a diagnosis of Asperger Syndrome. Complete
11 characteristics of the interview sample are presented in Table 2.
12
13
14
15
16
17

18
19 INSERT TABLE 2 ABOUT HERE
20
21

22 **Data Analysis**

23

24
25 Copies of transcribed interviews were made available to all fathers to address issues
26 of trustworthiness (Mason, 2012) by providing participants with a further opportunity
27 of verifying that the narrative given accurately reflected their views and experiences.
28 Analysis was guided by aspects of a grounded theory approach (Strauss & Corbin,
29 1998) where theory is “inductively derived from the study of the phenomenon which it
30 represents” (Ezzy, 2002, p.12). This approach involving a ‘constant comparison’
31 (Dey, 2004) of data on a line by line basis, leads to the development of meaning and
32 new understandings, by identifying patterns of similarity or difference. NVIVO 10, a
33 Computer Assisted Qualitative Data Analysis Software programme (CAQDAS)
34 (Bazeley & Richards, 2000) was employed during this process. Hutchison, Johnston,
35 & Breckon, (2010) reflected on the limitations of CAQDAS approaches in terms of
36 generating interpretations while acknowledging that such software can facilitate the
37 systematic handling and organisation of significant amounts of qualitative data by
38 enabling a rigour in the sorting, matching and connecting of data, which leads to the
39 formation of categories and sub-categories. Some aspects of analysis were informed
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 by a priori themes identified in the literature, especially relating to the
4
5 conceptualisation of sleeping problems. Other themes emerged inductively from the
6
7 data, such as those relating to impacts on families.
8
9

10 **Ethics**

11
12
13
14 The research received ethical approval from XXX University. Participants completing
15
16 the online survey were given information concerning research aims, provided with
17
18 assurances of confidentiality and anonymity and were made aware that should they
19
20 go on to complete the survey, consent would be assumed. Fathers interviewed
21
22 received further information, pertaining to their right to decline to answer any
23
24 question and to withdraw from the research at any time without consequences. None
25
26 did so.
27
28
29

30 **Findings**

31
32 All names used are pseudonyms and each father has been assigned a numeric
33
34 code: (F1 = Father 1) etc.
35
36
37

38 ***Sleep associated problems***

39
40
41 Two thirds of fathers interviewed reported a range of significant problems associated
42
43 with their children's sleeping patterns. These are categorised below.
44
45

46 **Bed-time resistance**

47
48
49 Half of fathers reported significant difficulties in getting children into bed, often
50
51 despite the range of strategies intended to facilitate this process (see below): '*And*
52
53 *then it may be a bath, watch a bit of telly, and then get him to bed - which is a lot*
54
55 *more challenging than getting him up in a morning*'. (F17). The transition to bed was
56
57
58
59
60

1
2
3 specifically highlighted by one man: *'It's getting him to move from doing whatever*
4 *he's doing to getting ready for bed.'* (F12). The bed-time routine was often a lengthy
5 one for parents: *'When Ethan's at home, I'm persuading him to go to bed at about*
6 *eleven... The 'going to bed' routine takes about 40-45 minutes'* (F10). Fathers
7 described the growing frustration involved in the evening as they themselves
8 became more tired: *'So, half past eight, he won't go to bed. Half past ten, he still*
9 *won't go to bed. Half past ten..., you're getting irate, you're tired, you've been up*
10 *and down the stairs a million times'*. (F9).

21 **Sleep onset**

22
23
24 A number of fathers described problems children experienced in falling asleep once
25 they were in bed: *'You have to sit with him...otherwise he'll just get out of bed again*
26 *and wander round...eight or nine times out of ten'*(F7). Several described how
27 children's difficulties in falling asleep had led to them sleeping in parents' beds: *'At*
28 *the beginning of our relationship [my partner] used to sleep in the bed with Helen to*
29 *get her to sleep'* (F14). Sometimes complex routines had developed so that children
30 could fall asleep in the parents' bed before being moved back to their own, which in
31 some cases had become difficult to modify. *'Simon gets into our bed, reads for a*
32 *while and falls asleep...And it has to follow the same pattern...but if you ask him not*
33 *to do that, if I'm on an earlier shift, and I actually wanted to go to bed, it's. "oh, no,*
34 *no, no, no, no"* (F18). For a minority of families, the situation was more extreme, with
35 one father stating: *'some nights he just doesn't go to sleep'* (F20) and another: *'So at*
36 *the time when he was 3, 4, 5 years old, he could go three, four days and nights*
37 *without sleep'* (F10).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Night waking

One of the most problematic aspects of children's sleeping difficulties, discussed by two thirds of fathers, was their inability to sleep for long periods through the night, often waking very early. Such disrupted nights could occur frequently: *'He will wake up very often – it could be any time between three o'clock in the morning and five in the morning' (F24)*. Once awake, children could stay awake for long periods: *'she'll be up for about four hours. So last night she woke up at about ten, finally went down at about half past two' (F5)*. For several families, acute problems with sleeping had been going on for a number of years: *'when he was 3, 4, 5 years old, he could go three, four days and nights without sleep... now - he's 14, he goes to bed at eleven. He falls asleep between one and two. He's up again at five.'*(F10). Several fathers reflected on the impact of caring for children through the night as well as during the day: *'With an autistic child there's no break. Yeah, you sleep, but you're half asleep and half awake, because if he wakes up, all the lights are on, all the curtains are open, and you're in bed upstairs' (F11)*.

Night-time roles

The roles of fathers and mothers in managing sleep associated problems differed between families, for a variety of reasons, including the relative demands of parents' employment, health issues, other caring responsibilities and parents' relative ability to cope with sleeplessness. Mothers and fathers often shared the task of taking their children with autism to bed: *'At the moment we have got an alternate going on, where it's one night me, then it's [my wife]' (F20)*. However, as children grew older, the undertaking could become increasingly problematic for some mothers: *'With [my wife]... he's as big as she is, and it's a harder battle for her.'*(F16). One father

1
2
3 explained his central role in the bed-time routine in relation to his wife's other caring
4 tasks: *'we have certain routines that have to be done for him to go downstairs, and*
5 *so I tend to take that on at night because Julia's had it all day, by and large, and*
6 *she's just about spent'. (F21).*
7
8
9
10

11
12
13
14
15 In ten families, fathers had taken the lead in managing children's night-time waking
16 for various reasons. One father cited his wife's job as key: *'some days he would get*
17 *up at four, five o'clock every morning - I would get up with him, because I realise my*
18 *wife [as a teacher] has got thirty kids all day' (F9).* For another family, family
19 finances and health problems influenced role decisions at night: *'It has to be me,*
20 *because my wife is an assistant head and [suffers from] fatigue so her sleep is*
21 *sacrosanct... she's the breadwinner in the house'. (F10).* In some instances, carer's
22 own sleeping patterns was central to role division: *'I'm not the best sleeper myself,*
23 *so it just made more sense that I did that particular duty. It just made it easier for my*
24 *wife' (F24).* Another man felt that he could cope better with night-time supervision
25 duties: *'I do most of it, because my wife doesn't cope very well with being woken up*
26 *in the night' (F5).* A separated father of children with autism explained his process of
27 decision-making regarding getting his children up in the mornings: *'If I get up at six*
28 *o'clock..., I'm going to be shattered by two. And it means it's going to be*
29 *difficult...because it's just me on my own looking after [children] with additional*
30 *needs, so that's quite hard.'* (F19)
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Impact on families

Fathers discussed a range of significant negative effects on their own health and working life, the quality of their partner relationship and the life experiences of their non-disabled children.

Health

Lack of sleep for parents was the most significant health issue raised, with its associated negative impact on daily functioning. Both parents were often affected: '*A few years ago we were getting no sleep. Literally two or three hours' broken sleep a night' (F16)*. One father referred to the range of effects of sleep loss on himself and his wife: '*we were physically and mentally drained. That sleep deprivation thing' (F18)*. For one man, the impact of chronic sleep deprivation over a number of years had resulted in profound consequences for his mental health: '*I had sleep deprivation-induced psychosis, which sounds pretty bad but it was actually worse than it sounds. It was something else' (F10)*. The potential consequences of sleep loss had affected one father's view on his own life expectancy:

'There'd been almost daily articles in the press about how lack of sleep is going to kill you... I made rather more life insurance arrangements than a lot of people would, because there is the very real prospect that I won't see retirement' (F5).

Employment

Seven fathers reported that their children's sleeping issues had caused significant problems in their working lives. One father described how extreme mental health problems associated with sleep deprivation had led to early retirement. Others

1
2
3 discussed the impact of long-term interrupted sleep on their performance in the
4
5 workplace.
6
7

8 *'But from [when he was] four to seven, I'd say, some days I was going in to*
9
10 *work and I was absolutely drained, absolutely drained - and having a full day's*
11
12 *work ahead of me' (F9).*
13
14

15 One father described how he had learnt to manage sleep loss and full-time
16
17 employment over a number of years:
18
19

20 *'For the past fourteen years...we've just adjusted as time goes on. The guys*
21
22 *at work don't understand how sometimes I can function, because I come in,*
23
24 *I'm tired as anything, you know' (F20).*
25
26
27

28 Sometimes fathers explained how colleagues had accommodated their difficulties:
29
30

31 *'But a few years ago we were getting no sleep. Literally two or three hours'*
32
33 *broken sleep a night. And I'd be going in to work... and [colleagues] would*
34
35 *just go 'Bad night with Seth?' and... If I fell asleep while I was parked up*
36
37 *somewhere it would just be like 'Mike's had a bad night with Seth'. (F16).*
38
39

40 Another father explained that his employer's response to his extreme tiredness due
41
42 to his child's night-time waking could be variable:
43
44

45 *'And they say to me 'What can we do to help?' I say 'The main thing I need is*
46
47 *tolerance. Because there are days when I will make mistakes and days when*
48
49 *I'll struggle...and people will seem sympathetic until you're a bit slow or you*
50
51 *make a mistake. In which case it's 'Why haven't you...?' (F5).*
52
53
54
55
56
57
58
59
60

Couple relationship

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Nine fathers described a range of negative effects of sleep problems on their partner relationships. The issue of bed-time resistance resulted in a lack of time for parents to spend together in the evenings: *'we never get any time to ourselves. We can't sort of sit and watch a late-night film because Jo is likely to just sort of wander down'* (F2). A father of a teenager with autism and obsessive compulsive disorder reflected on the impact on stressful bed-times over years on his partner relationship: *'When you're still faced with having to put your son to bed... every night, and that can take up to an hour, there's not a lot of time in the evening for socialising and relaxing... And so it has put a strain on our relationship in many respects.'* (F21). For one couple, the strain caused by their child's sleeping problems almost resulted in the breakdown of their relationship: *'And it got to the stage where we were on the point of divorce. We'd literally reached the point where we had no life'* (F18). Several fathers reported that children's sleeping difficulties affected not only the couple relationship inside the home but also their opportunities for socialising together outside it: *'We've never been able to have a night out since probably Charlie [now aged 12] was born'* (F4). The result of co-sleeping meant that some couples experienced long periods where they could not share a bed: *'I would sleep on a chair in his room. Simply because that was the only option'* (F10). Some couples often shared a bed for only part of the night due to the need for one parent to supervise their child with autism during the night: *'It could be any time between three o'clock in the morning and five in the morning. At which time I usually go into his room. There's a bed made up next to his bed'* (F24.)

Siblings

Four fathers described the ways in which the sleeping problems of their children with autism had affected their other typically developing children. In some cases, bed-time resistance issues resulted in lack of quality time to spend with other children, for example: *'You feel quite sad because you miss out on that time with her'*. (F20).

Night-time waking in children with autism also sometimes resulted in interrupted sleep for siblings: *'It usually has a knock-on effect, Hayley (child with autism) will wake Ethan up, Ethan will wake Sadie up....'* (F14). One father spoke of the way in which sleep loss was affecting the education of his typically developing children: *'her sister is missing a lot of sleep because of her, and so she's behind at school'* (F5).

Another man explained the need for special arrangements for his typically developing daughter at school: *'And the teachers are aware that there are these issues with sleep going on at home. Ellie [sibling] is allowed to sleep at her convenience..at school'* (F10).

Discussion

This study extends our knowledge of the nature of sleeping problems in children with autism and their impact on the daily experience of family members. It appears to be one of the first to explore father involvement and perspectives in this important area.

Fathers reported a range of significant sleep-related difficulties in bed-time resistance, sleep onset, sleep duration and night-time waking and emerged as closely involved in the management of these, both here and in an associated large scale survey (XXX), often sharing or taking lead responsibility for supervising bed-times and periods of night-time waking. Such findings resonate with a growing evidence base which highlights the important, though often overlooked, contribution

1
2
3 of fathers of children with autism in areas of personal care (XXX; (Meadan, Stoner, &
4 Angell, 2015); leisure (Mitchell & Lashewicz, 2015) and education (XXX). More
5
6 research is needed which explores the roles of both mothers and fathers in
7
8 managing sleep. In addition, fathers must be included in co-parenting approaches to
9
10 maximise the impact of such interventions. Braunstein et al., (2013) noted that many
11
12 parent interventions in the field of autism, as in others, fail to report the sex of
13
14 participants, thereby seeming to imply that gender is not a significant factor in
15
16 parental response, when in fact, comprehensive meta-analytic studies have shown
17
18 that men and women may respond differently to the methodology of parent
19
20 education programmes (Panter-Brick et al., 2014; Scourfield, Cheung, & Macdonald,
21
22 2014). In the area of sleep management, Malow et al., (2014), found the use of a
23
24 parent education approach to be effective in improving sleep onset in 80 children
25
26 with autism, although since neither the sex of participants was reported, nor a
27
28 gender-differentiated analysis undertaken, it cannot be assumed that the intervention
29
30 was or would be equally as effective with both parents.
31
32
33
34
35
36

37 Fathers reported significant negative physical and mental health effects of children's
38
39 sleeping problems. In several instances, these were acute, with fathers struggling to
40
41 cope with on-going sleep deprivation, alongside the daily demands of paid
42
43 employment. Such findings resonate with those of previous studies which reported
44
45 that **sleeping problems in children with ASD were found to be related to poorer**
46
47 **mental health (Hodge et al., 2013) and increased stress (Hoffman et al., 2008; Levin**
48
49 **and Scher, 2016) in mothers** and, in addition, raise questions regarding assertions
50
51 that mothers may be at greater risk of negative effects of children's sleeping
52
53 problems compared to fathers (see Polimeni et al., 2005, for example). Tilford et al.,
54
55 (2015) argued that further research is needed regarding the impact of sleeping
56
57
58
59
60

1
2
3 problems on the health of caregivers of children with autism and we would argue that
4 such investigations must include fathers, to gain a more comprehensive
5 understanding of impacts within families and associated gender-differentiated carer
6 needs. Hodge et al., (2015) found that
7
8
9
10

11
12 Greater focus on the effect of sleeping problems on siblings of children with autism
13 and implications for practice, is also required. Difficulties reported here are in
14 keeping with those of other studies where typically developing siblings of children
15 with autism have been shown to have an increased risk of sleeping problems (Chou
16 et al., 2012). Qualitative studies undertaken with siblings themselves would prove
17 especially insightful in understanding the nature and extent of impacts on their lives.
18
19
20
21
22
23
24

25
26 Several targets for intervention are indicated here, relating to increasing the
27 effectiveness of sleep management strategies and minimising the impact of
28 difficulties on family members. Regarding service provision, Ryan, Burton, &
29 Bromley, (2014) testified to the general shortage of sleep support services in the UK,
30 which represents a major shortcoming in view findings presented here. The
31 development of greater expertise amongst primary health care professionals is
32 needed, since as noted by Caulfield, (2011) health professionals have been slow to
33 recognise the importance of identifying and managing sleeping difficulties. The
34 greater use of nurse-led community-based sleep clinics where staff receive intensive
35 and specialist training has been found to be effective in improving quality of life for
36 families having children with learning disabilities and sleeping problems (Ryan et al.,
37 2014) and the further use of such an approach should be investigated for families
38 with children with autism.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 More father inclusive parent education approaches are required. Honaker & Meltzer,
4 (2016) argued for the development of technological approaches in this area such as
5 web-based modules or apps. Such approaches may be especially user-friendly for
6 fathers who are often less able to access face to face provision available during the
7 working week (Cutress & Muncer, 2014) and who may prefer online support and
8 information (Flynn and Stana, 2012; Seymour-Smith, 2013; Watkins & Jefferson,
9 2013).

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Limitations of the current study should be noted. Since the sample consists of men who were from an almost exclusively white ethnic background, two thirds of whom were educated to degree level or above, it cannot be said to be representative of all fathers of children with autism. Further research is needed which is effective in recruiting fathers from more diverse ethnic and educational backgrounds. In relation to ethnicity, for example, Waheed (2015) recommended greater use of a wider range of engagement activities including face-to-face presentations in local community venues and a strategic distribution of flyers. In addition, as noted above, survey data associated with this study indicated that 81% of fathers reported some degree of child sleep-related stress which rose to 100% of fathers interviewed. If we take such stress as a proxy for children experiencing sleeping problems, then fathers having children with such difficulties may be over-represented in both the survey and interview sample, although the rate of sleeping problems in the population of children with ASD could be high, estimated at 40%-80% of children (Polimeni, Richdale and Francis, 2005). Despite these limitations, it is argued that the current study does serve to extend knowledge, given the scarcity of father perspectives in autism and sleep management.

1
2
3 In conclusion, the present study found that the nature of sleeping problems in
4 children with autism resulted in a number of substantial negative effects on fathers,
5 mothers and siblings. Fathers were often closely involved in the management of
6 difficulties. Given the frequency of problems and the scale of negative impacts on
7 families, it is argued that sleeping problems in children with autism should be
8 regarded as a significant public health concern in the UK, urgently warranting an
9 increase in both research and family-centred service provision.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Brand, S., Jossen, S., Holsboer-Trachsler, E., Pühse, U., & Gerber, M. (2015). Impact of aerobic exercise on sleep and motor skills in children with autism spectrum disorders - a pilot study. *Neuropsychiatric Disease and Treatment*, 11, 1911–1920.
- Braunstein, V. L., Peniston, N., Perelman, A., & Cassano, M. C. (2013). The inclusion of fathers in investigations of autistic spectrum disorders. *Research in Autism Spectrum Disorders*, 7(7), 858–865.
- Brotherson, S. E., Dollahite, D. C., & Hawkins, A. J. (2005). Generative Fathering and the Dynamics of Connection between Fathers and Their Children. *Fathering: A Journal of Theory, Research, & Practice about Men as Fathers*, 3, 1–28.
- Bruni, O. (2010). The importance of sleep for children's well-being. *Sleep Medicine* 11 (2010) 599–600, 11, 599–600.
- Caulfield A. (2011). The specialist nurse in the field of childhood sleep disorders. *Community Practitioner*, 84(2), 32–35 4p.
- Chou, M.-C., Chou, W.-J., Chiang, H.-L., Wu, Y.-Y., Lee, J.-C., Wong, C.-C., & Gau, S. S.-F. (2012). Sleep problems among Taiwanese children with autism, their siblings and typically developing children. *Research in Autism Spectrum Disorders*, 6, 665–672.
- Cleary, M., Horsfall, J., & Hayter, M. (2014). Data collection and sampling in qualitative research: does size matter? *Journal of Advanced Nursing*, 70(3), 473-475.

- 1
2
3 Cutress, A. L., & Muncer, S. J. (2014). Parents' views of the National Autistic
4 Society's EarlyBird Plus Programme. *Autism: The International Journal of*
5 *Research & Practice*, 18(6), 651.
6
7
8
9
10 Delahaye, J., Kovacs, E., Sikora, D., Hall, T. A., Orlich, F., Clemons, T. E.,
11 Kuhlthau, K. (2014). The relationship between Health-Related Quality of Life
12 and sleep problems in children with Autism Spectrum Disorders. *Research in*
13 *Autism Spectrum Disorders*, 8, 292–303.
14
15
16
17
18 Dey, I. (2004). Grounded theory. In C. Seale, G. Gobo, G. J. F., & D. Silverman
19 (Eds.), *Qualitative Research Practice*. London: Sage.
20
21
22
23
24 Donaldson, S. O., Elder, J. H., Self, E. H., & Christie, M. B. (2011). Fathers'
25 Perceptions of Their Roles During In-Home Training for Children With Autism.
26 *Journal of Child & Adolescent Psychiatric Nursing*, 24(4), 200–207.
27
28 <http://doi.org/10.1111/j.1744-6171.2011.00300.x>
29
30
31
32
33 Flippin, M., & Crais, E. R. (2011). The Need for More Effective Father Involvement in
34 Early Autism Intervention. *Journal of Early Intervention*, 33(1), 24–50.
35
36
37
38 Flynn, M. A., & Stana, A. (2012). Social support in a men's online eating disorder
39 forum. *International Journal of Men's Health*, 11(2), 150–169.
40
41
42
43
44 Gallagher, S., Phillips, A. C., & Carroll, D. (2010). Parental stress is associated with
45 poor sleep quality in parents caring for children with developmental
46 disabilities. *Journal of Pediatric Psychology*, 35(7), 728–737.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Goldman, S. E., McGrew, S., Johnson, K. P., Richdale, A. L., Clemons, T., & Malow,
4
5 B. A. (2011). Sleep is associated with problem behaviors in children and
6
7 adolescents with Autism Spectrum Disorders. *Research in Autism Spectrum*
8
9 *Disorders, 5*, 1223–1229.
- 10
11 Goldman, S., Richdale, A., Clemons, T., & Malow, B. (2012). Parental Sleep
12
13 Concerns in Autism Spectrum Disorders: Variations from Childhood to
14
15 Adolescence. *Journal of Autism & Developmental Disorders, 42*(4), 531–538
16
17
18 8p.
- 19
20 Hartley, S. L., & Schultz, H. M. (2015). Support needs of fathers and mothers of
21
22 children and adolescents with autism spectrum disorder. *Journal of Autism*
23
24 *and Developmental Disorders, 45*(6), 1636–1648.
- 25
26 Herrmann, S. (2015). Article: Counting Sheep: Sleep Disorders in Children With
27
28 Autism Spectrum Disorders. *Journal of Pediatric Health Care*. Retrieved from
29
30 [http://ezproxy.leedsbeckett.ac.uk/login?url=http://search.ebscohost.com/login.](http://ezproxy.leedsbeckett.ac.uk/login?url=http://search.ebscohost.com/login.aspx?direct=true&db=edselp&AN=S0891524515002308&site=eds-live&scope=site)
31
32
33
34
35
36
37
38
39
40 Hirata, I., Mohri, I., Kato-Nishimura, K., Tachibana, M., Kuwada, A., Kagitani-
41
42 Shimono, K., Taniike, M. (2016). Sleep problems are more frequent and
43
44 associated with problematic behaviors in preschoolers with autism spectrum
45
46 disorder. *Research in Developmental Disabilities, 49-50*, 86–99.
- 47
48 Hodge, D., Hoffman, C., Sweeney, D., & Riggs, M. (2013). Relationship Between
49
50 Children's Sleep and Mental Health in Mothers of Children with and Without
51
52 Autism. *Journal of Autism & Developmental Disorders, 43*(4), 956–963 8p.
- 53
54 Hoffman, C. D., Sweeney, D. P., Lopez-Wagner, M. C., Hodge, D., Nam, C. Y., &
55
56 Botts, B. H. (2008). Children with Autism: Sleep Problems and Mothers'
57
58
59
60

- 1
2
3 Stress. *Focus on Autism and Other Developmental Disabilities*, 23(3), 155–
4
5 165.
6
7
8 Honaker, S. M., & Meltzer, L. J. (2016). Clinical review: Sleep in pediatric primary
9
10 care: A review of the literature. *Sleep Medicine Reviews*, 25, 31–39.
11
12 Jan J.E, Owens, J. A. Weiss, M. D. et al (2008) Sleep hygiene for children with
13
14 neurodevelopmental Disabilities. *Pediatrics*. 122, 6, 1343-1350.
15
16 Johnson, N. L., & Simpson, P. M. (2013). Lack of father involvement in research on
17
18 children with autism spectrum disorder: maternal parenting stress and family
19
20 functioning. *Issues in Mental Health Nursing*, 34(4), 220–228.
21
22
23 Levin, A., & Scher, A. (2016). Sleep problems in young children with autism
24
25 spectrum disorders: A study of parenting stress, mothers' sleep-related
26
27 cognitions, and bedtime behaviors. *CNS Neuroscience & Therapeutics*,
28
29 22(11), 921–927.
30
31
32 Kotagal, S., & Broomall, E. (2012). Review Article: Sleep in Children With Autism
33
34 Spectrum Disorder. *Pediatric Neurology*, 47, 242–251.
35
36 Lambert, A., Tessier, S., Rochette, A.-C., Scherzer, P., Mottron, L., & Godbout, R.
37
38 (2016). Poor sleep affects daytime functioning in typically developing and
39
40 autistic children not complaining of sleep problems: A questionnaire-based
41
42 and polysomnographic study. *Research in Autism Spectrum Disorders*, 23,
43
44 94–106.
45
46
47 Mason, J. (2012). *Qualitative researching* (2nd ed.). London: Sage.
48
49 Malow, B., Adkins, K., Reynolds, A., Weiss, S., Loh, A., Fawkes, D., ... Clemons, T.
50
51 (2014). Parent-Based Sleep Education for Children with Autism Spectrum
52
53 Disorders. *Journal of Autism & Developmental Disorders*, 44(1), 216–228 13p.
54
55
56
57
58
59
60

- 1
2
3 Matson, J. L., Ancona, M. N., & Wilkins, J. (2008). Sleep Disturbances in Adults with
4
5 Autism Spectrum Disorders and Severe Intellectual Impairments. *Journal of*
6
7 *Mental Health Research in Intellectual Disabilities*, 1(3), 129–139.
8
9
10 Mayes, S., Calhoun, S., Murray, M., & Zahid, J. (2011). Variables Associated with
11
12 Anxiety and Depression in Children with Autism. *Journal of Developmental &*
13
14 *Physical Disabilities*, 23(4), 325–337.
15
16 Meadan, H., Stoner, J. B., & Angell, M. E. (2015). Fathers of children with autism:
17
18 perceived roles, responsibilities, and support needs. *Early Child Development*
19
20 *and Care*, 1–17.
21
22
23 Meltzer, L. J. (2011). Factors associated with depressive symptoms in parents of
24
25 children with autism spectrum disorders. *Research in Autism Spectrum*
26
27 *Disorders*, 5, 361–367.
28
29
30 Meltzer, L. J. (2008). Brief report: Sleep in parents of children with autism spectrum
31
32 disorders. *Journal of Pediatric Psychology*, 33, 380–386.
33
34 Mitchell, J., & Lashewicz, B. (2015). More Than a Pal: The Generative Leisure Work
35
36 of Fathers Raising Children with Autism Spectrum Disorder. *Fathering*, 13(2),
37
38 130–145.
39
40
41 Moss, A., Gordon, J., & O'Connell, A. (2014). Impact of Sleepwise: an intervention
42
43 for youth with developmental disabilities and sleep disturbance. *Journal of*
44
45 *Autism & Developmental Disorders*, 44(7), 1695–1707.
46
47
48 Panter-Brick, C., Burgess, A., Eggerman, M., McAllister, F., Pruett, K., & Leckman, J.
49
50 F. (2014). Practitioner Review: Engaging fathers - recommendations for a
51
52 game change in parenting interventions based on a systematic review of the
53
54 global evidence. *Journal of Child Psychology & Psychiatry*, 55(11), 1187–
55
56 1212.
57
58
59
60

- 1
2
3 Papageorgiou, V., & Kalyva, E. (2010). Self-reported needs and expectations of
4
5 parents of children with autism spectrum disorders who participate in support
6
7 groups. *Research in Autism Spectrum Disorders*, 4, 653–660.
8
9
10 Polimeni M. A., Richdale. A. L., & Francis, A. P. (2005). A survey of sleep problems
11
12 in autism, Asperger's disorder and typically developing children. *Journal of*
13
14 *Intellectual Disability Research*, 49(4), 260–268.
15
16
17 Robson, C. (2011) *Real World Research*. London: Wiley.
18
19 Russell, S., & McCloskey, C. R. (2016). Parent Perceptions of Care Received by
20
21 Children with an Autism Spectrum Disorder. *Journal of Pediatric Nursing*, 31,
22
23 21–31.
24
25
26 Ryan, G., Burton, L., & Bromley, D. (2014). A nurse-led sleep service for children
27
28 and young people with disability. *Primary Health Care*, 24(6), 16–23.
29
30 Schreck, K. A., Mulick, J. A., & Smith, A. F. (2004). Sleep problems as possible
31
32 predictors of intensified symptoms of autism. *Research in Developmental*
33
34 *Disabilities*, 25(1), 57–66.
35
36
37 Scourfield, J., Cheung, S. Y., & Macdonald, G. (2014). Working with fathers to
38
39 improve children's well-being: Results of a survey exploring service provision
40
41 and intervention approach in the UK. *Children and Youth Services Review*,
42
43 40.
44
45
46 Seymour-Smith, S. (2013). A reconsideration of the gendered mechanisms of
47
48 support in online interactions about testicular implants: A discursive approach.
49
50 *Health Psychology*, 32(1), 91–99.
51
52
53 Strauss, A., & Corbin, J. (1990). *Basics of Qualitative Research: Grounded Theory*
54
55 *Procedures and Techniques*. London: Sage.
56
57
58
59
60

- 1
2
3 Tilford, J. M., Payakachat, N., Kuhlthau, K. A., Pyne, J. M., Kovacs, E., Bellando, J.,
4
5 ... Frye, R. E. (2015). Treatment for Sleep Problems in Children with Autism
6
7 and Caregiver Spillover Effects. *Journal of Autism and Developmental*
8
9 *Disorders, 45*(11), 3613–3623.
10
11
12 Turner, K. S., & Johnson, C. R. (2013). Behavioral interventions to address sleep
13
14 disturbances in children with autism spectrum disorders: a review. *Topics in*
15
16 *Early Childhood Special Education, 33*(3), 144–152.
17
18
19 Waheed, W., A. Hughes-Morley, A. Woodham, G. Allen, and P. Bower. (2015).
20
21 Overcoming Barriers to Recruiting Ethnic Minorities to Mental Health
22
23 Research: A Typology of Recruitment Strategies. *BMC, Psychiatry 15* (1):
24
25 555–111.
26
27
28 Watkins, D. C., & Jefferson, S. O. (2013). Recommendations for the use of online
29
30 social support for African American men. *Psychological Services, 10*(3), 323–
31
32 332.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Characteristics of survey sample

<i>Characteristic</i>	<i>N (%)</i>
<i>Age</i>	
21-29	11 (4%)
30-39	51 (17%)
40-49	159 (52%)
50-59	77 (25%)
60-69	8 (3%)
<i>Number of children with autism</i>	
1	257 (84%)
2	40 (13%)
3	8 (3%)
4	1 (1%)
<i>Relationship to child with autism</i>	
Biological father	287 (94%)
Step-father	13 (4%)
Adoptive father	4 (1%)
Foster father	2 (1%)
<i>Marital status</i>	
Married	242 (79%)
In a partnership	38 (12%)
Separated/Divorced/widowed	21 (6%)
Single	5 (2%)
<i>Educational background (highest level)</i>	
Completed primary/secondary school	26 (9%)
Passes at GCSE/ A level	61 (20%)
Vocational qualifications	59 (19%)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Degree/ postgraduate qualifications 160 (52%)

Employment status

Full-time 190 (62%)

Self-employed 42 (14%)

Retired/Not in paid work 46 (15%)

Part-time 27 (9%)

Ethnicity

White 286 (95%)

Mixed heritage 6 (2%)

Black or Black British 5 (2%)

Asian or Asian British 2 (1%)

Chinese 1 (1%)

Note percentages may not add up to 100% due to rounding. Some values were missing.

Table 2: Characteristics of interview sample

<i>Characteristic</i>	<i>N (%)</i>
<i>Age</i>	
21-29	2
30-39	4
40-49	12
50-59	7
<i>Number of children with autism</i>	
1	20
2	4
3	1
<i>Relationship to child with autism</i>	
Biological father	23
Step-father	2
<i>Marital status</i>	
Married	23
Living with partner	1
Separated/Divorced	1
<i>Educational background (highest level)</i>	
Passes at GCSE/ A level	1
Vocational qualifications	7

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Degree/ postgraduate qualifications 17

Employment status

Full-time 11

Self-employed 6

Retired/ Not in paid work 3

Part-time 5

Ethnicity

White 24

Black or Black British 1

