



LEEDS
BECKETT
UNIVERSITY

Citation:

Goff, L and Silva, MJ and Bordoli, C and Enayat, ZE and Cassidy, A and Maynard, M and Harding, S (2017) Longitudinal associations of skipping breakfast with ethnicity and cardiometabolic risk: the Determinants of Adolescence, now young Adults, Social well-being and Health Study (DASH). Proceedings of the Nutrition Society, 76 (OCE4). ISSN 1475-2719 DOI: <https://doi.org/10.1017/S0029665117003305>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4991/>

Document Version:

Article (Published Version)

Published Abstract

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.



Summer Meeting, 10–12 July 2017, Improving Nutrition in Metropolitan Areas

Longitudinal associations of skipping breakfast with ethnicity and cardiometabolic risk: the Determinants of Adolescence, now young Adults, Social well-being and Health Study (DASH)

L.M. Goff¹, M.J. Silva¹, C. Bordoli¹, Z.E. Enayat¹, A. Cassidy¹, M. Maynard² and S. Harding¹

¹Division of Diabetes & Nutritional Sciences, King's College London, UK and

²School of Health & Well-Being, Leeds Beckett University, UK

Ethnic inequalities in cardiometabolic disease^(1,2) may be explained by differences in diet and lifestyle. Poor dietary habits, such as skipping breakfast and consumption of fizzy drinks and fast foods are more common amongst ethnic minority children and adolescents^(3,4). The long-term effects of these childhood behaviours on adult cardiometabolic risk factors have not yet been investigated in an ethnically diverse population. We aimed to assess ethnic patterns in adolescent and young adult breakfast skipping and its influence on cardiometabolic risk in young adulthood amongst a diverse UK cohort.

The DASH cohort was recruited in 2002/03 and consisted of 6643 11–13 year olds, sampled to represent the main ethnic groups of the UK population. The 'DASH 10 years on' study is a longitudinal follow-up of a subset of the cohort who are now young adults (21–23 years). Participants had anthropometric measures (weight, BMI, waist circumference), blood pressure, total and HDL-cholesterol and HbA1c assessed and completed a short dietary behaviours questionnaire indicating how frequently they consume breakfast (daily, 3–4 days a week, 1–2 days a week, never/hardly ever).

The cohort consisted of 311 males (age 22.8 (95 % CI 22.7, 22.9) years; BMI 24.7 (95 % CI 24.3, 25.2) kg/m²) and 316 females (age 22.7 (95 % CI 22.6, 22.8) years; BMI 24.9 (95 % CI 24.3, 25.5) kg/m²). A total of 107 White British, 102 Black Caribbean, 132 Black African, 99 Indian, 111 Bangladeshi or Pakistani and 115 Other (mainly mixed) were included in the follow-up. In young adulthood regular breakfast skipping was reported by 56 % of participants; Black African participants were more likely to skip breakfast than White British (OR: 1.81, 1.04 to 3.17, $p = 0.004$). The highest proportion of breakfast skipping occurred amongst the Black Caribbean (66 %) and Black African (64 %) groups and the lowest amongst Indian participants (46 %). The impact of skipping breakfast during both adolescence and young adulthood on cardiometabolic risk factors during young adulthood were investigated using multivariate regression modelling. Skipping breakfast at 11–13 years was a significant determinant of BMI at 21–23 years (1.45 (95 % CI 0.61, 2.29), $p = 0.001$) as was skipping breakfast at 21–23 years, although the effect was slightly attenuated in this age group (0.92 (95 % CI 0.1, 1.73), $p = 0.027$). Skipping breakfast at both 11–13 years and 21–23 years were also important determinants of total cholesterol levels (11–13 years: 0.17 (95 % CI 0.01, 0.33), $p = 0.041$; 21–23 years: 0.23 (95 % CI 0.07, 0.38), $p = 0.003$).

This is the first longitudinal assessment of breakfast skipping and its impact on cardiometabolic risk factors amongst an ethnically diverse cohort of young adults in the UK. In this work we have recognised the detrimental impact of childhood breakfast skipping on cardiometabolic risk factors, such as BMI and cholesterol concentrations, in young adulthood. Furthermore we have identified distinct ethnic patterns in breakfast skipping, such that skipping breakfast is most prevalent amongst Black African and Caribbean groups and less common amongst Indians. Our findings provide a useful insight into dietary behaviours that health promotion campaigns could target in aiming to improve the diets of young people, and highlights the importance of targeting interventions to improve dietary behaviours such as breakfast consumption at specific groups of young adults in the population.

1. Becker E *et al.* (2006) *National Centre for Social Research*.
2. Zhang Q *et al.* (2009) *Ethnicity & Health* 14(5): 439–57.
3. Harding S *et al.* (2008) *Int J Epi* 37(1): 162–72.
4. Nicklas TA *et al.* (1998) *J Am Diet Assoc* 98(12): 1432–8.