



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Breen, S and Harrison, AJ and Kenny, IC (2010) HIP ROTATION RANGE OF MOTION AND ITS IMPACT ON LOWER LIMB ALIGNMENT ON LANDING. ISSN 1999-4168

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/746/>

Document Version:

Article (Published Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

# HIP ROTATION RANGE OF MOTION AND ITS IMPACT ON LOWER LIMB ALIGNMENT ON LANDING

Sarah Breen<sup>1</sup>, Drew Harrison<sup>1</sup>, and Ian Kenny<sup>1</sup>

Biomechanics Research Unit, University of Limerick, Limerick, Ireland<sup>1</sup>

The purpose of the present study was to compare lower limb alignment at initial ground contact between groups with normal and abnormal hip rotation range of motion. Male (n=8) and female (n=8) subjects performed an maximal drop jump diagonal side cut task ten to the left and ten to the right. Lower limb alignment was assessed through knee angle, hip angle, ankle angle, thigh rotation and shank rotation at initial foot contact. One significant difference was reported between groups for the knee angle variable on the non dominant side. This indicates that the only the knee angle variable is affected by unbalanced hip rotation range of motion and on the non-dominant side.


**KEY WORDS:** Anterior Cruciate Ligament, joint angles, limb dominance, segment rotations

**INTRODUCTION:** Anterior cruciate ligament (ACL) injuries are well recognised as one of the most common and serious sports injuries with upwards of 250,000 ACL injuries in the United States each year (Boden *et al.* 2000). Seventy two percent of non-contact ACL injuries occur at or shortly after foot strike (Myklebust *et al.* 1997, Olsen *et al.* 2004). Lower limb alignment at the moment of foot strike has been shown to effect valgus loading at the knee (McLean *et al.*, 2004a; Zeller *et al.*, 2003) with greater valgus loads increasing ACL injury risk. Numerous factors can alter lower limb alignment at landing such as hip joint neuromuscular control, strength and range of motion (ROM). Neuromuscular control at the hip has been suggested to be functional as a means of countering ACL injury inducing valgus loads (Besier *et al.* 2003, Lloyd and Buchanan 2001, Zhang and Wang 2001). Decreased ROM at the hip has also been linked to injury risk for conditions such as lower back pain (Verrall *et al.* 2007, Ireland and Wall 1990). Hip rotation ROM has also been shown to differentiate ACL injured subjects from the general population (Gomes *et al.* 2008); it is unclear if this would still be the case prospectively. Lower limb alignment at landing and hip function have been previously investigated in terms of their affect on valgus loading and ACL injury risk however there is no indication of any interaction between the two. This study aimed to investigate differences in initial contact lower limb alignment dependent variables (knee angle, hip angle, ankle angle, thigh rotation, shank rotation) between two subject groups divided according to hip rotation ROM balance (independent variable) normal with external rotation  $<10^{\circ}$  different from internal rotation (balanced) and abnormal with  $>10^{\circ}$  difference (unbalanced). It was hypothesised that the group with abnormal hip rotation would display different landing alignment to that of the group with normal hip rotation ROM.

**METHOD:** Subjects included eight males (age  $21 \pm 3$  yrs; height  $1.79 \pm 0.06$  m; mass  $76 \pm 6$  kg) and eight females (age  $21 \pm 2$  yrs; height  $1.68 \pm 0.07$  m; mass  $64 \pm 7$  kg). Passive hip rotation ROM assessment took place prone on a plinth style bed (Harris-Hayes *et al.* 2007) using manual pelvic stabilisation and a bio-med gravity based inclinometer for measurement of shank orientation (Cibulka *et al.* 1998). Following ROM and maximum drop jump height assessment subjects completed 20 trials of a dynamic task. This involved dropping from a 0.30 m bench, and performing an immediate drop jump up to reach and touch a target which was suspended at their maximum drop jump height. The suspended target triggered a directional cueing system which indicated which direction the subject had to run diagonally to on landing (10 left 10 right), set up is shown in Figure 1.

**Data Collection:** A six-camera high-speed motion analysis system (Eagle; Motion Analysis Corp., Santa Rosa, CA) (500Hz) was synchronised with an AMTI dual force platform system (1000 Hz). 43 retro-reflective markers were secured on the asis, psis, sacrum, iliac crest, greater trochanter, medial and lateral epicondyle and malleolus, upper and lower calcaneous, 2nd and 5th metatarsal of both legs four marker clusters were also placed on the thigh and shank. Each subject stood for a static trial prior to full data collection.

**Data Analysis:** Hip rotation groupings were defined as mentioned according to hip rotation ROM. The abnormal grouping was unidirectional demonstrating an increased level of external rotation. These groups had significantly different internal external rotation ratio's with



**Figure 1. Dynamic Task Set Up**

a mean difference of 29°\*. Knee hip and ankle angles were presented as anatomical angles thigh and shank rotations were presented as the difference between standing position (static trial) and the position at ground contact. The differences between the hip rotation groups were assessed by a one way anova and Cohen's d was used as a measure of effect size. Pearson's correlation was also used to assess correlations between ROM internal external rations and the dependant variables.

**RESULTS:** Average values for all lower limb alignment variables recorded at ground contact are presented in Table 1 for both groups. One significant difference was found between the groups for the knee angle variable on the dominant leg. The group with abnormal hip rotation ROM demonstrated significantly more knee extension on landing in their non-dominant leg that the normal hip rotation ROM group. There was a very strong effect size also shown for this variable.

**Table 1 Lower limb alignment variables at initial contact for both groups**

(°)	Normal Hip ROM		>Ext Hip ROM		p-value		Cohen's d	
	Dominant	Non-Dominant	Dominant	Non-Dominant	Dominant	Non-Dominant	Dominant	Non-Dominant
Knee Angle	159	<b>156*</b>	162	<b>168*</b>	0.258 <sup>a</sup>	<b>0.003*</b> <sup>f</sup>	0.59	<b>1.78</b>
Hip Angle	131	122	126	122	0.181 <sup>b</sup>	0.871 <sup>g</sup>	0.73	0.08
Ankle Angle	133	133	125	125	0.309 <sup>c</sup>	0.305 <sup>h</sup>	0.55	0.55
Thigh Rotation	32	26	34	33	0.8 <sup>d</sup>	0.317 <sup>i</sup>	0.13	0.52
Shank Rotation	19	14	10	23	0.26 <sup>e</sup>	0.144 <sup>j</sup>	0.59	0.78

**Power Analysis**

<sup>a</sup>= 0.196 <sup>b</sup>=0.259 <sup>c</sup>=0.166 <sup>d</sup>=0.057 <sup>e</sup>=0.195 <sup>f</sup>=0.897 <sup>g</sup>=0.053 <sup>h</sup>=0.168 <sup>i</sup>=0.162 <sup>j</sup>=0.302

When a Pearson's correlation was implemented on hip rotation range of motion and each dependent variable one distinct relationship was shown for knee angle on the non dominant leg with a correlation coefficient of 0.54\*, significant at 0.030 (Figure 2).




Figure 2. Knee Angle - Hip Rotation ROM Correlation

**DISCUSSION:** The purpose of this study was to investigate any differences in initial contact lower limb alignment variables (knee angle, hip angle, ankle angle, thigh rotation, shank rotation) between two subject groups divided according to hip rotation ROM. There was one significant difference shown between the two groups (non dominant knee angle) but overall the proposed hypothesis was not supported. There has been no previous research assessing the interaction between initial contact lower limb alignment and levels of hip rotation ROM. This investigation is therefore important in establishing that hip ROM does not have a significant effect on all of the landing alignment variables. The one variable that did show an interaction with hip rotation ROM was knee angle on the non dominant leg. Non-dominant knee angle at contact was significantly different between groups with a strong effect size and 29% of its variability was explained by each subject's hip rotation ROM (Pearson's correlation coefficient 0.54). The knee at ground contact in the abnormal hip rotation ROM group was significantly more extended which has been cited as a potential risk factor for ACL injury (Boden *et al.* 2000). The fact that this is only the case in the non-dominant limb is also interesting as limb dominance has also been assessed as a potential ACL injury risk factor. It is thought that more ACL injuries occur on the non-dominant side but this has not yet been proven conclusively (Matava *et al.* 2002). Overall the proposed hypothesis was not supported as only one of five variables differed between groups. As stated in the introduction many factors affect lower limb alignment at landing; it is plausible that factors such as neuromuscular control and strength around the hip joint may demonstrate a stronger effect than the variables utilised in this study on lower limb alignment at landing. Additional measures that were not employed in this study which would have provided further depth to the investigation was valgus angle and joint moments, future analysis of those variables for the given data is planned. Areas that may alter lower limb alignment on landing and may merit future research include neuromuscular control, and strength at the hip. Variables such as muscle strength can be easily targeted in injury prevention interventions for the production of safer lower limb alignments at landing.

**CONCLUSION:** One variable from five lower limb alignment variables was shown to be significantly different when compared in terms of hip rotation, this variable was knee angle on the dominant leg. This relationship between knee angle and hip rotation is important as the more extended position adopted by those with abnormal hip rotation ROM may place them at

an increase risk of ACL injury. Hip rotation ROM is easily targeted by stretching interventions which may act to decrease knee extension at ground contact and decrease injury risk. Future research is necessary to investigate the relationship between other hip joint functional measures such as strength and balance etc. This would assist in the development of other appropriate injury prevention programmes for the alteration of lower limb alignment at landing to decrease injury risk.

## REFERENCES:

- Besier, T., Lloyd, D. and Ackland, T. (2003) Muscle activation strategies at the knee during running and cutting maneuvers, *Medicine & Science in Sports & Exercise*, **35**, 119.
- Boden, B.P., Dean, G.S., Feagin Jr, J.A., Garrett Jr, W.E (2000) Mechanisms of anterior cruciate ligament injury *Orthopedics*, **23**, 573-578
- Cibulka, M. T., Sinacore, D. R., Cromer, G. S. and Delitto, A. (1998) Unilateral hip rotation range of motion asymmetry in patients with sacroiliac joint regional pain, *Spine*, **23**, 1009.
- Gomes, J. L. E., de Castro, J. V. and Becker, R. (2008) Decreased Hip Range of Motion and Noncontact Injuries of the Anterior Cruciate Ligament, *Arthroscopy: The Journal of Arthroscopic and Related Surgery*, **24**, 1034-1037.
- Harris-Hayes, M., Wendl, P. M., Sahrmann, S. A. and Van Dillen, L. R. (2007) Does stabilization of the tibiofemoral joint affect passive prone hip rotation range of motion measures in unimpaired individuals? A preliminary report, *Physiotherapy Theory and Practice*, **23**, 315-323
- Ireland, M. and Wall, C. (1990) Epidemiology and comparison of knee injuries in elite male and female United States basketball athletes, *Med Sci Sports Exerc*, **22**, S82.
- Lloyd, D. and Buchanan, T. (2001) Strategies of muscular support of varus and valgus isometric loads at the human knee, *Journal of biomechanics*, **34**, 1257-1267
- Matava, M.J., Freehill, A.K., Grutzner, S. And Shannon, W. (2002) Limb dominance as a potential etiologic factor in noncontact anterior cruciate ligament tears, *Journal of Knee Surgery*, **15**, 11-6
- Mc Lean, S.G., Lipfert, S.W. and van Den Bogart, A.J. (2004) Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Medicine and Science in Sports and Exercise*, **36**, 1008-1016.
- Myklebust, G., Maehlum, S., Engebretsen, L., Strand, T. and Solheim, E. (1997) Registration of cruciate ligament injuries in Norwegian top level team handball. A prospective study covering two seasons, *Scandinavian Journal of Medicine and Science in Sports*, **7**, 289-292.
- Olsen, O., Myklebust, G., Engebretsen, L. and Bahr, R. (2004) Injury Mechanisms for Anterior Cruciate Ligament Injuries in Team Handball: A Systematic Video Analysis, *American Journal of Sports Medicine*, **32**, 1002-1012.
- Verrall, G., Slavotinek, J., Barnes, P., Esterman, A., Oakeshott, R. and Spriggins, A. (2007) Hip joint range of motion restriction precedes athletic chronic groin injury, *Journal of Science and Medicine in Sport*, **10** 463-466.
- Zeller, B., McCrory, J., Kibler, W. and Uhl, T. (2003) Differences in Kinematics and Electromyographic Activity Between Men and Women during the Single-Legged Squat, *The American Journal of Sports Medicine*, **31**, 449.
- Zhang, L. and Wang, G. (2001) Dynamic and static control of the human knee joint in abduction-adduction, *Journal of biomechanics*, **34**, 1107-1115

*Acknowledgement:* The authors would like to thank the following for their contribution to the data collection process: Joseph Costello and Catherine Tucker. The Irish Research Council for Engineering and Science are also acknowledged for the funding provided which supported this work.