



LEEDS
BECKETT
UNIVERSITY

Citation:

Till, K and Jones, B (2015) Monitoring anthropometry and fitness using maturity groups within youth rugby league. *Journal of strength and conditioning research / National Strength & Conditioning Association*, 29 (3). 730 - 736. ISSN 1064-8011 DOI: <https://doi.org/10.1519/JSC.0000000000000672>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/88/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

1 **Monitoring anthropometry and fitness using maturity groups within youth rugby league**

2

3

Running Head: Anthropometry and fitness within maturity groups

4

5

 Kevin Till and Ben Jones

6

7

 Research Institute for Sport, Physical Activity and Leisure, Leeds Metropolitan University,

8

 Leeds, West Yorkshire, United Kingdom

9

10

11 Corresponding Author:

12

 Dr. Kevin Till

13

 Room 111, Fairfax Hall

14

 Research Institute for Sport, Physical Activity and Leisure

15

 Headingley Campus, Leeds Metropolitan University

16

 W.Yorkshire, LS6 3QS

17

 Phone: (044-11) 01132-832600 Ext: 25182

18

 Email: k.till@leedsmet.ac.uk

19

20

1

2 **Monitoring anthropometry and fitness using maturity groups within youth rugby league**

3

4 **Running Head:** Anthropometry and fitness within maturity groups

5

6

7

ABSTRACT

1
2 The purpose of the present study was to evaluate the anthropometry and fitness, and
3 change in these characteristics over time, of youth rugby league players by using maturity
4 status to determine annual categories instead of traditional chronological annual-age
5 grouping. One hundred and twenty one male rugby league players were assessed using
6 anthropometric (i.e., height, sitting height, body mass and sum of four skinfolds) and fitness
7 (i.e., vertical jump, medicine ball chest throw, 10m and 20m sprint and multi stage fitness
8 test; MSFT) measures over a 5 year period. Each player was classified into one of six
9 maturity groups based on their maturity offset (Years from Peak Height Velocity; i.e., 1.5
10 YPHV). MANOVA analyses identified significant ($p < 0.001$) main effects for maturity group
11 for cross-sectional characteristics and longitudinal change in performance over time. Analyses
12 demonstrated that more mature groups had greater anthropometric and fitness characteristics,
13 except for endurance performance (MSFT -2.5 YPHV = 1872 ± 18 m vs 2.5 YPHV = $1675 \pm$
14 275 m). For longitudinal changes in characteristics over time, a significant effect was only
15 identified for height and sitting height ($p < 0.05$). These findings provide comparative data for
16 anthropometric and fitness characteristics and change in performance over time in accordance
17 to maturity status within youth rugby league players. Classifying players into annual maturity
18 groups may be an additional or alternative assessment method for evaluating anthropometry
19 and fitness performance in adolescent populations. Further, tracking performance changes
20 over time, especially in relation to maturation, may reduce the limitations associated with
21 chronological annual-age grouping.

22

23

24 **Key Words:** Maturation, longitudinal, player assessment, talent identification, talent
25 development

INTRODUCTION

1
2 The assessment of anthropometric and fitness characteristics of adolescent athletes is
3 commonly used within research and practice across youth sports, with literature available that
4 presents comparative data within such populations (e.g., rugby union, 25; soccer, 11;
5 volleyball, 21). These anthropometric and fitness characteristics are often collected and
6 analysed by strength and conditioning coaches to assist with talent identification and monitor
7 the responses and development of physical characteristics in relation to various training
8 programmes. Traditionally in youth sport contexts, players are assigned, compete and are
9 selected within chronological annual-age categories (i.e., Under 13s) similar to educational
10 systems. As this chronological annual-age grouping process is common, athlete characteristics
11 are always presented, assessed and evaluated within such annual-age categories.

12 During adolescence, maturation (i.e., the timing and tempo of progress towards the
13 adult mature state, 16) varies considerably between individuals of the same chronological age
14 (4). As physical performance is related to biological maturation during adolescence (15,22),
15 boys advanced in biological maturity are generally better performers in physical tasks (e.g.,
16 speed, strength, power) than their later maturing peers (17). Since maturation and
17 chronological age rarely progress at the exact same rate (15), comparisons of characteristics
18 using chronological annual-age categories, can lead to youths being (dis)advantaged due to
19 their maturity status (2). These maturational (dis)advantages have resulted in the selection of
20 relatively older (5) and earlier maturing (18,24,28) players to representative levels within
21 youth sport. Although this relationship is apparent and it has been recommended to consider
22 maturity status in the evaluation of performance for over 15 years (3), only recently have
23 studies began to consider maturation in the evaluation of physical characteristics within youth
24 athletes (30,32). Based on the effect of maturity on performance and selection within

1 adolescent populations, it may seem more appropriate to consider individuals by maturity
2 instead of traditional chronological annual-age grouping systems.

3 Alongside, presenting data within chronological annual-age categories, current
4 research is predominantly cross-sectional with performance often only measured at one
5 specific time point. Recent recommendations suggest monitoring performance longitudinally
6 to assess the changes that occur in characteristics over time (34), which would allow the
7 evaluation of the development of characteristics within and among youth athletes to be more
8 easily identifiable (1). However, research observations tracking characteristics longitudinally
9 within adolescent athletes are limited (6,36), especially considering maturational status
10 (22,31).

11 Due to the physically demanding nature of rugby league, players are required to have
12 highly developed fitness capacities of power, strength, speed, agility and aerobic power (8).
13 Research to date in Australia (7,9,10) and the UK (30) has demonstrated increasing
14 anthropometric and fitness characteristics across youth annual-age categories and the
15 selection of earlier maturing players to talent development squads (i.e., Regional and
16 National, 28). Due to the relationship between maturation, anthropometry, fitness and
17 performance in youth rugby league this provides an ideal population to consider such
18 characteristics by maturity status.

19 Therefore, the primary purpose of the current study was to evaluate the
20 anthropometric and fitness characteristics within a rugby league academy by using maturity
21 status to determine annual categories instead of traditional chronological annual-age groups.
22 The second purpose was then to provide a longitudinal evaluation of the change in
23 anthropometric and fitness characteristics in relation to maturity status.

24

25

METHODS

1 **Experimental Approach to the Problem**

2 Rugby league players aged between 12.8 and 15.5 years from an English Super
3 League club's academy performed a testing battery at the start of each pre-season over a five-
4 year period (2007-2012). Players were assessed on anthropometric (height, body mass and
5 sum of four skinfolds), maturation (age at peak height velocity; PHV) and fitness (vertical
6 jump, medicine ball chest throw, 10m and 20m sprint and multi-stage fitness test)
7 characteristics. To evaluate anthropometric and fitness characteristics by maturity status,
8 players were assigned into annual maturity groups based on their maturity offset (Years from
9 PHV; YPHV) calculated by Mirwald et al. (20). Players that were assessed on consecutive
10 years were investigated for annual change in characteristics to examine longitudinal
11 development of characteristics based on maturity status.

12 **Subjects**

13 A total of 121 male, academy rugby league players (age = 14.40 ± 1.69 years) were
14 used in the study. Data was collected over a five-year period between 2007 and 2011 at the
15 Under 13s, 14s, 15s and 16s chronological annual-age categories. Players could potentially
16 join the academy programme at the Under 13s age category and leave the programme at the
17 Under 16s level (i.e., left the club or progressed to the Under 18s) but throughout this period
18 players were selected to or exited the programme at different stages. This resulted in a mixed
19 cross-sectional and longitudinal dataset whereby players were assessed between one and four
20 times. This data collection provided a total of 206 assessments with change in performance
21 data available on 85 occasions when players were assessed at consecutive age groups (i.e.,
22 Under 13s-14s).

23 Each player was categorised into one of six maturity offset groups (i.e., -2.5 YPHV (-
24 2.99 to -2.0), -1.5 YPHV (-1.99 to -1.0), -0.5 YPHV (-0.99 to 0.0), 0.5 YPHV (0.01 – 1.0),
25 1.5 YPHV (1.01 to 2.0) and 2.5 YPHV (2.01 – 3.0)). These categories were developed to

1 provide an annual category by maturity status instead of the traditional chronological annual-
2 age grouping. All experimental procedures were approved by the institutional ethics
3 committee with assent and parental consent provided along with permission from the rugby
4 league club.

5 **Procedures**

6 All pre-season testing was completed across two testing sessions in September each
7 year. All testing was undertaken by the lead researcher throughout the five-year period. A
8 standardised warm up including jogging, dynamic movements and stretches was used prior to
9 testing followed by full instruction and demonstrations of the assessments. Anthropometric
10 and fitness assessments were undertaken on all players within the academy, with the
11 procedures for each measure detailed below.

12 *Anthropometry:* Height and sitting height were measured to the nearest 0.1 cm using a
13 Seca Alpha stadiometer. Leg length was calculated by subtracting sitting height from standing
14 height. Body mass, wearing only shorts, was measured to the nearest 0.1 kg using calibrated
15 Seca alpha (model 770) scales. Sum of four skinfolds was determined by measuring four
16 skinfold sites (biceps, triceps, subscapular, suprailiac) using calibrated Harpenden skinfold
17 callipers (British Indicators, UK) in accordance with Hawes and Martin (12). Intraclass
18 correlation coefficients (ICC) and typical error measurements (TEM) for reliability of
19 skinfold measurements were $r = 0.954$ ($p < 0.001$) and 3.2%, respectively, indicating
20 acceptable reliability based on established criteria (i.e., $>.80$; 13).

21 *Maturity:* An age at peak height velocity (PHV) prediction equation (20) was used.
22 This involved a gender specific multiple regression equation including chronological age,
23 stature, sitting height, leg length, body mass and their interactions (24) being applied. The
24 equation in boys is $\text{Maturity Offset} = -9.236 + 0.0002708 \cdot \text{Leg Length} + \text{Sitting Height}$
25 $\text{interaction} - 0.001663 \cdot \text{Age} + \text{Leg Length interaction} + 0.007216 \cdot \text{Age and Sitting Height}$

1 interaction + 0.02292.Weight by Height ratio (20). The prediction equation has a 95%
2 confidence interval for boys of ± 1.18 years (20) and relationships with skeletal age have been
3 shown to be strong (i.e., $r=0.83$; 17). Maturity offset was determined by subtracting age at
4 PHV from chronological age and then allowed individuals to be assigned to a maturity offset
5 group.

6 *Lower-body Power:* A countermovement jump with hands positioned on hips was
7 used to assess lower body power via a just jump mat (Probotics, Hunstville, AL, USA). Jump
8 height was measured to the nearest 0.1 cm from the highest of three attempts (14). The ICC
9 and TEM for the vertical jump was $r = 0.903$ ($p < 0.001$) and 2.9%, respectively.

10 *Upper-body Power:* A 2 kg medicine ball (Max Grip, China) chest throw power (26).
11 Participants threw the ball horizontally as far as possible while seated with their back against
12 a wall. Distance was measured to the nearest 0.1m from where the ball landed to the wall with
13 the highest of three trials used as the score. The ICC and TEM for the medicine ball chest
14 throw was $r = 0.965$ ($p < 0.001$) and 0.6%, respectively.

15 *Speed:* Running speed was assessed over 10 m and 20 m using timing gates (Brower
16 Timing Systems, IR Emit, Draper, UT, USA). Participants were instructed to start in their
17 own time from a standing start 0.5 m behind the initial timing gate. Time was recorded to the
18 nearest 0.01s from the best of three attempts. ICC and TEM of the 10 m and 20 m sprints
19 were $r = 0.812$ ($p < 0.001$), 7.8% and $r = 0.852$, 4.5%, respectively.

20 *Endurance:* The multistage fitness test (MSFT; 23) was used to assess endurance
21 performance. Players were required to run 20 m shuttles, keeping to a series of beeps.
22 Running speed increased progressively until the players reached volitional exhaustion. Total
23 distance covered to the nearest 20 m was used to assess endurance performance. The ICC and
24 TEM for the MSFT has been reported as $r = 0.90$ ($p < 0.001$) and 3.1% (6).

25 *Statistical Analyses*

1 All analyses were conducted using SPSS 19.0 with mean and standard deviation (SD)
2 scores calculated for all dependant variables (i.e., anthropometric and fitness characteristics)
3 at each maturity offset group (i.e., -2.5 YPHV). Results are presented cross-sectionally by
4 each maturity group and longitudinally by analysing the change in performance between
5 assessments. MANOVA analyses were used to determine if differences existed between
6 dependant variables and the change in dependant variables between each maturity offset
7 group. A Bonferroni post-hoc analysis was used to determine where any significant
8 differences occurred. Significance levels were set at $p < 0.05$ with effect sizes (η^2) also
9 calculated.

11 RESULTS

12 Table 1 shows the anthropometric and fitness characteristics of all academy rugby
13 league players according to maturity offset group (i.e., -2.5 YPHV). MANOVA analyses
14 identified significant main effects for maturity offset group ($F_{5, 202} = 15.72$, $p < 0.001$, $\eta^2 = 0.47$,
15 $1 - \beta = 1.00$) with a significant large difference found across the groups for all variables with
16 univariate analyses presented in Table 1. Post-hoc analysis found chronological age was
17 significantly greater across the maturity offset groups except between -2.5 and -1.5 and
18 between 1.5 and 2.5 YPHV. Height, sitting height and body mass was also greater in the more
19 mature groups with skinfolds significantly greater in the 1.5 and 2.5 groups compared to the
20 other maturity offset groups.

21 Sprint speed was greater across the maturity groups, which showed significance
22 between -1.5 YPHV and the four greater maturity offset groups. Vertical jump performance
23 was also greater across the maturity offset groups with significance only demonstrated
24 between the -2.5 and -1.5 and 0.5 and 1.5 YPHV groups. Medicine ball chest throw was
25 significantly greater across the maturity offset groups. For MSFT distance there was no

1 significant difference between any maturational groups with the -2.5 YPHV group actually
2 the covering the greatest distance.

3 ***Insert Table 1 near here***

4 Table 2 shows the anthropometric and fitness changes with maturation over time.
5 MANOVA analyses identified a significant main effect for change in performance by
6 maturity offset group ($F_{4, 81} = 1.91$, $p=0.002$, $\eta^2=0.20$, $1-\beta = 0.99$) demonstrating that change
7 in performance was related to maturity status. Significant differences between maturity offset
8 groups for specific variables were found for height ($F_{4, 81} = 13.04$, $p<0.001$, $\eta^2=0.41$, $1-\beta =$
9 1.00) and sitting height ($F_{4,81} = 15.72$, $p=0.009$, $\eta^2=0.16$, $1-\beta = 0.98$) with the change between
10 the -1.5 to -0.5 and -0.5 to 0.5 YPHV groups significantly greater than the changes that
11 occurred between the 0.5 to 1.5 and 1.5 to 2.5 YPHV groups. No other significant differences
12 in change in performance were identified for any other variable due to the magnitude of
13 variation in the change in anthropometric and fitness characteristics.

14 ***Insert Table 2 near here***

15

16

DISCUSSION

17 The aims of the present study were to firstly evaluate the anthropometric and fitness
18 characteristics of junior rugby league players by using maturity status to determine annual
19 categories instead of traditional chronological annual-age groups and secondly to
20 longitudinally evaluate the change in performance in relation to maturation during the
21 adolescent period (Under 13s - 16s). Findings identified anthropometric characteristics were
22 greater as maturation increased across the six maturity offset groups with significant
23 differences identified for the change in height and sitting height between the maturity groups
24 with greater growth apparent at around PHV. For fitness characteristics, speed and lower and
25 upper body power developed with maturity status whereas maturity status had no effect on

1 endurance performance. No significant differences were identified for the change in fitness
2 performance across the maturity groups due to the magnitude of variation shown.

3 Cross-sectional examinations of chronological age, age at PHV and anthropometric
4 characteristics across the maturity offset group's revealed significant interactions.

5 Chronological age was greater and age at PHV was lower as maturity increased. This would
6 be expected as these variables contribute to the YPHV variable used to determine maturity
7 offset within this study and previous research (19,24). Therefore, using YPHV (i.e., maturity
8 offset) as an indicator of maturation includes both the assessment of chronological age and
9 maturation (i.e., age at PHV) providing an alternative to traditional chronological annual-age
10 group classifications. Height, sitting height and body mass were all significantly greater
11 across the maturity offset groups with the more mature players significantly taller and heavier
12 than the less mature players (e.g., Height, -1.5 YPHV = 154.6 ± 6.7 , 1.5 YPHV = 176.5 ± 4.7
13 cm). Findings are expected as these characteristics contribute to the prediction of age at PHV
14 (20), have been demonstrated to be strongly correlated to maturation (e.g., $p < 0.001$; 30) and
15 are related to the normal adaptations of growth, maturation and development during
16 adolescence (17). Sum of four skinfolds were significantly greater in the more mature players
17 (e.g., 1.5 YPHV = 38.9 ± 13.2 mm) compared to less mature players (e.g., -1.5 YPHV = 29.0
18 ± 4.4 mm). During adolescence, fat mass remains reasonably stable (4) with these findings
19 demonstrating that the more mature players selected to the academy possess greater body fat.
20 A possible explanation for this may be that earlier maturing players may have been selected to
21 the academy due to size advantages, in which previous research (28) highlighted increased
22 sum of skinfolds when earlier maturing forwards were compared to later maturing backs.
23 Therefore coaches may select players based on size and maturation, which may be
24 advantageous for forwards positions in rugby due to their game demands (i.e., physical
25 collisions and tackles).

1 Current findings demonstrated significant differences across maturity offset groups for
2 all fitness characteristics. Generally, fitness performance was greater in the more mature
3 groups for sprint speed (e.g., 20m sprint -2.5 YPHV = 1.98 ± 0.07 , 2.5 YPHV = 1.84 ± 0.07
4 s), vertical jump (e.g., -2.5 YPHV = 35.4 ± 4.2 , 2.5 YPHV = 42.8 ± 4.9 cm) and medicine
5 ball throw (e.g., -2.5 YPHV = 3.5 ± 0.4 , 2.5 YPHV = 6.3 ± 0.7 m) but not MSFT distance
6 (e.g., -2.5 YPHV = 1872 ± 186 , 2.5 YPHV = 1656 ± 251 m). These findings support previous
7 research that maturity is generally related to sprint and explosive performance (i.e., medicine
8 ball throw and vertical jump) during adolescence (18,33), which occurs due to increased
9 testosterone (17), increased muscle volume and size (27) and qualitative changes of the
10 muscle (i.e., contractile properties; 35). However, findings for endurance contradict existing
11 literature (18,33) and may be apparent due to differences in the playing positions (i.e.,
12 forwards have lower endurance performance than backs) amongst the maturity offset groups,
13 which is apparent in junior (28) and senior (9) rugby league players. The fact that significant
14 differences were not exclusively identified across all the maturity offset groups (e.g., vertical
15 jump no significant difference between -1.5 YPHV and 2.5 YPHV) support the notion that
16 advanced maturation is not always associated with better performance (28). The increase in
17 sum of four skinfolds (i.e., higher body fat percentage) with increasing maturity offset group
18 may have implications for fitness performance in the current sample due to the negative
19 association between skinfolds and physical performance, previously identified (28). This
20 finding suggests that skinfolds should be monitored regularly during adolescence to assess
21 body fat percentage, with training and nutritional interventions used appropriately to control
22 for excessive skinfolds that could negatively affect fitness performance.

23 Longitudinal examinations of change in characteristics within adolescent athletes are
24 limited (22), especially considering maturation (31). Current findings demonstrated
25 significant differences in the change in height and sitting height between the less and more

1 mature players as would be expected in relationship to age at PHV. These findings
2 demonstrate that monitoring height during adolescence should be considered in relation to
3 maturational status to understand an individual's potential growth. No significant differences
4 between maturity offset groups were identified for the change in performance for any fitness
5 variable. This is due to the large variability in the magnitude of change between maturity
6 groups (e.g., 20m sprint, -1.5 to -0.5 YPHV = -0.14 ± 0.12 s) demonstrating large individual
7 changes in fitness during adolescence, which may be related to changes in growth and
8 training status. Sprint speed improvements were increased around PHV, which may be related
9 to factors such as increased muscle mass and changes in the muscle-tendon architecture (36).
10 However, current findings contradict reports (22) that sprint performance is reduced leading
11 up to PHV. These longitudinal findings provide comparative data for the expected change in
12 performance in relation to maturity and provide an alternative to previous longitudinal
13 research (6,36), which use chronological annual-age groups. Such data could be applied to
14 estimate potential performance improvements based on current performance levels or used to
15 determine if young athletes are improving at an expected rate.

16 In conclusion, this study utilised a unique approach to classify anthropometric and
17 fitness characteristics into annual maturity categories, using a maturity offset (i.e., YPHV),
18 instead of traditional chronological annual-age grouping. The comparative data for
19 characteristics generally demonstrates an improvement in both anthropometric and fitness
20 measures in line with maturity, however some characteristics (i.e., MSFT distance) did not
21 follow this path suggesting that advanced maturation does not always result in superior
22 performance. These findings suggest that categorising players by maturity could be an
23 appropriate alternative or additional assessment method for evaluating player performance
24 alongside chronological annual-age categories, especially within adolescent athletes. The
25 longitudinal changes in performance demonstrate significant increases in height around age at

1 PHV with no further significant differences identified due to the magnitude of variation in
2 performance changes. Longitudinal monitoring should therefore be applied to allow current
3 performance and progress to be tracked to assist in identification, development and coaching
4 practices.

5

6

PRACTICAL APPLICATIONS

7 Due to the limitations of chronological annual-age grouping, considering maturity in
8 the evaluation of performance within adolescent athletes has recently been recommended (15,
9 34). National governing bodies, coaches, administrators and parents should assess and
10 consider maturation in the assessment and evaluation of youth athletes with YPHV (i.e.,
11 maturity offset) a possible alternative or additional method to chronological age for
12 classifying youth athletes. This approach would allow a more detailed assessment of an
13 athlete's current performance level, therefore assisting talent identification and development
14 processes alongside monitoring training adaptations. Measuring player characteristics and
15 performance by maturity offset would allow comparisons to be made in terms of biological
16 development instead of chronological age categories whereby differences in biological
17 maturation can be extensive (e.g., comparison of an early maturing, relatively older individual
18 vs a later maturing, relatively younger player). Likewise, comparing players by maturation
19 may reduce the emphasis placed on physical performance and size during selection, which has
20 resulted in relative age effects and maturational biases within youth rugby league (29) and
21 other sport contexts (e.g., soccer, 18). Lastly, tracking physical characteristics longitudinally
22 over time would assist in selection and development processes to attempt to differentiate
23 between current performance and potential for future development (32).

24

REFERENCES

1. Anderson, M, Hopkins, W, Roberts, A, and Pyne, D. Ability of test measures to predict competitive performance in elite swimmers. *J Sports Sci* 26: 123-130, 2008.
2. Armstrong, N, Welsby, JR, and Kirby, BJ. Peak oxygen uptake and maturation in 12 year olds. *Med Sci Sport Exerc* 30: 165-169, 1998.
3. Baxter-Jones, ADG. Growth and development of young athletes: Should competition levels be age related. *Sports Med* 20: 59-64, 1995.
4. Baxter-Jones, ADG, and Sherar, LB. Growth and Maturation. In: Armstrong N (eds). *Paediatric Exercise Physiology*. Elsevier, USA. 2007; 1-26.
5. Cogley, S, Baker, J, Wattie, N, and McKenna, J. Annual age-grouping and athlete development: A meta-analytical review of relative age effects in sport. *Sports Med* 39: 235-256, 2009.
6. Elferink-Gemser, MT, Visscher, C, Lemmink, KAPM, and Mulder, T. Multidimensional performance characteristics and standard of performance in talented youth field hockey players: A longitudinal study. *J Sports Sci* 25: 481-489, 2007.
7. Gabbett, TJ. A comparison of physiological and anthropometric characteristics among playing positions in junior rugby league players. *Br J Sports Med* 39: 675-680, 2005.
8. Gabbett, TJ. Applied Physiology of Rugby League. *Sports Med* 38: 119-138, 2008.
9. Gabbett, TJ, Kelly, J, and Pezet, T. A comparison of fitness and skill among playing positions in sub-elite rugby league players. *J Sci Med Sport* 11: 585-592, 2007.
10. Gabbett, TJ, Jenkins, DG, and Abernethy, B. Physiological and anthropometric correlates of tackling ability in junior elite and subelite rugby league players. *J Strength Cond Res*. 24: 2989-95, 2011.

- 1 11. Gravina, L, Gil, SM, Ruiz, F, Zubero, J, Gil, J, and Irazusta, J. Anthropometric and
2 physiological differences between first team and reserve soccer players aged 10-14
3 years at the beginning and end of the season. *J Strength Cond Res* 22: 1308-1314,
4 2008.
- 5 12. Hawes, MR, and Martin, AD. Human Body Composition. In: Eston R, Reilly T
6 (eds). *Kinanthropometry and exercise physiology laboratory manual: Tests,*
7 *procedures and data second edition.* Volume 1: Anthropometry. London, Routledge.
8 7-43. 2001
- 9 13. Hopkins, WG. Measures of reliability in sports medicine and science. *Sports Med*
10 30, 1-15, 2000.
- 11 14. Hunter, JP, and Marshall, RN. Effects of power and flexibility training on vertical
12 jump technique. *Med Sci Sports Exerc* 34: 478-486, 2002.
- 13 15. Lloyd, RS, Oliver, JL, Faigenbaum, AD, Myer, GD, and De Ste Croix, MB.
14 Chronological age vs. biological maturation: Implications for exercise programming
15 in youth. *J Strength Cond Res* 28: 1454-1464, 2014.
- 16 16. Malina, RM. Physical Growth and Biological Maturation of Young Athletes. *Exerc*
17 *Sport Sci Rev* 22: 389-433, 1994.
- 18 17. Malina, RM, Bouchard, C, and Bar-Or, O. *Growth, Maturation, and Physical*
19 *Activity.* (2nd ed.). United States of America; Human Kinetics. 2004.
- 20 18. Malina, RM, Eisenmann, JC, Cumming, SP, Ribeiro, B, and Aroso, J. Maturity-
21 associated variation in the growth and functional capacities of youth football (soccer)
22 players 13-15 years. *Eur J App Physiol* 91: 555-562, 2004.
- 23 19. Malina, RM, Coelho Silva, MJ, Figueiredo, AJ, Carling, C, and Beunen GP.
24 Interrelationships among invasive and non-invasive indicators of biological
25 maturation in adolescent male soccer players. *J Sports Sci* 30: 1705-1717, 2012.

- 1 20. Mirwald, RL, Baxter-Jones, ADG, Bailey, DA, and Beunen, GP. An assessment of
2 maturity from anthropometric measurements. *Med Sci Sport Exerc* 34: 689-694,
3 2002.
- 4 21. Nikolaidis, PT, Ziv, G, Arnon, N, and Lidor, R. Physical characteristics and
5 physiological attributes of female volleyball players--the need for individual data.
6 *J Strength Cond Res* 26: 2547-57, 2012.
- 7 22. Philippaerts, RM, Vaeyens, R, Janssens, M, Van Renterghem, B, Matthys, D, Craen,
8 R, Bourgois, J, Vrijens, J, Beunen, G, and Malina, RM. The relationship between
9 peak height velocity and physical performance in youth soccer players. *J Sports Sci*
10 24: 221-230, 2006
- 11 23. Ramsbottom, R, Brewer, J, and Williams, C. A progressive shuttle run test to
12 estimate maximal oxygen uptake. *Br J Sports Med* 22: 141-144, 1988.
- 13 24. Sherar, LB, Baxter-Jones, ADG, Faulkner, RA, and Russell, KW. Do physical
14 maturity and birth date predict talent in male youth ice hockey players? *J Sports Sci*
15 25: 879-886, 2007.
- 16 25. Smart, DJ, and Gill, ND. Effects of an off-season conditioning programme on the
17 physical characteristics of adolescent rugby union players. *J Strength Cond Res* 27,
18 708-717, 2013.
- 19 26. Stockbrugger, BA, and Haennel, RG. Contributing factors to performance of a
20 medicine ball explosive power test: a comparison between jump and non jump
21 athletes. *J Strength Cond Res* 17: 768-774, 2003.
- 22 27. Temfeno, A, Hugues, J, Chardon, K, Mandengue, SH, and Ahmaidi S. Relationship
23 between vertical jumping performance and anthropometric characteristics during
24 growth in boys and girls. *Eur J Pediatr* 168: 498-503, 2009.

- 1 28. Till, K, Cogley, S, O'Hara, J, Cooke, C, and Chapman, C. Anthropometric,
2 physiological and selection characteristics in high performance UK junior Rugby
3 League players. *Talent Dev Excellence* 2: 193-207, 2010.
- 4 29. Till, K, Cogley, S, Wattie, N, O'Hara, J, Cooke, C, and Chapman, C. The
5 prevalence, influential factors and mechanisms of relative age effects in UK Rugby
6 League. *Scand J Med Sci Sports* 20: 320-329, 2010.
- 7 30. Till, K, Cogley, S, O'Hara, J, Brightmore, A, Chapman, C, and Cooke, C. Using
8 anthropometric and performance characteristics to predict selection in junior UK
9 rugby league players. *J Sci Med Sport* 14: 264-269, 2011.
- 10 31. Till, K, Cogley, S, O'Hara, J, Chapman, C, and Cooke, C. Considering maturation
11 and relative age in the longitudinal evaluation of junior rugby league players. *Scand*
12 *J Sci Med Sport* 2013; doi: 10.1111/sms.12033
- 13 32. Till, K, Cogley, S, O'Hara, J, Chapman, C, and Cooke, C. An individualized
14 longitudinal approach to monitoring the dynamics of growth and fitness development
15 in adolescent athletes. *J Strength Cond Res* 27: 1313–1321, 2013.
- 16 33. Vaeyens, R, Malina, RM, Janssens, M, Van Renterghem, B, Bourgois, J, Vrijens, J,
17 and Philippaerts, RM. A multidisciplinary selection model for youth soccer: the
18 Ghent Youth Soccer Project. *Br J Sports Med* 40: 928-934, 2006.
- 19 34. Vaeyens, R, Lenoir, M, Williams, AM, and Philippaerts RM. Talent identification
20 and development programmes in sport: Current models and future directions. *Sports*
21 *Med* 38: 703-714, 2008.
- 22 35. Van Praagh, E, and Dore, E. Short term muscle power during growth and maturation.
23 *Sports Med* 32: 701-728, 2002.
- 24 36. Williams, CA, Oliver, JL, and Faulkner, J. Seasonal monitoring of sprint and jump
25 performance in a soccer academy. *Int J Sports Physiol Perform* 6: 264-275, 2011.

1 **Table 1: Anthropometric and fitness characteristics by annual maturity offset group and associated univariate analyses representing group**
 2 **differences**

	Maturity Offset Group (YPHV)						MANOVA		
	-2.5	-1.5	-0.5	0.5	1.5	2.5	F	P	η^2
	(n=6)	(n=19)	(n=30)	(n=55)	(n=67)	(n=29)			
Age (years)	12.48 ± 0.38	13.30 ± 0.68	13.92 ± 0.86 ^{a,b}	14.61 ± 0.71 ^{a,b,c}	15.22 ± 0.58 ^{a,b,c,d}	15.63 ± 0.44 ^{a,b,c,d}	57.05	<0.001	0.59
Age at PHV (years)	14.72 ± 0.43	14.55 ± 0.72	14.23 ± 0.81	13.99 ± 0.62 ^b	13.76 ± 0.49 ^{a,b,c}	13.38 ± 0.39 ^{a,b,c,d}	13.69	<0.001	0.26
Height (cm)	147.1 ± 5.2	154.6 ± 6.7	164.7 ± 6.0 ^{a,b}	171.8 ± 4.6 ^{a,b,c}	176.5 ± 4.7 ^{a,b,c,d}	180.5 ± 5.5 ^{a,b,c,d}	102.07	<0.001	0.72
Sitting Height (cm)	72.7 ± 3.2	76.8 ± 3.5 ^a	82.3 ± 3.5 ^{a,b}	86.4 ± 2.6 ^{a,b,c}	89.8 ± 2.2 ^{a,b,c,d}	92.7 ± 2.5 ^{a,b,c,d,e}	139.06	<0.001	0.78
Body Mass (kg)	37.9 ± 2.5	45.9 ± 4.7	54.7 ± 6.4 ^{a,b}	65.8 ± 6.2 ^{a,b,c}	74.3 ± 8.1 ^{a,b,c,d}	83.2 ± 10.0 ^{a,b,c,d,e}	107.35	<0.001	0.73
Skinfolds (mm)	24.8 ± 3.8	29.0 ± 4.4	27.6 ± 4.6	31.9 ± 8.6	38.9 ± 13.2 ^{a,b,c,d}	47.1 ± 16.6 ^{a,b,c,d,e}	14.24	<0.001	0.26
10m Sprint (s)	1.98 ± 0.07	1.96 ± 0.06	1.89 ± 0.09 ^b	1.87 ± 0.10 ^b	1.84 ± 0.08 ^{a,b}	1.84 ± 0.07 ^{a,b}	9.25	<0.001	0.19
20m Sprint (s)	3.46 ± 0.08	3.44 ± 0.11	3.28 ± 0.15 ^b	3.22 ± 0.17 ^{a,b}	3.17 ± 0.13 ^{a,b,c}	3.15 ± 0.11 ^{a,b,c}	16.92	<0.001	0.30
Vertical Jump (cm)	35.4 ± 4.2	39.1 ± 4.2	41.1 ± 5.8	43.6 ± 5.2 ^{a,b}	43.4 ± 6.4 ^{a,b}	42.8 ± 4.9	4.26	0.001	0.10
Medicine Ball Throw (m)	3.5 ± 0.4	4.0 ± 0.5	4.8 ± 0.7 ^{a,b}	5.6 ± 0.6 ^{a,b,c}	6.0 ± 0.6 ^{a,b,c,d}	6.3 ± 0.7 ^{a,b,c,d}	69.20	<0.001	0.64
MSFT Distance (m)	1872 ± 186	1547 ± 267	1637 ± 223	1670 ± 303	1675 ± 275	1656 ± 251	2.27	0.044	0.05

3 ^aSignificantly different to -2.5 ($P<0.05$); ^bSignificantly different to -1.5 ($P<0.05$); ^cSignificantly different to -0.5 ($P<0.05$); ^dSignificantly different to 0.5
 4 ($P<0.05$); ^eSignificantly different to 1.5 ($P<0.05$).

1 **Table 2: Change in anthropometric and fitness characteristics between annual maturity offset groups**

	Change between maturity offset groups (YPHV)				
	-2.5 to -1.5	-1.5 to -0.5	-0.5 to 0.5	0.5 to 1.5	1.5 to 2.5
	(n=6)	(n=13)	(n=19)	(n=30)	(n=17)
Height (cm)	5.4 ± 2.9	7.3 ± 2.1	6.1 ± 2.3	3.4 ± 2.0 ^{a,b}	2.0 ± 0.8 ^{a,b}
Sitting Height (cm)	2.5 ± 1.6	4.0 ± 2.5	3.7 ± 2.0	3.0 ± 2.1	1.6 ± 1.0 ^{a,b}
Body Mass (kg)	5.8 ± 2.0	8.5 ± 2.2	9.4 ± 3.4	6.7 ± 3.5	5.7 ± 5.8
Skinfolds (mm)	2.5 ± 3.6	-1.0 ± 3.0	1.8 ± 3.3	2.4 ± 6.0	-2.2 ± 9.0
10m Sprint (s)	-0.03 ± 0.03	-0.07 ± 0.07	-0.05 ± 0.09	-0.06 ± 0.08	-0.03 ± 0.04
20m Sprint (s)	-0.04 ± 0.02	-0.14 ± 0.12	-0.13 ± 0.12	-0.10 ± 0.10	-0.07 ± 0.06
Vertical Jump (cm)	4.1 ± 3.0	3.0 ± 4.4	3.7 ± 4.1	2.8 ± 4.1	2.1 ± 3.1
Medicine Ball Throw (m)	0.55 ± 0.37	0.64 ± 0.35	0.81 ± 0.38	0.63 ± 0.45	0.53 ± 0.30
MSFT Distance (m)	15 ± 64	142 ± 192	111 ± 206	83 ± 250	35 ± 226

2 ^aSignificantly different to -1.5 to -0.5 YPHV ($P < 0.01$); ^bSignificantly different to -0.5 to 0.5 YPHV ($P < 0.01$)

1 37.
2