



LEEDS
BECKETT
UNIVERSITY

Citation:

O'Mahoney, L and Stavropoulos-Kalinoglou, A and Birch, K and Ajjan, R and Price, OJ and Campbell, M (2019) Vascular Structure and Functional Responses to Consecutive High-Fat Feeding between Insulin Treatment Regimens in Adults with Type 1 Diabetes and Matched Controls. *Diabetes*, 68 (S1). ISSN 0012-1797 DOI: <https://doi.org/10.2337/db19-799-P>

Link to Leeds Beckett Repository record:

<http://eprints.leedsbeckett.ac.uk/5794/>

Document Version:

Article

Published abstract

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Title: Vascular Structure and Functional Responses to Consecutive High-Fat Feeding between Insulin Treatment Regimens in Adults with Type 1 Diabetes and Matched Controls.

Authors: L O'Mahoney¹, A Stavropoulos-Kalinoglou¹, KM Birch², RA Ajjan², OJ Price¹, and MD Campbell^{2,3}

¹Institute for Sport, Physical Activity, and Leisure, Leeds Beckett University, Leeds, UK

²Multidisciplinary Cardiovascular Research Group, University of Leeds, Leeds, UK

³School of Food Science and Nutrition, University of Leeds, Leeds, UK

Abstract: Background Impaired vascular health is prevalent in type 1 diabetes (T1D); however, it remains unknown whether different insulin treatment regimens mediate indices of vascular structure or function.

Methods Sixteen individuals with T1D receiving either multiple daily injection therapy (MDI; n=8; age: 32±13years; BMI:26.0±5.9kg.m²; HbA1c:53.7±11.2mmol/mol [7.1±3.2%]) or continuous subcutaneous insulin infusion (CSII; n=8; age:35±18years; BMI:26.3±4.6kg.m²; HbA1c: 58.6±9.7mmol/mol [7.5±3.0%]) and ten matched controls (CON; age:31±13years; BMI: 24.3±2.9kg.m²) consumed two high fat (HF) meals at 4-hour intervals. Carotid artery intima-media thickness (CIMT) and flow mediated dilation (FMD) was assessed at baseline, with further FMD assessment at 3-hrs following the ingestion of each meal using high resolution B-mode ultrasound. Bolus insulin dose was standardised using the carbohydrate-counting method. **Results** CIMT was significantly higher in individuals with T1D compared to controls (p=0.039); treatment stratification within T1D revealed MDI mediated this effect (MDI vs. CON: p=0.049; CSII vs. CON: p=0.112). FMD remained unchanged following the first meal (p=0.204) but was significantly impaired following the second meal (p<0.001); post-hoc analysis revealed MDI mediated this effect of impaired FMD after the second meal (MDI vs. CON: p=0.048; CSII vs. CON: p=0.416). **Conclusions** Our findings indicate that patients treated with MDI therapy have higher CIMT (a structural marker of subclinical atherosclerosis) compared to controls but not CSII therapy. FMD was impaired following a second HF meal irrespective of a diabetes status. Considering the pre-existing heightened cardiovascular disease risk in T1D therapeutic strategies to reduce postprandial risk warrants further research.