



LEEDS
BECKETT
UNIVERSITY

Citation:

Gregg, E and Nicholson, G and Beggs, C and Bissas, A (2019) Is trunk strength associated with functional mobility in older women? In: International Society of Posture & Gait Research World Congress 2019, Edinburgh, Scotland.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/6062/>

Document Version:

Conference or Workshop Item (Published Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

IS TRUNK STRENGTH ASSOCIATED WITH FUNCTIONAL MOBILITY IN OLDER WOMEN?

Emily Gregg, Gareth Nicholson, Clive B. Beggs & Athanassios Bissas

Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, UK

BACKGROUND AND AIM: Age-related declines in muscle strength have been associated with reduced functional capacity, postural instability and increased risk of falls in older adults. Studies have generally focussed on the significance of lower limb strength, however more recently the importance of trunk muscle strength has been explored, which may have been previously overlooked (Granacher et al., 2013). A limited number of studies have investigated the relationship between trunk strength and function, but this was tested under isometric contractions (Shahtahmassebi et al., 2017). Therefore, the purpose of this study was to investigate the relationship between dynamic (concentric) trunk strength and functional mobility in older women.

METHODS: A sample of 40 older women (age: 68.9 ± 3.73 years) participated in this study. To assess functional mobility, participants completed the timed up and go (TUG), 30 second chair stand test (CST), and a timed stair climb task (ascent and descent). Both self-selected normal and fast gait speed (GS) were also recorded. Concentric trunk flexion and extension strength data were collected using an isokinetic dynamometer (Biodex, USA). Measurements were conducted in the seated-compressed position in line with the manufacturer's guidelines. Following five sub-maximal trials, peak torque was recorded during three maximal flexion and extension trials at two testing speeds ($20^\circ/\text{s}$ and $45^\circ/\text{s}$). Peak torque was then normalised to body mass. Pearson's correlations were performed to investigate relationships between trunk strength and functional mobility measures. The level of significance was set at $p \leq 0.05$.

RESULTS: A number of associations between trunk strength and functional mobility measures were revealed. Greater trunk extension strength was correlated with superior performance in the CST ($45^\circ/\text{s}$: $r=0.33$, $p=0.037$), stair ascent ($20^\circ/\text{s}$: $r=-0.32$, $p=0.043$; $45^\circ/\text{s}$: $r=-0.56$, $p<.001$) and stair descent tasks ($45^\circ/\text{s}$: $r=-0.47$, $p=0.002$). Trunk extension strength was also associated with GS during normal ($20^\circ/\text{s}$: $r=0.37$, $p=0.019$; $45^\circ/\text{s}$: $r=0.48$, $p=0.004$) and fast trials ($45^\circ/\text{s}$: $r=0.34$, $p=0.034$). Trunk flexion strength was correlated with TUG ($45^\circ/\text{s}$: $r=-0.42$, $p=0.007$), stair ascent ($20^\circ/\text{s}$: $r=-0.41$, $p=0.008$; $45^\circ/\text{s}$: $r=-0.47$, $p=0.002$) and stair descent times ($20^\circ/\text{s}$: $r=-0.44$, $p=0.005$; $45^\circ/\text{s}$: $r=-0.42$, $p=0.007$). Trunk flexion strength was also associated with normal ($20^\circ/\text{s}$: $r=0.55$, $p<.001$; $45^\circ/\text{s}$: $r=0.56$, $p<.001$) and fast GS ($20^\circ/\text{s}$: $r=0.32$, $p=0.047$; $45^\circ/\text{s}$: $r=0.34$, $p=0.031$).

CONCLUSIONS: These findings suggest that there is a moderate-strong relationship between muscle strength in the trunk area and a range of functional mobility measures in older women. These results have important practical implications which could be used to inform the inclusion of trunk exercises in interventions targeting functional mobility for older adults.

REFERENCES: Granacher et al. *Sports Med* (2013) 43:627-641; Shahtahmassebi et al. *Scientific reports* (2017) 7:10907