



LEEDS
BECKETT
UNIVERSITY

Citation:

Chantler, S and Wood-Martin, R and Holliday, A and Davison, G and Crabtree, DR and Readhead, C and Jones, B (2024) The Frequency and Severity of Gastrointestinal Symptoms in Rugby Players. International Journal of Sports Medicine. pp. 1-9. ISSN 1439-3964 DOI: <https://doi.org/10.1055/a-2206-4751>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/10542/>

Document Version:

Article (Accepted Version)

© 2024, Thieme. All rights reserved.

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

The frequency and severity of gastrointestinal symptoms in rugby players

Abstract

Assess self-reported frequency and severity of gastrointestinal symptoms (GIS) at rest and around rugby training and match play in male and female rugby union players. An online questionnaire was sent to registered rugby union players (sevens or fifteens). Thirteen GIS were assessed alongside perceptions of appetite around rugby and rest using Likert and visual analogue scales. Questions investigating a range of medical and dietary factors were included. 325 players (male n=271, female n=54) participated in the study. More frequent GIS (at least one GIS experienced weekly/more often) was reported by players at rest (n=203; 62%) compared to around rugby (n=154; 47%). Overall severity of GIS was low (mild discomfort), but a portion of players (33%) did report symptoms of moderate severity around rugby. Female players reported more frequent and severe symptoms compared to male counterparts ($p<0.001$). Self-reported appetite was significantly lower after matches compared to training. There were no dietary or medical factors associated with GIS severity scores. This study describes GIS characteristics in male and female rugby union players. Half of players experienced some form of GIS which may affect nutrition, training, or performance, thus should be a consideration for practitioners supporting this cohort.

Keywords: rugby, gastrointestinal symptoms, appetite, sport

The frequency and severity of gastrointestinal symptoms in rugby players

Introduction

Depending on the style and duration of the activity, 30-93% of athletes report a range of gastrointestinal symptoms (GIS) around exercise [1]. Different symptoms, often categorised into upper or lower gastrointestinal tract, can impair training, performance, or wellbeing [2]. The aetiology of GIS around exercise can be complex, with no single causal factor. Exercise-associated gastrointestinal syndrome highlights two mechanistic avenues for GIS [3]: the circulatory gastrointestinal and the neuroendocrine gastrointestinal pathways. Exercise intensity, duration, dietary intake, time of day, sleep, medication and the external environment may influence the experience of GIS in athletes [4–9].

Data describing the frequency and severity of GIS in athletes is predominantly from endurance exercise and there is limited research investigating GIS in team sports[10]. Rugby union is a high intensity intermittent team sport that includes specific risk factors which may predispose players to gastrointestinal distress and changes in appetite (e.g. high training volume, collisions, use of pain medication). Repeated training sessions with high intensity efforts and collisions contribute to the physical demands on players and reflect in their recovery and energy demands thus creating a unique team sport environment [11–13]. Food choices, food volume and energy balance may influence GIS [14,15] but gastrointestinal profiles have not been included in any investigations of dietary intakes in rugby players thus far.

Changes in appetite do not appear part of the traditional assessment of gastrointestinal distress around exercise, although loss of appetite may be a barrier to exogenous fuelling. Loss of appetite may be experienced concomitant to other GIS, such as nausea or bloating [14]. Appetite is likely critical in the post-exercise period for team sports when nutrition is prioritised as part of recovery and where high daily calorie intakes are necessary, as is the case in rugby [16]. The continued growth of women's rugby is an important consideration, as female athletes report higher levels of GIS compared to males in endurance athletes but is unknown in team sport athletes [17,18].

Therefore, the aim of this study was to establish self-reported frequency and severity of GIS and appetite responses associated with rugby and rest in rugby union players.

Materials and Methods

Participants

Male and female senior rugby union players, from both fifteens and sevens, were invited to participate in the study. Online anonymous questionnaires were sent via coaches or national governing body. Coaches were recruited through existing rugby networks across different levels of play. Multiple governing bodies were invited to participate. Those who participated distributed the

questionnaire via the lead nutritionist, the nutritionist at the club or the players association. The participant information sheet was shared prior to the study and this information and informed consent were included in the online questionnaire (Supplementary material). Informed consent was obtained from all participants. Ethics approval was granted by the Ethics Advisory Committee at an institution affiliated with one of the authors. Additional permissions were obtained from the Irish and South African rugby federations.

Questionnaire

An online questionnaire (Qualtrics^{XM}), written in the English language, was used for convenience and effective distribution (Supplementary material). Self-reported characteristics (age, body mass, height, playing level, primary position of play, training hours/week) were included to define participants [19]. Modified Likert scales were used to assess the frequency and severity of thirteen GIS [20,21] at rest and around rugby. Rest and around rugby were not pre-defined for participants and left to the participants interpretation. The lack of specific definition to the rugby and rest environments was identified as a limitation after the study. Perceptions of appetite pre and post-training and pre and post-match were included as a separate GIS using visual analogue scales (VAS) [22,23]. Two items relating to constructs of appetite: feelings of hunger and desire to eat were included. A combination of dichotomous and ranking questions were used to assess the various associated risk factors for GIS reported in other studies. Risk factors included antibiotic use, NSAIDs, fibre intakes (via fruit and vegetables), alcohol post-exercise, hydration pre-exercise, probiotics (in supplements), fermented foods and high protein intakes. These risk factors were drawn from a variety of research [24–29].

Statistical analyses

Statistical analyses were conducted using the Statistical Package for the Social Sciences software programme (SPSS, Version 26). Data are presented as median (interquartile range (IQR)). VAS data for perceptions of appetite are reported in arbitrary units (AU) rather than millimetres, due to screen/phone size. All GIS data were non-normally distributed. Wilcoxon rank tests were used to compare differences in GIS frequencies at rest and around rugby (separate GIS), as well as changes in GIS severity scores between rest and rugby conditions in all, male and female groups. Chi-squared tests were used to compare the frequencies of any GIS (weekly or more often) or GIS severity (>4) between rest and rugby conditions. Spearman rank-order correlation was used to analyse the relationship between GIS severity scores and associated risk factors mentioned above. Statistical significance was accepted at a level of $p < 0.05$. Graphs were created by the authors using R (version 4.2.0).

Results

Participant characteristics

A total of 325 players (271 male and 54 female) completed the questionnaire. Characteristics of the sample are presented in Table 1. Half the sample were International/National level [19]. None of the characteristics (e.g., body mass, age, training hours/week) had any significant relationship with GIS frequency or severity ($p>0.05$).

Insert Table 1

Gastrointestinal symptom frequency and severity

At least one GIS (experienced weekly or more often) was reported by 62% (n=203) of players at rest and by 47% (n=154) of players around rugby (Table 2, $p<0.001$). In a separate question, almost half of the players (48%) reported that they 'probably' or 'definitely' experienced more severe GIS around matches compared to training (any symptom), however only diarrhoea ($p<0.001$) and urgent need to defecate ($p=0.48$) had a higher frequency reported around rugby compared to rest for females, whilst diarrhoea ($p=0.05$) had a higher frequency around rugby compared to rest for males (Table 2). Other GIS were reported as higher at rest compared to rugby conditions.

Insert Table 2

Median rating of severity of individual symptoms ranged between 1-3 (no discomfort to mild discomfort) (Figure 1). However, 42% and 33% of players reported at least one symptom with moderate discomfort or worse (≥ 4) at rest or around rugby, respectively (Table 2).

Insert Figure 1

Combined severity scores for upper, lower and total GIS were significantly higher at rest compared to around rugby for the whole group (Table 3, $p<0.001$, $p=0.002$, $p<0.001$ for upper, lower, and total GIS scores respectively).

Insert Table 3

Changes in appetite

Sixty percent of players reported lower feelings of hunger and desire to eat after a match compared with before a match, while 40% reported lower feelings of hunger and desire to eat after training compared with before training (Figure 2). After matches, 39% of all participants reported that their loss of appetite resolved within an hour, while 46% reported they needed one to two hours for their appetite to return to normal.

Insert Figure 2

Sex-based differences

Female participants reported higher frequency and severity of GIS (Table 2, Table 3, Figure 1) and lower feelings of hunger and desire to eat around rugby compared to males (Figure 2). Male players reported lower total severity of GIS around rugby compared to rest ($p<0.001$), whereas female

players reported no difference between severity scores around rugby compared to rest (Table 3). The median scores translate as minor discomfort (rating of 2), apart from diarrhoea (around rugby) which had a median severity of mild discomfort (rating of 3). In male players, median feelings of hunger after matches were lower than after training (35 vs. 45 AU, $p=0.54$, Figure 2) as was desire to eat (34 vs. 45 AU, $p=0.003$, Figure 2). which was similar for female players (feelings of hunger; 35 vs 53 AU, $p<0.001$, desire to eat; 31 vs. 52 AU, $p<0.001$, Figure 2).

Risk Factors for GIS

None of the risk factors assessed (medication or dietary) in the questionnaire were found to be associated to the frequency or severity of GIS scores (Table 4, all $p>0.05$).

Insert Table 4

Discussion

This is the first study of self-reported GIS characteristics and perceptions of appetite in high level rugby union players. Approximately half of players experienced at least one symptom weekly or more often at rest or around rugby. Players reported more frequent and severe GIS at rest compared to rugby for the majority of symptoms, but the overall severity was low. Female players experienced more frequent and severe GIS compared to their male counterparts. Measures of appetite were lower post-match compared to post-training, with females reporting lower levels of hunger and desire to eat compared to males. There were no associations between the frequency or severity of GIS and any of the external factors assessed. GIS profiles in rugby players should be assessed in order to highlight risk factors and possible intervention options for those who are affected.

The frequency of reported GIS is lower than in a previous mixed sport sample (86%) [30] but aligns with findings from various endurance studies [1,31]. The range of frequencies across individual symptoms (4-48%) in this study mirrors those previously reported where certain GIS (e.g. bloating) are more common than others [30,32]. Diarrhoea and urge to defecate were two GIS that were elevated around rugby compared to rest. This may be as a response to sympathetic activation that may influence specific lower GIS [33]. Rugby may expose players to higher rates of psychological stress due to concerns about performance. Psychological stress has been correlated with GIS during endurance running [34]. Unfortunately, there is no published research on the psychological stress state in rugby players but a 'nervous tummy' is supported anecdotally by players and in recent studies in runners [35] and team athletes [10].

The lower overall frequency and severity of GIS around rugby compared to rest (47% vs. 62%) may be influenced by training or dietary adaptations. Players involved in a fulltime rugby programmes (~50% of participants) will participate in a range of training modalities [36]. Rugby training sessions can be 45-60 minutes [36,37]; a duration that may be too short to elicit significant GIS but may still

contribute to positive adaptations [38]. Hypothetically, repeated rugby sessions may improve gastrointestinal tolerance over time, via decreasing levels of splanchnic hypoperfusion in response to increasing cardiovascular fitness. Different exercise modalities and different styles of training (fifteens vs. sevens) may also alter the magnitude of the GI response [29,38–40]. In this current study, participants did not report any perceived difference in GIS between different training sessions, including sessions with contact-based drills. Full contact-based drills or collisions in match play increase the metabolic requirements of players [36,41], but seem not to influence the experience of GIS reported here. This may also be in part to the limits placed on contact-based drills during training over time[42]. Further, there may be a secondary improvement mediated via the microbiome [7,43]. Specific symptom frequency, for example flatulence, may be influenced or modulated by certain bacterial species [44]. Athletes have been found to have altered microbiome profiles compared to non-athletes, including rugby players [45–47]. However, the impact of these differences in microbiome on GIS is unknown. A third of players still reported GIS of moderate severity or worse around rugby which may cause discomfort around training or rugby matches, possibly negatively influencing performance [48]. The experience of athletes of GIS in relation to match day performance will be useful to explore further.

Dietary adaptations may include pre-training dietary exclusions [49], or training in a euhydrated, healthy state [50]. Participants in this study at elite/sub-elite level may approach rugby training with specific dietary habits learned through experience, limiting risk of GI distress. This may go alongside adaptations to higher carbohydrate intakes. High volumes of carbohydrate fluid given repeatedly over two weeks has been shown to reduce GIS during exercise [51], while high carbohydrate intakes may also induce an increased content of glucose transporter protein (GLUT4 and GLUT5) in the gastrointestinal tract [52]. Conversely, higher volumes of food eaten to meet higher caloric requirements may aggravate specific GIS such as flatulence and bloating at rest [43,53]. Therefore, dietary exposures aligned with a training programme may minimise GIS around rugby. Monitoring GIS frequency and severity reported over a preseason and during a season may give more insight around possible training and dietary adaptations.

This study showed GIS to be more common in female compared to males, with sex-based changes in colonic transit time and visceral hypersensitivity being proposed amongst the possible mechanisms [17,18,48]. Menstrual phase may affect female rugby players, who previously reported abdominal pain, nausea or cramping as common GIS that negatively affect training during their menstrual cycle [54]. Unfortunately, neither menstrual phase nor contraceptive use were assessed in this study in the female players. As there appears to be no sex-based difference seen in the gastrointestinal endothelial response to exercise [55], an endocrine related mechanism may be more likely. Female players may need screening for GIS and changes in relation to their menstrual cycle phase. Female participants in this study, although from national representative teams, may train and compete in part-time environments and have less nutrition support compared to males,

as well as lower training ages. This may also differentiate any training and dietary associated adaptations mentioned above.

Self-reported measures of appetite (feelings of hunger, desire to eat) were lower post-match compared to post-training and most players reported this took an hour or more to resolve. This coincides with the clearance time for markers of gut endothelial damage, splanchnic reperfusion [56,57] and metabolites such as lactate [58,59]. The reported suppression of appetite seen here with rugby is concordant with appetite responses after continuous [60,61]; and intermittent high-intensity exercise in laboratory studies of non-athletic populations [62,63]. Previous data has not demonstrated any differences in appetite responses to exercise in male and female athletes [64,65]. These data may reflect both the physiological intensity and psychological stress of match scenarios as mentioned previously. More investigation into the differences between sevens and fifteens players would be useful, but were not within the scope of this study. Appetite, as a GIS, requires practical nutrition support; however, validated nutritional strategies for increasing appetite and promoting feeding post-exercise have yet to be established.

Based on the current literature, this is the first study to describe self-reported rates of NSAIDs use in rugby players. Acute NSAID use has been shown to increase gut cell damage with and without exercise [66–68]. Chronic NSAID use has been associated to higher levels of GIS in non-athletes [69], but there is limited consensus on the impact of chronic NSAID use on GIS and performance in athletes [70]. Rates of reported NSAID use were lower than studies in endurance runners (57–60% of race finishers) [2,71] and collegiate American football players (~50% players during the season) [72]. Lower levels seen here may reflect improved awareness in players and stricter protocols around NSAID prescription due to publicised reports of now-retired rugby players describing high NSAID use for pain management and consequent GIS. Continued education to players of all levels around appropriate NSAID use will only benefit future gastrointestinal outcomes. A third of players reported antibiotics in the previous year, but this is difficult to contextualise with limited data on antibiotic use in team sport outside of major competitions [73]. Antibiotics have been shown to disrupt host microbiome, aggravate GIS (antibiotic associated diarrhoea) and recurrent use may impact long term gastrointestinal outcomes [74] and performance [75]. It would be important for future research to elucidate microbiome recovery in athletes post-antibiotics, or any concomitant strategies (e.g. probiotics) that may limit any detrimental effects of antibiotics on gastrointestinal health [76].

Dietary factors may influence GIS. Fruit and vegetable, alcohol, probiotic, and prebiotic intakes and pre-match hydration status were included in the questionnaire as these have been implicated in general gastrointestinal health and GIS around exercise [25,77–79]. Although there were no relationships found with GIS in this study, practitioners can still consider that these factors may play a role in GIS in individual cases. Investigation of dietary strategies used in endurance athletes to

proactively manage GIS has been published [80] which may give some insight into other dietary interactions.

Limitations

Using an online questionnaire has limitations. Although efforts were made to be comprehensive, superficial questions make some of the application of these data challenging. While rest and rugby were chosen to establish the change in environment, there may be differences in interpretation around when a player would transition from one to the other. More in depth studies should consider the differences between sevens and fifteens environments. There is currently no consensus on what frequency (e.g., weekly) or severity (e.g., moderate) of GIS is meaningful to athletes for wellbeing or performance. This is especially relevant for GIS like burping compared to diarrhoea, where performance associated impact may not be equivalent. This questionnaire was cross-sectional and some form of season-wide surveillance from a baseline would give more insight into the ability for the gut to adapt in contact-based, highly demanding sports. Ensuring that the current GIS questionnaires available is validated for team sports, as they have been done for endurance exercise, may be useful as there are limited data from team sports [48,81,82].

This study highlights self-reported GIS in high level male and female rugby players. Low severity GIS are common, but more so at rest. With limited data in team sport, these data may assist with the awareness for practitioners and development of interventions for the individuals who are affected. Future research focusing on the chronic impact of rugby training, NSAID use, collisions and dietary changes on the GIS profile over time (a season) would be useful to promote gastrointestinal health for wellbeing and performance.

References

- [1] De Oliveira EP, Burini RC, Jeukendrup A. Gastrointestinal complaints during exercise: Prevalence, etiology, and nutritional recommendations. *Sports Medicine* 2014; 44: 79–85. doi:10.1007/s40279-014-0153-2
- [2] Hoffman MD, Fogard K. Factors related to successful completion of a 161-km ultramarathon. *Int J Sports Physiol Perform* 2011; 6: 25–37. doi:10.1123/ijsp.6.1.25
- [3] Costa RJS, Snipe RMJ, Kitic CM, et al. Systematic review : exercise-induced gastrointestinal syndrome — implications for health and intestinal disease. *Aliment Pharmacol Ther* 2017; 246–265. doi:10.1111/apt.14157
- [4] Costa RJS, Gaskell SK, McCubbin AJ, et al. Exertional-heat stress-associated gastrointestinal perturbations during Olympic sports: Management strategies for athletes preparing and competing in the 2020 Tokyo Olympic Games. *Temperature* 2020; 7: 58–88. doi:10.1080/23328940.2019.1597676
- [5] Gaskell SK, Rauch CE, Parr A, et al. Diurnal versus Nocturnal Exercise-Impact on the Gastrointestinal Tract. *Med Sci Sports Exerc* 2020; doi:10.1249/MSS.0000000000002546
- [6] Pals KL, Chang RT, Ryan AJ, et al. Effect of running intensity on intestinal permeability. *J Appl Physiol* 1997; 82: 571–576. doi:10.1152/jappl.1997.82.2.571
- [7] Smith KA, Pugh JN, Duca FA, et al. Gastrointestinal pathophysiology during endurance exercise: endocrine, microbiome, and nutritional influences. *Eur J Appl Physiol* 2021;
- [8] Wilson PB. Associations between sleep and in-race gastrointestinal symptoms: An observational study of running and triathlon race competitors. *Sleep Science* 2020; 13: 293–297. doi:10.5935/1984-0063.20200029
- [9] Wilson PB, Ingraham SJ. Glucose-fructose likely improves gastrointestinal comfort and endurance running performance relative to glucose-only. *Scand J Med Sci Sports* 2015; 25: e613–e620. doi:10.1111/sms.12386
- [10] Wilson PB, Fearn R, Pugh J. Occurrence and Impacts of Gastrointestinal Symptoms in Team-Sport Athletes: A Preliminary Survey. *Clin J Sport Med* 2022; doi:10.1097/JSM.0000000000001113
- [11] Roe G, Till K, Darrall-Jones J, et al. Changes in markers of fatigue following a competitive match in elite academy rugby union players. *South African Journal of Sports Medicine* 2016; 28: 2. doi:10.17159/2078-516x/2016/v28i1a1411
- [12] Bradley WJ, Cavanagh BP, Douglas W, et al. Quantification of training load, energy intake, and physiological adaptations during a rugby preseason: A case study from an elite European rugby union squad. *J Strength Cond Res* 2015; 29: 534–544. doi:10.1519/JSC.0000000000000631
- [13] Costello N, Deighton K, Preston T, et al. Collision activity during training increases total energy expenditure measured via doubly labelled water. *Eur J Appl Physiol* 2018; 118: 1169–1177. doi:10.1007/s00421-018-3846-7

- [14] Costa RJS, Gill SK, Hankey J, et al. Perturbed energy balance and hydration status in ultra-endurance runners during a 24 h ultra-marathon. *British Journal of Nutrition* 2014; 112: 428–437. doi:10.1017/S0007114514000907
- [15] Wilson PB, Rhodes GS, Ingraham SJ. Saccharide Composition of Carbohydrates Consumed during an Ultra-endurance Triathlon. *J Am Coll Nutr* 2015; 34: 497–506. doi:10.1080/07315724.2014.996830
- [16] Costello N, Deighton K, Preston T, et al. Are professional young rugby league players eating enough? Energy intake, expenditure and balance during a pre-season. *Eur J Sport Sci* 2019; 19: 123–132. doi:10.1080/17461391.2018.1527950
- [17] Pugh JN, Lydon K, O'Donovan CM, et al. More than a gut feeling: What is the role of the gastrointestinal tract in female athlete health? *Eur J Sport Sci* 2021; 1–10. doi:10.1080/17461391.2021.1921853
- [18] Ter Steege RWF, Van Der Palen J, Kolkman JJ. Prevalence of gastrointestinal complaints in runners competing in a long-distance run: An internet-based observational study in 1281 subjects. *Scand J Gastroenterol* 2008; 43: 1477–1482. doi:10.1080/00365520802321170
- [19] McKay AKA, Stellingwerff T, Smith ES, et al. Defining Training and Performance Caliber: A Participant Classification Framework. *Int J Sports Physiol Perform* 2022; 17: 317–331. doi:10.1123/ijsp.2021-0451
- [20] Pugh JN, Sage S, Hutson M, et al. Glutamine supplementation reduces markers of intestinal permeability during running in the heat in a dose-dependent manner. *Eur J Appl Physiol* 2017; 117: 2569–2577. doi:10.1007/s00421-017-3744-4
- [21] Svedlund J, Sjodin I, Dotevall G. GSRS-A Clinical Rating Scale for Gastrointestinal Symptoms in Patients with Irritable Bowel Syndrome and Peptic Ulcer Disease. 1988
- [22] Blundell J, De Graaf C, Hulshof T, et al. Appetite control: Methodological aspects of the evaluation of foods. *Obesity Reviews* 2010; 11: 251–270. doi:10.1111/j.1467-789X.2010.00714.x
- [23] Flint A, Raben A, Blundell J, et al. Reproducibility, power and validity of visual analogue scales in assessment of appetite sensation in single test meal studies Obesity-associated arterial hypertension-pathophysiology and treatment View project Collaboration with NIHS on RNA-seq data View. Article in *International Journal of Obesity* 2000; 24: 38–48
- [24] Snipe RMJ, Costa RJS. Does the temperature of water ingested during exertional-heat stress influence gastrointestinal injury, symptoms, and systemic inflammatory profile? *J Sci Med Sport* 2018; 21: 771–776. doi:10.1016/j.jsams.2017.12.014
- [25] Costa RJS, Camões-Costa V, Snipe RMJ, et al. Impact of exercise-induced hypohydration on gastrointestinal integrity, function, symptoms, and systemic endotoxin and inflammatory profile. *J Appl Physiol* 2019; 126: 1281–1291. doi:10.1152/jappphysiol.01032.2018
- [26] Fujimori S, Gudis K, Sakamoto C. A review of anti-inflammatory drug-induced gastrointestinal injury: Focus on prevention of small intestinal injury. *Pharmaceuticals* 2010; 3: 1187–1201. doi:10.3390/ph3041187

- [27] Hsu YJ, Huang WC, Lin JS, et al. Kefir supplementation modifies gut microbiota composition, reduces physical fatigue, and improves exercise performance in mice. *Nutrients* 2018; 10. doi:10.3390/nu10070862
- [28] Shing CM, Peake JM, Lim CL, et al. Effects of probiotics supplementation on gastrointestinal permeability, inflammation and exercise performance in the heat. *Eur J Appl Physiol* 2014; 114: 93–103. doi:10.1007/s00421-013-2748-y
- [29] van Wijck K, Pennings B, van Bijnen AA, et al. Dietary protein digestion and absorption are impaired during acute postexercise recovery in young men. *Am J Physiol Regul Integr Comp Physiol* 2013; 304. doi:10.1152/ajpregu.00294.2012
- [30] Pugh JN, Fearn R, Morton JP, et al. Gastrointestinal symptoms in elite athletes: Time to recognise the problem? *Br J Sports Med* 2018; 52: 487–488. doi:10.1136/bjsports-2017-098376
- [31] Pugh JN, Kirk B, Fearn R, et al. Prevalence, severity and potential nutritional causes of gastrointestinal symptoms during a marathon in recreational runners. *Nutrients* 2018; 10. doi:10.3390/nu10070811
- [32] de Lira CAB, Viana RB, Mesquista KP, et al. Frequency and intensity of gastrointestinal symptoms in exercisers individuals at rest and during physical exercise: An internet-based survey. *Intest Res* 2019; 17: 537–545. doi:10.5217/IR.2018.00162
- [33] Brouns F, Beckers E. Is the Gut an Athletic Organ? Digestion, Absorption and Exercise. *Sports Medicine* 1993; 15: 242–257. doi:10.2165/00007256-199315040-00003
- [34] Wilson PB. Perceived life stress and anxiety correlate with chronic gastrointestinal symptoms in runners. *J Sports Sci* 2018; 36: 1713–1719. doi:10.1080/02640414.2017.1411175
- [35] Urwin CS, Main LC, Mikocka-Walus A, et al. The Relationship Between Psychological Stress and Anxiety with Gastrointestinal Symptoms Before and During a 56 km Ultramarathon Running Race. *Sports Med Open* 2021; 7. doi:10.1186/s40798-021-00389-5
- [36] Bradley WJ, Cavanagh BP, Douglas W, et al. Quantification of training load, energy intake, and physiological adaptations during a rugby preseason: A case study from an elite European rugby union squad. *J Strength Cond Res* 2015; 29: 534–544. doi:10.1519/JSC.0000000000000631
- [37] Morehen JC, Bradley WJ, Clarke J, et al. The assessment of total energy expenditure during a 14-day in-season period of professional rugby league players using the doubly labelled water method. *Int J Sport Nutr Exerc Metab* 2016; 26: 464–472. doi:10.1123/ijsnem.2015-0335
- [38] Stuempfle KJ, Hoffman MD. Gastrointestinal distress is common during a 161-km ultramarathon. *J Sports Sci* 2015; 33: 1814–1821. doi:10.1080/02640414.2015.1012104
- [39] Edwards KH, Ahuja KD, Watson G, et al. The influence of exercise intensity and exercise mode on gastrointestinal damage. *Applied Physiology, Nutrition, and Metabolism* 2021; 6: 1–6. doi:10.1139/apnm-2020-0883

- [40] Pfeiffer B, Stellingwerff T, Hodgson AB, et al. Nutritional intake and gastrointestinal problems during competitive endurance events. *Med Sci Sports Exerc* 2012; 44: 344–351. doi:10.1249/MSS.0b013e31822dc809
- [41] Costello N, Deighton K, Preston T, et al. Collision activity during training increases total energy expenditure measured via doubly labelled water. *Eur J Appl Physiol* 2018; 118: 1169–1177. doi:10.1007/s00421-018-3846-7
- [42] World Rugby. Contact load guidelines. 2021
- [43] Clarke SF, Murphy EF, O’Sullivan O, et al. Exercise and associated dietary extremes impact on gut microbial diversity. *Gut* 2014; 63: 1913–1920. doi:10.1136/gutjnl-2013-306541
- [44] Seo M, Heo J, Yoon J, et al. Methanobrevibacter attenuation via probiotic intervention reduces flatulence in adult human: A non-randomised paired-design clinical trial of efficacy. *PLoS One* 2017; 12: 1–20. doi:10.1371/journal.pone.0184547
- [45] Petersen LM, Bautista EJ, Nguyen H, et al. Community characteristics of the gut microbiomes of competitive cyclists. *Microbiome* 2017; 1–13. doi:10.1186/s40168-017-0320-4
- [46] Barton W, Penney NC, Cronin O, et al. The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level. *Gut* 2018; 67: 625–633. doi:10.1136/gutjnl-2016-313627
- [47] Chong PP, Chin VK, Looi CY, et al. The microbiome and irritable bowel syndrome - A review on the pathophysiology, current research and future therapy. *Front Microbiol* 2019; 10: 1–23. doi:10.3389/fmicb.2019.01136
- [48] Wilson PB, Fearn R, Pugh J. Occurrence and Impacts of Gastrointestinal Symptoms in Team-Sport Athletes: A Preliminary Survey. *Clinical Journal of Sport Medicine* 9900;
- [49] Parnell JA, Wagner-Jones K, Madden RF, et al. Dietary restrictions in endurance runners to mitigate exercise-induced gastrointestinal symptoms. *J Int Soc Sports Nutr* 2020; 17: 1–10. doi:10.1186/s12970-020-00361-w
- [50] Pugh JN, Impey SG, Doran DA, et al. Acute high-intensity interval running increases markers of gastrointestinal damage and permeability but not gastrointestinal symptoms. *Applied Physiology, Nutrition and Metabolism* 2017; 42: 941–947. doi:10.1139/apnm-2016-0646
- [51] Miall A, Khoo A, Rauch C, et al. Two weeks of repetitive gut-challenge reduce exercise-associated gastrointestinal symptoms and malabsorption. *Scand J Med Sci Sports* 2018; 28: 630–640. doi:10.1111/sms.12912
- [52] Horowitz M, Cunningham KM, Wishart JM, et al. The effect of short-term dietary supplementation with glucose on gastric emptying of glucose and fructose and oral glucose tolerance in normal subjects. *Diabetologia* 1996; 39: 481–486. doi:10.1007/BF00400681
- [53] Cremonini F, Camilleri M, Clark MM, et al. Associations among binge eating behavior patterns and gastrointestinal symptoms: A population-based study. *Int J Obes* 2009; 33: 342–353. doi:10.1038/ijo.2008.272

- [54] Findlay RJ, Macrae EHR, Whyte IY, et al. How the menstrual cycle and menstruation affect sporting performance: Experiences and perceptions of elite female rugby players. *Br J Sports Med* 2020; 1–7. doi:10.1136/bjsports-2019-101486
- [55] Snipe RMJ, Costa RJS. Does biological sex impact intestinal epithelial injury, small intestine permeability, gastrointestinal symptoms and systemic cytokine profile in response to exertional-heat stress? *J Sports Sci* 2018; 36: 2827–2835. doi:10.1080/02640414.2018.1478612
- [56] van Wijck K, Lenaerts K, van Loon LJC, et al. Exercise-Induced splanchnic hypoperfusion results in gut dysfunction in healthy men. *PLoS One* 2011; 6: e22366. doi:10.1371/journal.pone.0022366
- [57] Schellekens DHSM, Hundscheid IHR, Leenarts CAJ, et al. Human small intestine is capable of restoring barrier function after short ischemic periods. *World J Gastroenterol* 2017; 23: 8452–8464. doi:10.3748/wjg.v23.i48.8452
- [58] Coutts A, Reaburn P, Abt G. Heart rate, blood lactate concentration and estimated energy expenditure in a semi-professional rugby league team during a match: a case study. *J Sports Sci* 2003; 21: 97–103. doi:10.1080/0264041031000070831
- [59] Islam H, Townsend LK, McKie GL, et al. Potential involvement of lactate and interleukin-6 in the appetite-regulatory hormonal response to an acute exercise bout. *J Appl Physiol* 2017; 123: 614–623. doi:10.1152/jappphysiol.00218.2017
- [60] Broom DR, Stensel DJ, Bishop NC, et al. Exercise-induced suppression of acylated ghrelin in humans. *J Appl Physiol* 2007; 102: 2165–2171. doi:10.1152/jappphysiol.00759.2006
- [61] King NA, Burley VJ, Blundell JE. Exercise-induced suppression of appetite: effects on food intake and implications for energy balance. *Eur J Clin Nutr* 1994; 48: 715–724
- [62] Deighton K, Barry R, Connon CE, et al. Appetite, gut hormone and energy intake responses to low volume sprint interval and traditional endurance exercise. *Eur J Appl Physiol* 2013; 113: 1147–1156. doi:10.1007/s00421-012-2535-1
- [63] Holliday A, Blannin A. Appetite, food intake and gut hormone responses to intense aerobic exercise of different duration. *Journal of Endocrinology* 2017; 235: 193–205. doi:10.1530/JOE-16-0570
- [64] Alajmi N, Deighton K, King JA, et al. Appetite and Energy Intake Responses to Acute Energy Deficits in Females versus Males. *Med Sci Sports Exerc* 2016; 48: 412–420. doi:10.1249/MSS.0000000000000793
- [65] Thackray AE, Deighton K, King JA, et al. Exercise, appetite and weight control: Are there differences between men and women? *Nutrients* 2016; 8: 1–18. doi:10.3390/nu8090583
- [66] Clarkson PM. Exertional rhabdomyolysis and acute renal failure in marathon runners. *Sports Medicine* 2007; 37: 361–363. doi:10.2165/00007256-200737040-00022
- [67] Lambert GP, Boylan M, Laventure JP, et al. Effect of aspirin and ibuprofen on GI permeability during exercise. *Int J Sports Med* 2007; 28: 722–726. doi:10.1055/s-2007-964891

- [68] Van Wijck K, Lenaerts K, Van Bijnen AA, et al. Aggravation of exercise-induced intestinal injury by ibuprofen in athletes. *Med Sci Sports Exerc* 2012; 44: 2257–2262. doi:10.1249/MSS.0b013e318265dd3d
- [69] Sostres C, Gargallo CJ, Lanas A. Nonsteroidal anti-inflammatory drugs and upper and lower gastrointestinal mucosal damage. *Arthritis Res Ther* 2013; 15: 1–8
- [70] Cornu C, Grange C, Regalin A, et al. Effect of Non-Steroidal Anti-Inflammatory Drugs on Sport Performance Indices in Healthy People: a Meta-Analysis of Randomized Controlled Trials. *Sports Med Open* 2020; 6. doi:10.1186/s40798-020-00247-w
- [71] Whatmough S, Mears S, Kipps C. The use of non-steroidal anti-inflammatories (NSAIDs) at the 2016 London marathon. *Br J Sports Med* 2017; 51: 409 LP – 409. doi:10.1136/bjsports-2016-097372.317
- [72] Holmes N, Cronholm PF, Duffy AJIII, et al. Nonsteroidal Anti-Inflammatory Drug Use in Collegiate Football Players. *Clinical Journal of Sport Medicine* 2013; 23
- [73] Drew MK, Vlahovich N, Hughes D, et al. A multifactorial evaluation of illness risk factors in athletes preparing for the Summer Olympic Games. *J Sci Med Sport* 2017; 20: 745–750. doi:10.1016/j.jsams.2017.02.010
- [74] Francino MP. Antibiotics and the human gut microbiome: Dysbioses and accumulation of resistances. *Front Microbiol* 2016; 6: 1–11. doi:10.3389/fmicb.2015.01543
- [75] Hughes RL. A Review of the Role of the Gut Microbiome in Personalized Sports Nutrition. *Front Nutr* 2020; 6. doi:10.3389/fnut.2019.00191
- [76] McFarland L V. Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of *Clostridium difficile* disease. *Am J Gastroenterol* 2006; 101: 812–822. doi:10.1111/j.1572-0241.2006.00465.x
- [77] Conlon MA, Bird AR. The impact of diet and lifestyle on gut microbiota and human health. *Nutrients* 2015; 7: 17–44. doi:10.3390/nu7010017
- [78] Bishehsari F, Ph D, Desai V, et al. Alcohol and Gut-Derived Inflammation. *Alcohol Res* 2013; 38: 163–171
- [79] West NP, Colbey C, Vider J, et al. Nutrition Strategies for Gut Health and Immune Function — What Do We Know and What Are the Gaps? *Sports Science Exchange* 2017; 28: 1–5
- [80] Scrivin R, Costa RJS, Pelly F, et al. An exploratory study of the management strategies reported by endurance athletes with exercise-associated gastrointestinal symptoms. *Front Nutr* 2022; 9. doi:10.3389/fnut.2022.1003445
- [81] Scrivin R, Costa RJ, Pelly F, et al. Development and validation of a questionnaire investigating endurance athletes practices to manage gastrointestinal symptoms around exercise. *Nutrition & Dietetics* 2021; 1–10. doi:10.1111/1747-0080.12674
- [82] Gaskell SK, Snipe RMJ, Costa RJS. Test–retest reliability of a modified visual analog scale assessment tool for determining incidence and severity of gastrointestinal symptoms in response to exercise stress. *Int J Sport Nutr Exerc Metab* 2019; 29: 411–419. doi:10.1123/ijsnem.2018-0215

Figure legends:

Figure 1: Median (IQR) gastrointestinal symptom severity scores by symptom between female (grey) and male (dark grey) players; at rest (A) and around rugby (B); Hburn, Heartburn; Burp, Burping; U.abp, Upper abdominal pain; Naus, Nausea; Vomit, Vomiting; Cramp, Cramping; Bloat, Bloating; L. abp, Lower abdominal pain; Flat, Flatulence; Const, Constipation; Defec, , Urgent need to defecate; Stool, change in stool consistency; * denotes a significant difference between male and female players ($p<0.05$) Gastrointestinal symptom severity; 1. No discomfort 2. Minor discomfort 3. Mild discomfort 4. Moderate discomfort 5. Moderately severe discomfort 6. Severe discomfort 7. Very severe discomfort

Figure 2: Median (IQR) and range of ratings of perceived feelings of hunger and desire to eat after training and matches using a Visual analogue scale (Hunger training; feeling of hunger post training Hunger match, feelings of hunger post match, Desire training, desire to eat post training, Desire match, desire to eat post match) in male (A) and female (B) and players; * denotes a significant difference between pre and post training or pre and post match, $p<0.01$

Table titles and legends

Table 1. Characteristics of participants

Data are presented as percentage (number) of players; kg, kilograms; cm, centimetres; #denotes a significant difference between males and females; * denotes a significant difference between forwards and backs of the same sex, $p<0.05$

Table 2. Self-reported frequency and severity of gastrointestinal symptoms reported at rest and around rugby

Data are presented as the percentage (number) of players who reported any gastrointestinal symptom experienced weekly or more often (weekly, more than once a week or daily); GIS, gastrointestinal symptom, ^a denotes a significant difference the frequencies reported between rest and rugby, ($p<0.05$), ^b denotes a significant difference between male and female players for the reported frequency of an individual GIS ($p<0.05$), *denotes a significance between rest and rugby for any GIS reported as a severity of ≥ 4 , ($p<0.05$)

Table 3 Severity scores of self-reported gastrointestinal symptoms reported by rugby players at rest and around rugby

Data is reported as median [interquartile range] of the GIS severity scores, * denotes a significant difference between female and male players for the comparative score, ($p<0.05$), ^a denotes a significant difference between rest and rugby environments, ($p<0.05$)

Table 4. Frequency of participants reporting different non-exercise associated risk factors for gastrointestinal symptoms (all) and in relation to reported GIS at rest or around rugby (n=329)

NSAIDs, Non-steroidal anti-inflammation medication, GIS represents players who reports one of more gastrointestinal symptom with a frequency of weekly or more often at rest or around rugby

Table 1. Characteristics of participants

Characteristics	Male (n=271)	Female (n=54)
-----------------	--------------	---------------

Country of origin	<i>Ireland</i>	42% (115)	63% (34)
	<i>England</i>	32% (86)	2% (1)
	<i>South Africa</i>	24% (64)	17% (9)
	<i>Scotland</i>	1% (3)	18% (10)
	<i>Australia</i>	1% (2)	
Rugby union format	<i>Fifteens</i>	84% (226)	52% (28)
	<i>Sevens</i>	4% (11)	22% (12)
	<i>Both</i>	12% (32)	14% (14)
Level of play	<i>Elite/International</i>	21% (58)	
	<i>Highly trained/ National level</i>	59% (158)	100% (54)
	<i>Trained / Amateur divisions</i>	20% (54)	
Age (years)	<i>18-24</i>	63% (171)	48% (26)
	<i>25-35</i>	35% (96)	50% (27)
	<i>>35</i>	1% (3)	2% (1)
Primary playing position	<i>Forwards</i>	53% (143)	40% (22)
	<i>Backs</i>	43% (118)	52% (28)
	<i>n/a</i>	4% (9)	8% (4)
Hours of training/week	<i>3-<5</i>	3% (8)	
	<i>5-<7</i>	15% (40)	2% (1)
	<i>7-10</i>	27% (74)	30% (16)
	<i>>10</i>	49% (132)	63% (34)
Self-reported body mass (kg)	<i>All</i>	100.0 ± 12.9 [#]	72.2 ± 9.9
	<i>Forwards</i>	108.1 ± 8.8 [*]	78.7 ± 9.8 [*]
	<i>Backs</i>	89.1 ± 8.4	66.3 ± 5.9
Self-reported stature (cm)	<i>All</i>	185.2 ± 7.5 [#]	167.7 ± 6.5
	<i>Forwards</i>	188.2 ± 7.0 [*]	170.4 ± 5.6
	<i>Backs</i>	181.6 ± 6.5	165.9 ± 6.9

Data are presented as percentage (number) of players; kg, kilograms; cm, centimetres; [#]denotes a significant difference between males and females; ^{*} denotes a significant difference between forwards and backs of the same sex, $p < 0.05$

Table 2. Self-reported frequency and severity of gastrointestinal symptoms reported at rest and around rugby

	<i>All (n=325)</i>	<i>Male (n=271)</i>	<i>Female (n=54)</i>
--	--------------------	---------------------	----------------------

	At rest	Around rugby	At rest	Around rugby	At rest	Around rugby
Upper GIS % (n)						
Heartburn	6% (22) ^a	4% (14)	7% (21) ^a	4% (13)	1% (1)	1% (1)
Burping	27% (88) ^a	11% (35)	27% (74) ^a	9% (24)	25% (14) ^a	16% (9)
Upper abdominal pain	4% (12)	4% (13)	2% (5)	2% (6)	13% (7)	13% (7)
Nausea	7% (23)	7% (24)	6% (18)	7% (19)	7% (4)	9% (5)
Vomiting	1% (4)	1% (4)	1% (2) ^a	1% (4)	3% (2)	0% (0)
Stomach cramps/ gurgling	14% (45)	15% (48)	11% (29)	13% (34)	30% (16)	26% (14)
Lower GIS % (n)						
Bloating	20% (65)	15% (48)	15% (41) ^a	11% (30)	44% (24) ^a	33% (18)
Lower abdominal pain	6% (20)	6% (20)	4% (10)	4% (10)	19% (10)	19% (10)
Flatulence	48% (156)	28% (92)	48% (129) ^a	27% (73)	50% (27) ^a	37% (20)
Constipation	5% (15)	5% (17)	4% (10)	4% (10)	9% (5)	13% (7)
Diarrhoea	8% (27)	19% (61)	7% (20) ^a	15% (42)	13% (7) ^a	35% (19)
Urgent need to defecate	11% (35)	13% (43)	11% (29)	13% (35)	11% (6) ^a	15% (8)
Change in stool consistency	25% (80)	22% (71)	25% (68) ^a	18% (46)	22% (12)	28% (15)
Any upper GIS ≥ once a week	37% (120)	28% (93)	34% (92) ^a	26% (71)	52% (28)	41% (22)
Any lower GIS ≥ once a week	59% (189)	42% (137)	57% (153)	39% (104)	67% (36)	61% (33)
Any GIS ≥ once a week	62% (203)	47% (164) ^a	60% (164)	43% (118) ^a	72% (39) ^b	67% (36)
Players reporting any GIS severity ≥4	42% (138)	33% (106)*	40% (108)	29% (75)*	64% (35)	57% (31)

Data are presented as the percentage (number) of players who reported any gastrointestinal symptom experienced weekly or more often (weekly, more than once a week or daily); GIS, gastrointestinal symptom, ^a denotes a significant difference the frequencies reported between rest and rugby, ($p < 0.05$), ^b denotes a significant difference between male and female players for the reported frequency of an individual GIS ($p < 0.05$), *denotes a significance between rest and rugby for GIS severity, reported as a severity of ≥ 4 , ($p < 0.05$)

Table 3: Severity scores of self-reported gastrointestinal symptoms reported by rugby players at rest and around rugby

	All (n=325)	Male (n=271)	Female (n=54)
--	--------------------	---------------------	----------------------

	At rest	Around rugby	At rest	Around rugby	At rest	Around rugby
GIS severity score: upper	9 [7-13]	8 [6-11] ^a	9 [7-12]	7 [6-11] ^a	12 [8-15]	9 [7-13]*
GIS severity score: lower	11 [8-16]	11 [7-16] ^a	11 [8-15]	10 [7-14] ^a	15 [11 – 21]	17 [11-22]*
Total GIS severity score	21 [16-29]	19 [14-27] ^a	20 [15-27]	18 [13-26] ^a	26 [20 – 38]	26 [19-36]*

Data is reported as median [interquartile range] of the GIS severity scores, *denotes a significant difference between female and male players for the comparative score, ($p < 0.05$), ^a denotes a significant difference between rest and rugby environments, ($p < 0.05$)

Table 4. Frequency of participants reporting different non-exercise associated risk factors for gastrointestinal symptoms (all) and in relation to reported GIS at rest or around rugby (n=329)

Non-exercise risk factors	Overall frequency	GIS at rest	GIS around rugby
Take oral probiotics regularly (>70% of the time)	23%	14% (47)	14% (44)
Include prebiotic or fermented food at least once a week	22%	12% (39)	10% (31)
Eat 5 or more servings of vegetables per day	32%	21% (69)	17% (53)
Include an alcoholic drink once a week or more after rugby	34%	24% (78)	16% (51)
Take NSAIDs 2-3 times a month or more often	22%	14% (44)	12% (39)
Take antibiotics once or more over the last 12 months	33%	7% (24)	20% (64)

NSAIDs, Non-steroidal anti-inflammation medication, GIS represents players who reports one of more gastrointestinal symptom with a frequency of weekly or more often at rest or around rugby

Figure 1




Figure 2

