



LEEDS
BECKETT
UNIVERSITY

Citation:

Manono, G and Thiong'o, S and Wishitemi, BE (2022) "Perceived effects of climate change on plants and its potential impact on tourism - a perspective of the local residents of Maasai Mara National Reserve, Kenya." In: Okech, R and Kieti, D and van der Duim, V.R., (eds.) Tourism, Climate Change and Biodiversity in Sub-Saharan Africa. African Studies Centre Leiden (ASCL) April 2023, The Netherlands, pp. 27-47. ISBN 9789054481966

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/10969/>

Document Version:

Book Section (Published Version)

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

© 2022 R. Okech, D. Kieti, V.R. van der Duim

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

2

Perceived effects of climate change on plants and its potential impact on tourism – a perspective of the local residents of Maasai Mara National Reserve, Kenya

G. Manono, S.M. Thiong’o and B.E. Wishitemi

2.1 Introduction

The magnitude of the climate changes in the 21st century is comparable to the level of warming during the last deglaciation (Buizert et al., 2014; Corrick et al., 2020). For instance, according to the Intergovernmental Panel on Climate Change (IPCC, 2014), the global climate has warmed up by approximately 1.18°C over the period from 1880 to 2020. This is also supported by other agencies such as the Goddard Institute of Space Sciences and the American Meteorological Society that postulate that global temperatures will continue to rise (Lindsey & Dahlman, 2021; Hawkins et al., 2017). Furthermore, extreme precipitation events will become more intense and frequent, while oceans will continue to warm, acidify and rise. A study by Thomas et al. (2004), which modelled the expected impact of gradual climate change on 1,103 species, predicted that 15-37% of these species would be committed to extinction by 2050. The Kenyan Government (GoK) contends that both minimum (nighttime) and maximum (daytime) temperatures have increased by 0.7°C-2.0°C and 0.2°C-1.3°C, respectively, since the early 1960s (GoK, 2010). Rainfall has become more irregular, unpredictable, and intense (Mwenda et al., 2020; Omony, Wakhungu & Oteng’i, 2015). Whereas precipitation has generally decreased in the main rainfall season between March and May, it has increased in the previously much shorter rain season between October and December. Moreover, severe floods are occurring (or are predicted to occur) along the coastal strip and the northern parts of the country in seasons which are normally dry such as that between September and February (Parry et al., 2012; Marigi, 2017; UNEP, 2021; GoK, 2010).

Human activities have been found to be by far the major cause of climate change, through their continuous release of greenhouse gases and aerosols into the atmosphere, through changing land surfaces, and through depleting the stratospheric Ozone Layer (Abdollahbeigi, 2020; IPCC, 2014; Foukal et al., 2006; Krishnan et al., 2020; Schmale, Zieger & Ekman, 2021). No wonder an increasing band of researchers are suggesting that the current age be named the 'Anthropocene', a new division of geological time, in recognition of the tremendous impact that *Homo sapiens* has on the planet (Zalasiewicz et al., 2015). Like other developing economies in Africa and further afield that rely on tourism as a generator of foreign exchange, Kenya depends heavily on the outdoor recreational opportunities presented by its natural environment, which include pristine nature, spectacular landscapes and seascapes, rare species, and wildlife in their natural habitat (Akama, 1999; GoK, 2010; Maingi, 2020; Nampushi & Nankaya, 2020; World Bank Group, 2010). Apart from these, idyllic climatic conditions are an enabler of yet another major tourism pull factor – beach tourism (Smith, 1993; Hein et al., 2009; Papageorgiou, 2016).

Adverse impacts of climate change, however, threaten to scupper this source of livelihood, through their assault on biodiversity. Vegetation forms a substantial segment of the earth's biomass (Fleming, 2015; Thompson, 2018), and hence, the greatest impacts of climate change in the country result from its influence on plants. Moreover, plants are the autotrophs in an ecosystem, and hence, their depletion will affect other organisms, for instance, animals. Paleo-ecological evidence suggests that organisms respond to climate change usually by migrating, whereas evolutionary adaptation plays only a minor role (Huntley, 1991). Given the limited dispersal and/or migratory capacity of most plants, they are likely to be prone to climate change effects. Drought may cause some tree species to disappear and, as a consequence, affect both vegetation structure and species composition (February et al., 2007). Models of future biome distributions in tropical South America predict the substitution of Amazonian forest cover by savannah-like vegetation (Salazar et al., 2007; Lapola et al., 2009). Barlow and Peres (2008) expect forest dieback in West and Southern Africa as a result of climate change. Parmesan and Yohe (2003) reported that alpine herbs are shifting poleward, 6.1 km per decade on average. Further, studies of plant phenology have attributed longer growing seasons, earlier onset of flowering, and earlier harvesting times to climate warming (Parmesan, 2006).

Evidence has suggested that climate change is caused by both natural and man-made factors over a period of time (Earth Science Communications, 2021; Oreskes, 2004; USEPA, 2020; USGS, 2021). The natural processes im-

plicated in climate change include volcanic eruptions, variations in the sun's intensity, as well as very slow changes in ocean circulation or land surfaces which occur on time scales of decades, centuries or longer. The influence of diverse external factors on the climate can be broadly estimated using the concept of radiative forcing – a measure of the influence a factor has in altering the balance of incoming and outgoing energy in the Earth. A positive radiative forcing warms the surface (for instance, greenhouse gases), while a negative radiative forcing cools the surface (for example, some types of aerosols) (IPCC, 2001).

The most important greenhouse gases that have been found to cause positive radiative forcing include carbon (IV) oxide (CO_2), methane (CH_4), and nitrous oxide/nitrogen (I) oxide (N_2O) (Dupar, 2020; Shakoor et al., 2021). In addition, halocarbon gases that have been found to be both ozone-depleting, and greenhouse gases include trichlorofluoromethane (CFCl_3), dichlorodifluoromethane (CF_2Cl_2), chlorodifluoromethane (CHF_2Cl) and 1, 1, 1, 2-tetrafluoromethane ($\text{CF}_3\text{CH}_2\text{F}$) (IPCC, 2014; IPCC, 2001). The radiative forcing due to increases in the greenhouse gases from 1750 to 2000 is estimated to be 2.43 Wm^{-2} : 1.46 Wm^{-2} from CO_2 (60%); 0.48 Wm^{-2} from CH_4 ; 0.34 Wm^{-2} from the halocarbons; and 0.15 Wm^{-2} from N_2O (IPCC, 2001).

Amidst this inundation of information on climate change and how deleterious its impacts are or could be, it remains underexposed how people, especially those living in marginal areas and likely to be adversely affected by climate change, perceive climate change effects on plants. For instance, an assessment carried out by Mutimba et al. (2010) on climate change vulnerability and adaptation preparedness in Kenya, concluded that climate change awareness, especially in the countryside, was quite low. A Gallup poll carried out between 2007 and 2008 by Pelham (2009) reported that 56% of Kenyans had some knowledge of global warming, whereas 44% had no notion of climate change whatsoever. However, also those who had some knowledge of global warming, were often not well versed in various climate change issues, such as adaptation and mitigation arguments (Pelham, 2009). This is in agreement with related studies locally and further afield (see Sraku-Lartey et al., 2020; Nash et al., 2019; Takakura et al., 2021; Mutekwa, 2009; Van Aalst et al., 2008 and Scheffran et al., 2012). The studies concur that these locals, given their living in remote locations, strained economic capabilities and/or low literacy levels, profess their ignorance about climate change and its effects. According to a study by Froehlich and Al-Saidi (2018), a reason for this might be that some view climate change as a natural occurrence and its management as a government policy issue.

Therefore, this study seeks to establish the perceived causes and manifestations of climate change amongst the community¹ in, and adjacent to, the MMNR, and to determine the perceived effects of climate changes on plant communities in the region.




Figure 2.1
The Maasai Mara ecosystem (Source: Farr et al. 2019)

2.2 Materials and Methods

The study was carried out in the Maasai Mara National Reserve and its environs (the Mara ecosystem) (Figure 2.1). Located approximately 180 km west of Nairobi, the reserve adjoins the Serengeti National Park, forming an extensive wildlife dispersal area (Maasai Mara National Reserve, 2021; UNESCO, 2021). To the north, east and west of the reserve are large parcels of land demarcated as group ranches, owned and inhabited by the semi-nomadic pastoral Maasai people.

This study employed both a survey research design, which enabled it to obtain the requisite information from a large segment of the populace over a short period of time, and an exploratory research design, which allowed for

¹ The term local/community residents in the context of this study refers to people living in and adjacent to the MMNR. It includes both the staff of MMNR (managers), community leaders, and the local rural folk.

deeper probing of respondents' attitudes, preferences and opinions (Oso & Onen, 2008). These designs used questionnaires and face to face interviews respectively.

The target population consisted of 1,500 residents of MMNR, including opinion leaders of the local community, local community members, and managers of the MMNR. This study collected data from 400 respondents, constituting 27% of the target population. This was close enough to the caveat suggested by Mugenda & Mugenda (2003) and Montgomery (1977), who stated that 30% of the accessible population would suffice for a descriptive study if the population units were more than 30. To ensure a proportionate representation, the sample consisted of 200 community members and staff members of the MMNR. The field study was conducted between the months of May and June of 2013². A sampling frame of the respondents was obtained from either the reserve or the villages which served to select the participants for the study, using simple random sampling. The study employed two methods: questionnaires and interview schedules, to collect both qualitative and quantitative data. The data collection tools were administered by the researcher, two supervisors and six trained enumerators. Descriptive statistics, for instance, were used to describe, summarise, and organise the data. Chi-square (χ^2) cross-tabulations were used to test if there were any significant relationships between the study variables. Means from the study were compared using t-tests and Analysis of Variance (ANOVA). The t-tests were used to compare two independent groups. More than two means were compared using Analysis of Variance (ANOVA) and post-hoc analysis carried out by Tukey's Honestly Significant Difference (HSD) test.

Frequencies were used to analyze the perceived causes and manifestations of climate change amongst the Maasai community in and adjacent to the MMNR. Structural Equation Modelling-Path Analysis (SEMPATH), implemented using the Analysis of Moment Structures (AMOS), was used to examine the perceived effects of climate change on plant communities in MMNR. The model (Figure 2.2) hypothesised that climate change could influence changes in plant species distribution, species composition, changes in adaptation strategies, changes in species diversity, the emergence of alien plant species, changes in vegetation cover, and extinction of plant species. Climate change was specified as an exogenous measured variable, while the perceived effects were postulated as observed endogenous variables.

² The data, though collected a little earlier, has not been presented elsewhere and is thus scientifically valid.




Figure 2.2
Path diagram of perceived effects of climate change on plants; spp. = species

All the above statistical tests were analysed with the Statistical Package for Social Sciences (SPSS) version 18, and STATA version 12. All statistical tests were two-tailed. Significant levels were measured at a 95% confidence level, with significant differences recorded at $p < 0.05$.

2.3 Results

Response rate

From a target population of 1,500 local residents and a sample of 400 respondents, the study gathered 386 usable responses. This was a response rate of 96.5%, which was considered sufficient for analysis (Mugenda & Mugenda, 2003).

Sample characteristics

Gender distribution (Table 2.1) showed that the respondents were disproportionately male (staff: male=80%, female=20% and local community: male=94%, female=6%). The preponderance of male respondents in the study may be adduced to the traditional cultural roles among the Maasai people, the predominant community living around MMNR (Allegretti, 2018). Older men usually take the advisory role in their homestead, whereas women build houses and are responsible for childcare and all household chores, and young men are in charge

of security. On the other hand, young boys herd livestock (Bitange, 2005). Since women are tied down to tasks that occur mainly around the home, one is likely to interact with men rather than women when carrying out a study.

Most of the respondents were aged between 31 and 40 years (staff=70%, local community =84.5%), suggesting that the bulk of the respondents had lived long enough to be aware of changes in the climate. Fewer respondents belonged to the 21-30 years (staff=25.5%, local community=11.5%) and 41-50 years (staff=4.5%, local community=4%) age brackets. The results indicated that while the majority of staff had secondary education (n=110, 55%), the bulk of the local community possessed primary education (n=154, 77%). Interestingly, there were slightly more people with college or university education among the local community (15.5%) compared to the staff (12%), suggesting that pockets of the local community are reasonably well educated.

Table 2.1
Respondents' Characteristics

Biographic information	Respondent type	Categories	Per cent
Respondents' gender	Staff	Male	80.0
		Female	20.0
		Total	100.0
	Local community	Male	94.0
		Female	6.0
		Total	100.0
Respondent's age	Staff	21-30 years	25.5
		31-40 years	70.0
		41-50 years	4.5
		Total	100.0
	Local community	21-30 years	11.5
		31-40 years	84.5
		41-50 years	4.0
		Total	100.0
Highest education level	Staff	None	5.0
		Primary	28.0
		Secondary	55.0
		College	5.0
		University	7.0
	Local community	Total	100.0
		None	2.0
		Primary	77.0
		Secondary	5.5
		College	6.5
University	9.0		
Total	100.0		

Source: Survey data (2013)

Awareness of climate change in the community

Awareness of climate change among the respondents was estimated, as this knowledge could help shape the appropriate interventionist strategies. The study found that 96% (n=192) and 97% (n=194) of employees of the MMNR and the neighbouring Maasai community had heard of climate change, respectively. This suggested that the climate change message could be pervasive in the community. A χ^2 cross-tabulation to determine if awareness of climate change was influenced by membership of the local community or staff was found to be non-significant ($\chi^2=0.29$, $df=1$, $p=0.586$), indicating no significant differences between the staff and members of the local community with respect to awareness of climate change.

However, the majority of the respondents (n=308, 77%) considered themselves as 'somewhat knowledgeable' about climate change as compared to only 20% (n=78) and 2% (n=7), who felt that they were 'knowledgeable' or 'experts', respectively. The results suggested that although most respondents knew something about climate change, they might be ignorant of the technical details of the process.

Respondents' gender ($\chi^2 = 5.857$, $df=1$, $p=.016$); age ($\chi^2 = 76.037$, $df=2$, $p<.001$) and education ($\chi^2 = 63.09$, $df=4$, $p<.001$) were found to significantly influence their perception of their own knowledge about climate change. More females (Table 2.1.) perceived themselves to be knowledgeable about climate change (33.3%) compared to the males (18.3%), whereas more males were found to be somewhat knowledgeable (81.7%) compared to the females (66.7%). This could be related to the fact that the females in the study were older and better educated than the males. Furthermore, it could be deduced that the younger and less educated members of the local community were reluctant to participate in the study – an aspect that could be attributed to the cultural structure of the Maasai community, where the young are customarily supposed to serve their community through chores (Obeja, 2015; Kerubo, 2016). This conclusion was buttressed by the demographics that respondents aged between 41-50 years had the highest proportion of knowledgeable people with respect to climate change (76.9%) compared with those in the 31-40 years age bracket (11.2%). The results in Table 2.2 indicate that education was positively correlated with the level of knowledge about climate change, with respondents having college and university education rating themselves highest with regard to climate change knowledge (68.2% and 55.2%, respectively).

Table 2.2

The relationship between climate change knowledge and biographical variables

Respondents' category			Level of knowledge		Total	
			Somewhat knowledgeable	Knowledgeable		
Gender	Male	Frequency	276	62	338	
		Percentage	81.7	18.3	100.0	
	Female	Frequency	32	16	48	
		Percentage	66.7	33.3	100.0	
	Total	Frequency	308	78	386	
		Percentage	79.8	20.2	100.0	
Age	21-30 years	Frequency	36	34	70	
		Percentage	51.4	48.6	100.0	
	31-40 years	Frequency	269	34	303	
		Percentage	88.8	11.2	100.0	
	41-50 years	Frequency	3	10	13	
		Percentage	23.1	76.9	100.0	
	Total	Frequency	308	78	386	
		Percentage	79.8	20.2	100.0	
	Education	None	Frequency	6	2	8
			Percentage	75.0	25.0	100.0
Primary		Frequency	184	25	209	
		Percentage	88.0	12.0	100.0	
Secondary		Frequency	98	20	118	
		Percentage	83.1	16.9	100.0	
College		Frequency	7	15	22	
		Percentage	31.8	68.2	100.0	
University		Frequency	13	16	29	
		Percentage	44.8	55.2	100.0	
Total		Frequency	308	78	386	
		Percentage	79.8	20.2	100.0	

Perceived causes and manifestations of climate change

The respondents perceived that climate change could be caused by deforestation, human settlement, agriculture, overutilisation of natural resources, greenhouse emissions, infrastructure and overharvesting of indigenous trees (Table 2.3). Most respondents (81%) considered deforestation to be the most important cause of climate change, followed by overharvesting of indigenous trees (72%) and human settlement (71%). The findings are corroborated by other studies carried out by Bennet (2017), Schlamadinger et al. (2005), and

Malhi et al. (2008), among others, who documented the deleterious effects of the loss of forest cover. The least important cause, according to the respondents, were greenhouse emissions (11%), followed by the construction of infrastructure (20%).

Most respondents (n=311, 78%) reported that erratic rainfall patterns were the most important manifestation of climate change, followed by droughts (n=286, 72%), floods (n=213, 53%), and lastly, increased temperatures (n=184, 46%).

Table 2.3
Frequencies of response on causes of climate change

Name of variable	Causes of climate change									
	Least important		Less important		Important		More important		Most important	
	Fq	%	Fq	%	Fq	%	Fq	%	Fq	%
Deforestation	21	5.3	22	5.5	12	3.0	21	5.3	324	81.0
Human settlement	11	2.8	51	12.8	30	7.5	22	5.5	286	71.5
Agriculture	11	2.8	50	12.5	32	8.0	174	43.5	133	33.3
Overutilisation of natural resources	13	3.3	20	5.0	54	13.5	107	26.8	206	51.5
Greenhouse emissions	258	64.5	18	4.5	47	11.8	33	8.3	44	11.0
Construction of infrastructure	20	5.0	249	62.3	29	7.3	23	5.8	79	19.8
Overharvesting of indigenous trees	22	5.5	11	2.8	38	9.5	41	10.3	288	72.0

Key: Fq = Frequency

Perceived effects of climate change on plants

An overwhelming proportion (99%, n=198) of local community members felt that climate change affected plants, compared to 81% (n=162) of the staff of MMNR who held a similar opinion. Most respondents (Table 2.4) felt that climate change influenced changes in plant species composition, species distribution patterns, changes in plants' adaptation strategies, changes in plant species diversity, the emergence of alien plant species, and changes in vegetation cover (each accounting for 16% of the 2529 responses). However, very few respondents thought that climate change was causing the extinction of plant species (6% of the 2529 responses).

For this question, the number of responses (2529) was more than the number of respondents in the study (400), showing that most respondents felt that climate change had more than one influence on plants (the question was a multiple response type question). Additionally, the total percentage of cases was 632.3, indicating that, on average, each respondent felt that climate change had six (632.3/100) types of influence on plants.

Table 2.4
Perceived impacts of climate change on plants

Plant impacts	Responses		Per cent of cases
	N	Per cent	
Changes in plant species diversity	394	15.6	98.5
Changes in plant species composition	396	15.7	99.0
Changes in plant species distribution patterns	397	15.7	99.3
Emergence of alien plant species	393	15.5	98.3
Extinction of plant species	162	6.4	40.5
Changes in plants' adaptation strategies	396	15.7	99.0
Changes in vegetation cover	391	15.5	97.8
Total	2529	100.0	632.3

Face-to-face interviews with older members of the community supported the above findings. One elder asserted:

After flooding, new or alien plants grow on bare ground, hence colonising the whole area, thus affecting the original vegetation. This brings about changes in plant species and the distribution patterns of plants. The floods and drought have brought changes in this area, hence the vegetation cover change.

Another stated:

The change in temperature, soils, and rainfall patterns has brought significant changes in the following: changes in plant species, changes in plant distribution patterns and introduction of alien species of plants. Some plants have been forced to adapt to the current state of the climate. You get some plant species growing where you least expect them.

To determine whether the perceived effects of climate change influenced plants in the MMNR, a SEMPATH analysis was conducted. The resultant path diagram is presented in Figure 2.3.




Figure 2.3
SEMPATH output of the perceived effects of climate change on plants

All the regression coefficients for the model were significantly different from zero beyond the 0.01 level. Except for the perceived effect on the extinction of plant species (coefficient = -0.37), all the path coefficients for the remaining six hypothesised relationships were positive. This indicated that when climate change increases, it causes changes in plant species diversity, plants' adaptation strategies, and species distribution. It also causes changes in the plants' vegetation cover, causes the emergence of alien species, and brings about changes in species composition. However, climate change was perceived to reduce the extinction of plant species. This implied that the respondents viewed climate change as having minimal effects on this variable.

2.4 Discussion

This study found a high awareness of climate change in the community (about 97%), which was unlike studies by Mutimba et al. (2010) and Pelham (2009), who reported that only 56% of Kenyans had some knowledge of global warming, whereas 44% knew nothing. The findings could suggest

that climate change mitigation initiatives, public sensitisation, and awareness programmes such as the National Climate Change Response Strategy 2010 as well as other reforestation and carbon offset projects undertaken by the government, NGOs, and the media might have been effective. Other current projects include the Kenya Adaptation Action Plan (2015-2030), and the National Climate Change Action Plan (2018-2022). Although most respondents were found to know something about climate change, they were likely to be ignorant of the technical details of the process, as supported by Pelham (2009) who found out that many Kenyans who reported knowledge of climate change were not altogether well versed in various climate change issues such as adaptation and mitigation processes.

The perception of respondents about their own knowledge of climate change was found to be influenced by their gender, age and education level, with older age, better education and being female being associated with greater knowledge. Many studies report that in public, men demonstrate greater scientific knowledge and scientific literacy than do women (Miller, 2007; Hayes, 2001), largely due to differences in the way men and women experience science and mathematics education. In addition, many studies demonstrate that women are less confident in displaying their scientific knowledge and abilities than do men (for instance, McCright, 2010; Jacobs & Simpkins, 2006). From the findings, the percentage of female respondents was small but those who participated in the survey were more knowledgeable than their male counterparts. However, this is contrary to the documented expectations and might be related to the fact that the females in the study were older and better educated than the males. In this vein, older people have often been found to be more knowledgeable about climate change than their younger counterparts. For example, Ogunleye and Yekinni (2012) found a positive correlation between the age of crop farmers in Ilorin East, Nigeria and their knowledge of climate change.

Deforestation was perceived as the strongest cause of climate change, followed by overharvesting of indigenous trees, agriculture, and human settlement, while the least important causes were considered to be greenhouse emissions and the construction of infrastructure. Deforestation or logging has been a major concern amongst Kenyan conservationists. According to Cochrane and Laurence (2002), logging does not only lead to loss of habitat for animals in the forests but also to changes in the microclimatic environment, the erosion of soil and modification of fire regimes. The impact depends on the type of logging: commercial mechanised logging with heavy equip-

ment, or local exploitations of timber through, for example, pit-swaying and firewood collection.

Forests play a huge role in the carbon cycle on our planet by absorbing carbon (IV) oxide and releasing oxygen during the day. When forests are cut down, not only does their carbon absorption cease, but the carbon stored in the trees is released into the atmosphere as CO₂ if the wood is burned, or even if it is left to rot after the deforestation process (Karl & Trenberth, 2003). Hence, deforestation contributes to climate change by increasing the level of carbon (IV) oxide, the most dominant human-influenced greenhouse gas. It is estimated that more than 1.5 billion tons of carbon dioxide are released into the atmosphere due to deforestation, mainly the cutting and burning of forests, every year. In fact, whereas cars and trucks have been found to account for about 14 per cent of global carbon emissions, 15 per cent is usually added to deforestation (Stott et al., 2000). Overharvesting of indigenous trees, agriculture, and human settlement contribute to climate change for similar reasons as does deforestation, because all these activities involve the cutting down of some trees. In addition, agricultural activities produce gases such as methane (CH₄), nitrogen (I) oxide (N₂O), nitrogen (II) oxide (NO) and ammonia (NH₃), which are all implicated for their radiative or chemical effects in the atmosphere (IPCC, 2014; Li, 2000; Bollman & Conrad, 1998).

This study found that the climate change impacts on plants that were perceived as important were changes in plant species, plants' adaptation strategies, changes in the distribution of plants, changes in vegetation cover, the emergence of alien species and changes in plant composition, whereas the least important impact was considered to be the extinction of plant species. These findings are in line with research by Mackey and Currie (2001), which indicated that temperature and water availability account for more than 75% of the variability in plant species richness over broad spatial scales. Changes in climate will also alter interactions between species, including patterns of competition, symbiosis, mutualism, predation, and dominance. Climate change, in the form of altering rainfall patterns (for instance, increased droughts) and temperature, could explain the observable changes in plant species, their distribution and vegetation cover in this study.

Changes in plant distribution and an extended range of pests and pathogens caused by climate change can allow for an invasion by alien species (McCarty, 2001). For instance, *Prosopis juliflora* ('mathenge') has become dominant in important ecosystems of Baringo, Tana River, Garissa and other semi-arid areas of the country. In addition, an excessive growth of some tree species

has been observed, including that of *Acacia reficiens* (acacia) after the 1997 El-Nino in North-Eastern Province (NEP), suppressing the growth of various species that make up grasslands for wildlife and livestock (Mutimba et al., 2010). Increases in temperature could lead to a shift of vegetation to higher elevations, which are cooler, while some species could become extinct. Indeed, across the country, some tree species, including *Melia volkensii*, *Terminalia spinosa*, *Delonix elata*, and *Hyphenea corriaceae* in North Eastern Province, as well as *Psychotria* species in the Taita Hills, Coast Province, have either gone extinct, or else their numbers have dramatically reduced. In addition, the projected rise in temperatures and long periods of drought could lead to more frequent and more intense fires, with estimates showing that Kenya has lost more than 5,700 ha of forests per year to forest fires over the past 20 years (Mutimba et al., 2010).

The vulnerability of natural resources such as plants to climate change could have major implications for Kenya's tourism industry, because long-term environmental shifts could alter the destination attractiveness for vacationers (Papageorgiou, 2016; Martín, 2005; Forster et al., 2012). Unless carefully monitored, the impacts could lead to the downgrade and ultimately the collapse of the climate-sensitive coastal, marine and wildlife industry in Kenya.

2.5 Conclusion and recommendations

This study has shown that practically every respondent drawn from the employees of MMNR and the local community has heard of climate change, demonstrating that awareness of climate change in the locale can be very pervasive. No significant differences were found between the MMNR staff and members of the local community with respect to their awareness of climate change, suggesting the ubiquitousness of the climate change message among the two groups. However, the results suggested that although most respondents knew something about climate change, they could still be quite ignorant of the technical details of the phenomenon (Mutimba et al., 2010; Nanyingi et al., 2012). The perception of respondents about their knowledge of climate change was also found to be influenced by gender, age and education level. Deforestation was perceived as the strongest cause of climate change, followed by overharvesting of indigenous trees, agriculture and human settlement, greenhouse emissions and construction of infrastructure. Erratic rainfall patterns were found to be the most important manifestation of climate change, followed by droughts, floods, and increased temperatures. Most respondents felt that climate change influenced changes in plant species com-

position, species distribution patterns, changes in plants' adaptation strategies, changes in plant species diversity, the emergence of alien plant species, and changes in vegetation cover.

The study recommends that the government and other environmental agencies should disseminate appropriate knowledge about climate change to the public and take immediate steps to discourage habitat destruction (for instance, deforestation, infrastructural and superstructural development in and adjacent to protected areas, overharvesting of trees, large-scale agriculture, indiscriminate harvesting of trees, and human settlement in forests) as this contributes to climate change. Most importantly, climate change's pernicious impacts on nature need to be controlled locally, regionally and internationally. For instance, marine and terrestrial spatial and seasonal quantifications of climate resources through Tourism Climatic Indices could inform sustainability planning for outdoor tourism in Kenya.

References

- Abdollahbeigi, M. (2020). Non-climatic factors causing climate change. *Journal of Chemical Reviews*, 2(4): 292-308.
- Akama, J.S. (1997). Tourism development in Kenya: Problems and policy alternatives. *Progress in Tourism and Hospitality Research*, 3(2): 95-105.
- Allegratti, A. (2018). Respatialising culture, recasting gender in peri-urban sub-Saharan Africa: Maasai ethnicity and the 'cash economy' at the rural-urban interface, Tanzania. *Journal of Rural Studies*, 60: 122-9.
- Barlow, J. & Peres, C. (2008). Fire-mediated dieback and compositional cascade in an Amazonian forest. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363: 1787-94.
- Bennett, L. (2017). *Deforestation and Climate Change*. Washington: Climate Institute.
- Buizert, C., Gkinis, V., Severinghaus, J.P., He, F., Lecavalier, B.S., Kindler, P., & Brook, E.J. (2014). Greenland temperature response to climate forcing during the last deglaciation. *Science*, 345(6201): 1177-80.
- Bitange, N. (2005). Maasai entrepreneurship and change. *Journal of Small Business and Entrepreneurship*, 18(2): 66-75.
- Corrick, E.C., Drysdale, R.N., Hellstrom, J.C., Capron, E., Rasmussen, S.O., Zhang, X., & Wolff, E. (2020). Synchronous timing of abrupt climate changes during the last glacial period. *Science*, 369(6506): 963-9.
- Dupar, M. (2020). *The IPCC's Special Report on Climate Change and Land: What's in it for Africa?* Retrieved from <https://www.africaportal.org/publications/ipccs-special-report-climate-change-and-land-whats-it-africa/>

- Earth Science Communications (2021). *The Causes of Climate Change*. Retrieved from <https://climate.nasa.gov/causes/>
- Farr, M.T., Green, D.S., Holekamp, K.E., Roloff, G.J., & Zipkin, E.F. (2019). Multispecies hierarchical modeling reveals variable responses of African carnivores to management alternatives. *Ecological Applications*, 29(2): e01845.
- February, E., West, A., & Newton, R. (2007). The relationship between rainfall, water source and growth for an endangered tree. *Austral Ecology*, 32: 397-402. Retrieved from <http://www.iiste.org/Journals/index.php>
- Fleming N. (2015). *Which organism has had the biggest impact on the planet?* Retrieved from <http://www.bbc.com/earth/story/20150211-whats-the-most-dominant-life-form>
- Food and Agriculture Organisation (FAO) (2011). *Wildlife in a changing climate*. Retrieved from <http://www.fao.org/forestry/30085-0822ac7d034de02f6e2d8245b97bde5db.pdf>
- Forster, J., Schuhmann, P.W., Lake, I.R., Watkinson, A.R., & Gill, J.A. (2012). The influence of hurricane risk on tourist destination choice in the Caribbean. *Climatic Change*, 114(3): 745-68.
- Foukal, P., Frohlich, C., Spruit, H., & Wigley, T. (2006). Variations in solar luminosity and their effect on the earth's climate. September 14, 2006. *Nature*, 443: 161-6.
- Froehlich, P., & Al-Saidi, M. (2018). Local community perception of climate change adaptation in Egypt. *IOP Conference Series: Earth and Environmental Science*, 191(1): 012003).
- GoK (2010). *National climate change response strategy*. Retrieved from <http://www.environment.go.ke/wp>
- Hawkins, E., Ortega, P., Suckling, E., Schurer, A., Hegerl, G., Jones, P., & Van Oldenborgh, G.J. (2017). Estimating changes in global temperature since the preindustrial period. *Bulletin of the American Meteorological Society*, 98(9): 1841-56.
- Hayes, B.C. (2001). Gender, scientific knowledge, and attitudes toward the environment. *Political Research Quarterly*, 54: 657-71.
- Hein, L., Metzger, M.J., & Moreno, A. (2009). Potential impacts of climate change on tourism; a case study for Spain. *Current Opinion in Environmental Sustainability*, 1(2): 170-8.
- Huntley, B. (1991). How plants respond to climate change: Migration rates, individualism and the consequences for plant communities. *Annals of Botany*, 67(1): 15-22.
- Jacobs, J.E., & Simpkins, S.D. (2006). *Leaks in the Pipeline to Math, Science, and Technology Careers*. San Francisco: Jossey-Bass.
- Kenya Wildlife Service (2010). *Mount Kenya National Park/Natural Forest/ Lewa Wildlife Conservancy (extension)*. Retrieved from www.whc.unesco.org/en/tentativelists/5510/
- Kerubo, I.P. (2016). *African indigenous education as practiced by the Maasai of Kenya*. Master's thesis. University of Nairobi.

- Krishnan, R., Gnanaseelan, C., Sanjay, J., Swapna, P., Dhara, C., Sabin, T., & Niyogi, D. (2020). Introduction to climate change over the Indian region. In Krishnan, R. (Eds.), *Assessment of Climate Change over the Indian Region*. Singapore: Springer, pp. 1-20.
- Lindsey R. & Dahlman L. (2021). *Climate Change: Global Temperature*. Retrieved from <https://www.climate.gov/news-features/understanding-climate/climate-change-global-temperature>.
- Maasai Mara National Reserve (2021). *Maasai Mara National Reserve – Welcome to Maasai Mara National Reserve*. Retrieved from <https://www.maasaimarakenyapark.com/>
- Machogu, O.J. (2014). *Assessment of the environmental impacts of wildlife based tourism in Kenya's protected areas: A case study of Maasai Mara National reserve*. Master's Thesis. University of Nairobi.
- Mackey, R., & Currie, D. (2001). The diversity-disturbance relationship: Is it generally strong and peaked? *Ecology*, 82(12): 3479-92.
- Maingi, S.W. (2020). Case Study 6: Mainstreaming overtourism education for sustainable behavioural change in Kenya's tourism industry context. In: Séraphin, H., Gladkikh, T. & Thanh, V. *Overtourism* (pp. 383-99). Cham: Palgrave Macmillan.
- Malhi, Y., Roberts, J.T., Betts, R.A., Killeen, T.J., Li, W., & Nobre, C.A. (2008). Climate change, deforestation, and the fate of the Amazon. *Science*, 319(5860): 169-72.
- Marigi, S.N. (2017). Climate change vulnerability and impacts analysis in Kenya. *American Journal of Climate Change*, 6(01): 52.
- Martin, M.B.G. (2005). Weather, climate and tourism a geographical perspective. *Annals of Tourism Research*, 32(3): 571-91.
- McCright, A. (2010). The effects of gender on climate change knowledge and concern in the American public. *Population and Environment*, 32: 66-87.
- McCarty, J. (2001). Ecological consequences of recent climate change. *Conservation Biology*, 15(2): 320-31.
- Medani M. (2014). Is tourism always beneficial? A case study from Masai Mara National Reserve, Narok, Kenya. *The Pacific Journal of Science and Technology* 5(1):458-483
- Miller, J.D. (2007). *The impact of college science courses for non-science majors on adult scientific literacy*. Paper presented at the 'The Critical role of college science courses for non-majors'-Symposium at the Annual Meeting of the American Association for the Advancement of Science. February 18. San Francisco.
- Montgomery, D.C. (1997). *Design and analysis of experiments* (4th edition). New York: John Wiley and Sons.
- Mugenda, M., & Mugenda G. (2003). *Research Methods: Quantitative and Qualitative Approaches*. Nairobi: ACTS Press.
- Mutekwa, V.T. (2009). Climate change impacts and adaptation in the agricultural sector: The case of smallholder farmers in Zimbabwe. *Journal of Sustainable Development in Africa*, 11(2): 237-56.

- Mutimba, S., Mayieko, S., Olum, P., & Wanyama, K. (2010). *Climate Change Vulnerability and Adaptation Preparedness in Kenya*. Nairobi: Heinrich Böll Stiftung, East and Horn of Africa.
- Mwenda, B., Kiambi, D., Kungu, J., Van De Gevel, J., Farda, C., & Morimoto, Y. (2020). Seasonal climate dynamics, perceptions and multiple risk adaptations: Lessons from smallholder mixed agro ecosystems in Semi-arid Kenya. *Journal of Agricultural Extension and Rural Development*, 12(3): 76-90.
- Nampushi, J., & Nankaya, J. (2020). An assessment of the Maasai residents' views on tourism in the Maasai Mara national reserve, Kenya. *International Journal of Tourism & Hospitality Review*, 7(2): 1-11.
- Nanyingi, M.O., Kiama, S.G., Thumbi, S., & Muchemi, G.M. (2012). *Climate change vulnerability, adaptation and mitigation of livestock systems in Kenya*. Wangari Maathai Institute for Environmental Studies and Peace, College of Agriculture and Veterinary Science, University of Nairobi, Nairobi, Kenya.
- Nash, N., Capstick, S., Whitmarsh, L., Chaudhary, I., & Manandhar, R. (2019). Perceptions of local environmental issues and the relevance of climate change in Nepal's Terai: Perspectives from two communities. *Frontiers in sociology*, 4(60).
- Obeja, R. (2015). *Fatherhood among the Maasai*. Doctoral dissertation. Universitat Internacional de Catalunya.
- Ogutu, J.O, Piepho, H.-P., Said, M.Y., Ojwang, G.O., Njino, L.W., & Kifugo S.C., (2016). Extreme wildlife declines and concurrent increase in livestock numbers in Kenya: What are the causes? *PLoS ONE*, 11(9): e0163249. Retrieved from <https://doi.org/10.1371/journal.pone.0163249>,
- Oso, W.Y., & Onen, D. (2008). *A General Guide to Writing Research Proposals and Reports: A Handbook for Beginning Researchers* (2nd Edn). Kampala: Makerere University Printery.
- Ottichilo, W.K., De Leew, J., Skidmore, A., Prins, H.H., & Said, M.Y. (2000). Population trends of large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya, between 1977 and 1997. *African Journal of Ecology*, 38: 202-16.
- Papageorgiou, M. (2016). Coastal and marine tourism: A challenging factor in Marine Spatial Planning. *Ocean & coastal management*, 129: 44-8.
- Parry, J.E., Echeverria, D., Dekens, J., & Maitima, J. (2012). *Climate risks, vulnerability and governance in Kenya: A review*. Commissioned by: Climate Risk Management Technical Assistance Support Project (CRM TASP), a joint initiative of the Bureau for Crisis Prevention and Recovery and the Bureau for Development Policy of UNDP.
- Pelham, B.W. (2009, April 22nd). *Awareness, opinions about global warming vary worldwide: Many unaware, do not necessarily blame human activities*. Retrieved from <http://www.gallup.com/poll/117772/awareness-opinions-global-warming-vary-worldwide.aspx>

- Ogunleye, K., & Yekinni, O. (2012). Crop farmers' knowledge level of climate change in Ilorin East Local Government Area of Nigeria. *I1STE Journal*. Retrieved from the 10th IPCC (2014): *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. Geneva, Switzerland: IPCC, 151.
- Omoyo, N.N., Wakhungu, J., & Oteng'i, S. (2015). Effects of climate variability on maize yield in the arid and semi-arid lands of lower eastern Kenya. *Agriculture & Food Security*, 4(1): 1-13.
- Oreskes N. (2004). <https://science.sciencemag.org/content/306/5702/1686.full> - aff-1 The Scientific Consensus on Climate Change. *Science*, 306 (5702), p. 1686.
- Parnesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421: 37-42.
- Parnesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Ann Rev Ecol Syst*, 37: 637-69.
- Scheffran, J., Brzoska, M., Brauch, H.G., Link, P.M., & Schilling, J. (Eds.) (2012). *Climate change, human security and violent conflict: Challenges for societal stability* (Vol. 8). Dordrecht: Springer Science & Business Media.
- Schlamadinger, B., Ciccicarese, L., Dutschke, M., Fearnside, P.M., Brown, S., & Murdiyarso, D. (2005). Should we include avoidance of deforestation in the international response to climate change?. In: D. Mudiyarso & H. Herawati(eds.), *Carbon Forestry: Who Will Benefit?* Center for International Forestry Research(CIFOR), Bogor, Indonesia. 215 pp.
- Schmale, J., Zieger, P., & Ekman, A.M. (2021). Aerosols in current and future Arctic climate. *Nature Climate Change*, vol. 11(2): 95-105.
- Serneels, S. & Lambin, E. (2001). Impact of land-use changes on the wildebeest migration in the northern part of the Serengeti–Mara ecosystem. *Journal of Biogeography*, 28, 391-407.
- Shakoor, A., Shahbaz, M., Farooq, T.H., Sahar, N.E., Shahzad, S.M., Altaf, M.M., & Ashraf, M. (2021). A global meta-analysis of greenhouse gases emission and crop yield under no-tillage as compared to conventional tillage. *Science of the Total Environment*, 750: 142299.
- Smith, K. (1993). The influence of weather and climate on recreation and tourism. *Weather*, 48(12): 398-404.
- Sraku-Lartey, M., Buor, D., Adjei, P.O.W., & Foli, E.G. (2020). Perceptions and knowledge on climate change in local communities in the Offinso Municipality, Ghana. *Information Development*, 36(1): 16-35.
- Takakura, H., Fujioka, Y., Ignatyeva, V., Tanaka, T., Vinokurova, N., Grigorev, S., & Boyakova, S. (2021). Differences in local perceptions about climate and environmental changes among residents in a small community in Eastern Siberia. *Polar Science*, 27: 100556.

- Thompson A. (2018). *Plants are the world's dominant life-form*. Retrieved from <https://www.scientificamerican.com/article/plants-are-the-worlds-dominant-life-form/>
- United Nations Education, Scientific and Cultural Organisation (UNEP) (2021). *Climate change could spark floods in world's largest desert lake: new study*. Retrieved from <https://www.unep.org/news-and-stories/story/climate-change-could-spark-floods-worlds-largest-desert-lake-new-study>
- United Nations Education, Scientific and Cultural Organisation (UNESCO) (2021). *The African Great Rift Valley - The Maasai Mara*. Retrieved from <https://whc.unesco.org/en/tentativelists/5512/>
- United States Environmental Protection Agency (USEPA) (2020). *Climate change science - Causes of climate change*. Retrieved from https://19january2017snapshot.epa.gov/climate-change-science/causes-climate-change_.html
- UNICEF (2020). *Troubled waters: Responding to climate change in Turkana*. Retrieved from <https://www.unicef.org/kenya/stories/Responding-to-climate-change-in-Kenya-semi-arid-regions>
- United States Geological Survey (USGS) (2021). *Why is climate change happening and what are the causes?* Retrieved from https://www.usgs.gov/faqs/why-climate-change-happening-and-what-are-causes-1?qt-news_science_products=0#qt-news_science_products
- Van Aalst, M.K., Cannon, T., & Burton, I. (2008). Community level adaptation to climate change: The potential role of participatory community risk assessment. *Global Environmental Change*, 18(1): 165-79.
- World Bank Group (2010). *Kenya's tourism: Polishing the jewel*. Retrieved from <http://documents1.worldbank.org/curated/en/610171468089674111/pdf/696870ESW0P1020e0Jewel0final0report.pdf>
- Zalasiewicz, J., Waters, C., Williams, M., Barnosky, A., Cearreta, A., Crutzen, P., & Oreskes, N. (2015). When did the Anthropocene begin? A mid-twentieth century boundary level is stratigraphically optimal. *Quaternary International*, 383: 196-203.