



LEEDS
BECKETT
UNIVERSITY

Citation:

Roe, G and Sawczuk, T and Tooby, J and Hudson, S and White, R and Mackay, L and Owen, C and Starling, L and Cross, M and Falvey, É and Hendricks, S and Kemp, S and Rasmussen, K and Readhead, C and Salmon, D and Stokes, K and Tucker, R and Jones, B (2024) Training and MatchRelated Head Acceleration Events in Top Level Domestic Senior Women's and Men's Rugby Union: A MultiLeague Instrumented Mouthguard Study. *Scandinavian Journal of Medicine & Science in Sports*, 34 (10). pp. 1-9. ISSN 0905-7188 DOI: <https://doi.org/10.1111/sms.14744>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/11451/>

Document Version:

Article (Published Version)

Creative Commons: Attribution 4.0

© 2024 The Author(s)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.






The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

ORIGINAL ARTICLE OPEN ACCESS

Training and Match-Related Head Acceleration Events in Top Level Domestic Senior Women's and Men's Rugby Union: A Multi-League Instrumented Mouthguard Study

Gregory Roe¹  | Thomas Sawczuk¹ | James Tooby¹  | Sam Hudson^{2,3} | Ryan White¹ | Lois Mackay^{1,4}  | Cameron Owen^{1,5}  | Lindsay Starling⁶ | Matt Cross^{1,7} | Éanna Falvey^{6,8} | Sharief Hendricks^{1,9}  | Simon Kemp^{10,11} | Karen Rasmussen¹² | Clint Readhead^{9,13} | Danielle Salmon⁶ | Keith Stokes^{2,3,11} | Ross Tucker^{6,14} | Ben Jones^{1,5,7,9,15}

¹Carnegie Applied Rugby Research (CARR) Centre, Carnegie School of Sport, Leeds Beckett University, Leeds, UK | ²Centre for Health and Injury and Illness Prevention in Sport, University of Bath, Bath, UK | ³UK Collaborating Centre on Injury and Illness Prevention in Sport (UKCCIIS), University of Bath, Bath, UK | ⁴England Netball, Loughborough, UK | ⁵England Performance Unit, Rugby Football League, Manchester, UK | ⁶World Rugby, Dublin, Ireland | ⁷Premiership Rugby, London, UK | ⁸School of Medicine & Health, University College Cork, Cork, Ireland | ⁹Division of Physiological Sciences and Health Through Physical Activity, Lifestyle and Sport Research Centre, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa | ¹⁰London School of Hygiene and Tropical Medicine, London, UK | ¹¹Rugby Football Union, Twickenham, UK | ¹²New Zealand Rugby Union, People Safety & Wellbeing, Wellington, New Zealand | ¹³South Africa Rugby Union, Cape Town, South Africa | ¹⁴Institute of Sport and Exercise Medicine (ISEM), Department of Exercise, University of Stellenbosch, Stellenbosch, South Africa | ¹⁵School of Behavioural and Health Sciences, Faculty of Health Sciences, Australian Catholic University, Brisbane, Queensland, Australia

Correspondence: Gregory Roe (g.roe@leedsbeckett.ac.uk)

Received: 16 July 2024 | **Revised:** 17 September 2024 | **Accepted:** 27 September 2024

Funding: This work was supported by World Rugby.

ABSTRACT

The aim of this study was to investigate the difference in head acceleration event (HAE) incidence between training and match-play in women's and men's players competing at the highest level of domestic rugby union globally. Players from Women's (Premiership Women's Rugby, Farah Palmer Cup) and Men's (Premiership Rugby, Currie Cup) rugby union competitions wore instrumented mouthguards during matches and training sessions during the 2022/2023 seasons. Peak linear (PLA) and angular (PAA) acceleration were calculated from each HAE and included within generalized linear mixed-effects models. The incidence of HAEs was significantly greater in match-play compared to training for all magnitude thresholds in both forwards and backs, despite players spending approximately 1.75–2.5 times more time in training. For all HAEs (PLA > 5 g and PAA > 400 rad/s²), incidence rate ratios (IRRs) for match versus training ranged from 2.80 (95% CI: 2.38–3.30; men's forwards) to 4.00 (3.31–4.84; women's forwards). At higher magnitude thresholds (PLA > 25 g; PAA > 2000 rad/s²), IRRs ranged from 3.64 (2.02–6.55; PAA > 2000 rad/s² in men's backs) to 11.70 (6.50–21.08; PAA > 2000 rad/s² in women's forwards). Similar trends were observed in each competition. Players experienced significantly more HAEs during match-play than training, particularly at higher magnitude thresholds. Where feasible, HAE mitigation strategies may have more scope for HAE reduction if targeted at match-play, particularly where higher magnitude HAEs are the primary concern. However, the number of HAEs associated with different training drills requires exploration to understand if HAEs can be reduced in training, alongside optimizing match performance (e.g., enhancing contact technique).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Scandinavian Journal of Medicine & Science In Sports* published by John Wiley & Sons Ltd.

1 | Introduction

Rugby union is a contact sport played globally by both men and women [1]. In contact sports, there is concern regarding the potential long-term effects of repetitive head impacts on brain health [2, 3]. Accordingly, it is important for stakeholders to understand the contexts in which head accelerations most commonly occur (e.g., training vs. competition). Such information can be used by researchers and policymakers to appropriately design and implement mitigation strategies [4, 5]. Furthermore, coaches and practitioners may wish to use this data to guide player load management strategies. Finally, this research may provide players with a greater understanding of the head accelerations associated with the activities involved in their sport.

Instrumented mouthguards (iMGs) have been implemented within rugby union to approximate head acceleration events (HAEs) [6]. However, the focus has been primarily on quantifying and characterizing HAEs during match-play [7–9]. A study in amateur men's players identified that the proportion of HAEs experienced during training compared to matches was 35% [7]. Nevertheless, HAEs have not been quantified within an elite training setting. There are differences in physical characteristics [10], greater training loads [11], collision intensities [12] and incidence of head injuries [13] at higher levels of competition compared to amateur settings. Therefore characterizing HAE exposure from training in elite players is a significant gap in the literature. Moreover, there has yet to be any study quantifying HAEs in training in women's players at any playing level. Therefore, the aim of the present study was to quantify and compare the incidence of HAEs in training and match-play in women's and men's players competing at the highest level of domestic rugby union, across a range of competitions globally.

2 | Methods

2.1 | Study Design

A prospective observational study was conducted in rugby union players from two semi-professional women's and two professional men's rugby union competitions who partook in the World Rugby deployment of iMGs (Prevent Biometrics; Minneapolis, MN, USA) during the 2022/2023 seasons. These included the highest level of domestic women's competition in England (Premiership Women's Rugby) and New Zealand (Farah Palmer Cup), and the highest level of domestic men's competition in England (Premiership Rugby) and South Africa (Currie Cup). Players wore iMGs during both rugby training and match-play. For each competition, iMG data were used to determine the HAE incidence in forwards and backs during training and matches. Incidence was expressed using two different exposure time denominators: *per typical match* or *typical training week*, which was based on median match playing and training times, respectively; and *per hour*, which allowed overall comparison with normalized exposure time. The number of players, training sessions, matches and total min per competition and positional group are detailed

in Table 1. Ethics approval was received from the University Ethics Committee (REF: 108638).

2.2 | Instrumented Mouthguards

All players underwent 3D dental scans and were provided with custom-fit iMGs (Prevent Biometrics; Minneapolis). The iMG contained an accelerometer and gyroscope that sampled at 3200 Hz with measured ranges of $\pm 200\text{g}$ and $\pm 35\text{rad/s}$. The coupling of the iMG to the upper dentation was determined by way of infrared proximity sensors. Proximity sensor data were processed in-house by Prevent Biometrics and provided the research team with timestamps of on-the-teeth periods in which the iMG was properly worn by the user. Laboratory validation of the Prevent Biometrics iMG yielded a concordance correlation coefficient of 0.984 (95% CI: 0.977–0.989), while field-based video-verification analysis yielded a positive predictive value of 0.94 (0.92–0.95) and a sensitivity value 0.75 (0.67–0.83) during on-field video-verification validation [14].

A discretized period of kinematics (-10 and $+40\text{ms}$ from the trigger point) was stored for each HAE and linear kinematics were transformed to the estimated head center of gravity (CoG) using the relative acceleration equation (Equation 1). The head was assumed to be a rigid body, therefore angular acceleration values were not transformed. Each HAE was classified as a true positive or false positive by an in-house Prevent Biometrics algorithm based on infrared proximity sensor readings and kinematics. Linear and angular kinematics from all HAEs were initially filtered by Prevent Biometrics using a four-pole, zero phase, low-pass Butterworth filter with a 200 Hz cut-off frequency. Linear acceleration was directly measured by the accelerometer and angular acceleration was calculated by Prevent Biometrics via differentiation of the angular velocity signal recorded by the gyroscope. Another in-house Prevent Biometrics algorithm classified HAEs based on the level of noise in the signal, events classified the level of noise remaining in each signal as low ($n = 70\,635$), moderate ($n = 4153$), or severe ($n = 1542$). An additional filter was applied by Prevent Biometrics to HAEs classified with moderate and severe noise with 100 and 50 Hz cut-off frequencies, respectively. Peak linear acceleration (PLA) and peak angular acceleration (PAA) values were calculated by extracting peak resultant values from each HAE. Following this processing, only HAEs which exceeded both 5g and 400rad/s^2 were used in the analysis, in order to ensure that HAEs included in the data set were a result of contact events, based on previous recommendations [9, 15]. The use of this threshold and Prevent Biometrics' algorithm resulted in a positive predictive value of 0.99 in a previous rugby union study [9], therefore HAEs were not directly video verified for false positives:

$$\vec{a}_h = \vec{a}_m + \vec{\omega} \times r_{mh} + \vec{\omega} \times (\vec{\omega} \times r_{mh}) \quad (1)$$

The relative acceleration equation, where \vec{a}_h is the linear acceleration at the head CoG with respect to time, \vec{a}_m is the linear acceleration at the iMG sensor location with respect to time, $\vec{\omega}$ is the angular acceleration with respect to time, r_{mh} is the position vector from iMG sensor location to the head CoG, and $\vec{\omega}$ is angular velocity with respect to time. The transformation vector (r_{mh}) was $(-0.082, 0.009, -0.065)$ in meters.

TABLE 1 | The number of players, player matches (PM)/training weeks (PTW), total hours, median training/playing time (interquartile range; IQR) and HAEs (all, > 25 g, > 2000 rad/s²) for training and matches per position and competition. W = women's competition, M = men's competition.

| | Players | Player matches/ player training weeks | Total player hours | HAEs | | | |
|----------------------------------|---------|---|--------------------------|--|-----------------|------------|----------------------------------|
| | | | | Median playing/ weekly training time (IQR) | All recorded | PLA > 25 g | PAA > 2000 rad/s ² |
| Matches (forwards) | | | | | | | |
| Women's: all competitions | 269 | 1038 (PM) | 892 | 58 (27–80) | 12930 | 1173 | 1142 |
| Premiership Women's Rugby (W) | 114 | 577 (PM) | 505 | 60 (27–80) | 7620 | 705 | 727 |
| Farah Palmer Cup (W) | 155 | 461 (PM) | 387 | 56 (27–76) | 5310 | 468 | 415 |
| Men's: all competitions | 238 | 1567 (PM) | 1306 | 56 (24–77) | 38305 | 4795 | 3592 |
| Premiership Rugby (M) | 143 | 1097 (PM) | 933 | 57 (26–80) | 27653 | 3460 | 2558 |
| Currie Cup (M) | 95 | 470 (PM) | 373 | 52 (22–70) | 10652 | 1335 | 1034 |
| Matches (backs) | | | | | | | |
| Women's: all competitions | 180 | 684 (PM) | 687 | 74 (40–80) | 6187 | 633 | 637 |
| Premiership Women's Rugby (W) | 71 | 355 (PM) | 357 | 80 (40–80) | 2992 | 259 | 294 |
| Farah Palmer Cup (W) | 109 | 329 (PM) | 329 | 71 (45–80) | 3195 | 374 | 343 |
| Men's: all competitions | 141 | 863 (PM) | 848 | 72 (40–80) | 14770 | 1838 | 1658 |
| Premiership Rugby (M) | 81 | 564 (PM) | 563 | 73 (42–80) | 10459 | 1328 | 1164 |
| Currie Cup (M) | 60 | 299 (PM) | 285 | 71 (28–80) | 4311 | 510 | 494 |
| Training (forwards) | | | | | | | |
| Women's: all competitions | 156 | 358 (PTW) | 538 | 143 (132–162) | 914 | 42 | 36 |
| Premiership Women's Rugby (W) | 31 | 68 (PTW) | 84 | 113 (99–138) | 162 | 9 | 12 |
| Farah Palmer Cup (W) | 125 | 290 (PTW) | 454 | 144 (133–166) | 752 | 33 | 24 |
| Men's: all competitions | 141 | 359 (PTW) | 503 | 129 (114–138) | 2429 | 168 | 127 |
| Premiership Rugby (M) | 90 | 239 (PTW) | 335 | 128 (122–138) | 1834 | 124 | 94 |
| Currie Cup (M) | 51 | 120 (PTW) | 168 | 149 (97–153) | 595 | 44 | 33 |
| Training (backs) | | | | | | | |
| Women's: all competitions | 92 | 190 (PTW) | 286 | 144 (134–166) | 401 | 18 | 25 |
| Premiership Women's Rugby (W) | 18 | 36 (PTW) | 38 | 138 (113–138) | 41 | 0 | 1 |
| Farah Palmer Cup (W) | 74 | 154 (PTW) | 248 | 162 (141–166) | 360 | 18 | 24 |
| Men's: all competitions | 58 | 111 (PTW) | 144 | 126 (112–135) | 394 | 27 | 40 |
| Premiership Rugby (M) | 39 | 79 (PTW) | 105 | 128 (114–135) | 316 | 23 | 35 |
| Currie Cup (M) | 19 | 23 (PTW) | 39 | 97 (97–125) | 78 | 4 | 5 |

Note: A player match is the product of the number of matches in the study and the number of players participating in them within each position-competition group (e.g., 269 women's forwards who participated in 1038 matches between them). Player training week is the product of the number of training weeks involved in the study and the number of players participating in them within each position-competition group (e.g., 141 men's forwards who participated in 359 training weeks between them). Total player hours is the accumulated total number of hours played/trained across all players within each position-competition group.

2.3 | Training Data

Due to the limited number of camera angles typically used to film rugby union training, it was not possible to visually inspect if all players were wearing their iMGs at all appropriate times during training (e.g., as a minimum iMG on-teeth during contact and off-teeth during the rest period within a drill). However, in a subsample of players from Premiership Rugby (men, $n = 17$, 11 forwards, 6 backs across 3 clubs) and Premiership Women's Rugby (women, $n = 24$, 16 forwards and 8 backs across 3 clubs), video footage was available in which each player was identifiable throughout a training session. Across 17 training sessions and 71 player-training-session observations, for all drills in which an iMG was worn, players only removed their iMG in between training activities or when they were not directly involved in the training activity in which there was the potential for contact (e.g., subbing in and out, or between repetitions during a training drill). Based on these findings, it was assumed that all players wore their iMG during training activities in which they were at risk of being involved in a contact event. However, given that the subsample was from a single country, it is possible that these practices were not ubiquitous across all competitions.

For each competition, weekly training video footage was requested from all participating clubs. Video data for a given training session were included in the study if an accurate start timestamp was provided, and the content of all drills could be discerned (i.e., no drills were undertaken off camera). For each training session, the start and stop times and contact level (classified using the rugby football union's definitions of contact training, unpublished) of each drill were coded by two experienced analysts, reaching a total of 325h of video footage analyzed. Only iMG data occurring on the date of the training session, between the start and stop times of each session were included in the training dataset. Accelerometer, gyroscope and proximity sensor data recorded by iMGs were synchronized to video footage and visualized using Matlab (MathWorks, UK, version R2023a). Three authors visually inspected proximity sensor data to determine whether players wore their iMG for each drill. A player's data were only included if they wore an iMG in all training drills on a given day with the exception of non-contact training.

Players typically only wore their iMGs during the main training days of the week during which contact training is typically undertaken (i.e., no training sessions 24h prior to a match, or the first training session after a match within 48h). Therefore, only data from the main training days of the week were included in the study. The total number of training weeks and minutes for each competition used in the final analysis are detailed in Table 1.

2.4 | Match-Data

Video analysis provided by Opta (StatsPerform, Chicago, IL, USA) were used to identify match exposure for each player-match. Only iMG data recorded by players participating in the match, on the date of the match, and between each player's start and stop playing time were included in the match dataset. Accelerometer, gyroscope, and proximity sensor data were synchronized to Opta data for each match, which contained timestamps of all contact

events (tackles, ball carries, rucks, mauls, and scrums). Proximity sensor data were used to identify whether the iMG was being worn for each contact-event and only player matches where the instrumented player wore their iMG for at least 90% of their contact events were included to ensure iMG adherence and minimize false negatives [8, 9].

2.5 | Statistical Analysis

To quantify and compare the HAE incidence in training and match-play for women's and men's players, generalized linear mixed models assuming a Poisson distribution were used. Sex, type (training or match play), position (forward or back) and the logarithm of minutes played were included as fully factorial fixed effects allowing slopes of HAEs accumulation per minutes to be estimated for each sex, type, and position combination. Random slopes for the number of HAEs per minute of exposure time were added for each athlete and allowed to vary between training and match play. A random effect for the week, clustering players within matches and training weeks together was also included. All statistical analyses were completed in R (version 4.3.2) using the *glmmTMB* [16] and *emmeans* [17] packages.

Incidence was estimated with a predicted function using each generalized linear mixed model using the various denominators. These denominators included *per typical match* or *per typical training week*, whereby the median minutes played per player-match and median minutes trained per week were used, respectively; and *per training hour* or *per match hour*, to allow comparison using a normalized exposure time. Median minutes played (used for the *per typical match* denominator) was calculated using playing time provided for each player-match in Opta data, while median minutes trained (used for the *per typical training week*) was calculated using only those players with two or more sessions in the training week. These median exposure times were calculated separately for forwards and backs, and men's and women's players and are provided in Table 1.

In addition to different denominators, incidence was expressed using different thresholds. Firstly, the incidence of *all HAEs* (i.e., those measured by the iMG which exceeded both 5g and 400 rad/s²) was used to estimate the exposure of HAEs exceeding those usually measured from non-contact events [9, 15]. Additionally, the incidence was calculated using higher HAE thresholds: separately, HAEs exceeding 25g, and HAEs exceeding 2000 rad/s² were used to capture the incidence of higher magnitude HAEs. Thresholds beyond this did not reach model convergence. Incidence rate data are visualized in Figures 1 and 2, and provided in numerical form in Table S1.

Incidence rate ratios were used to compare rates between training and match play within positions and sexes (e.g., men's forwards *incidence per match hour* vs. men's forwards *incidence per training hour*). Results are presented as means (95% confidence intervals). In situations where very limited data were present around the median number of minutes played, or very few HAEs were present, confidence intervals were likely to be inflated.




FIGURE 1 | Head acceleration event incidence for all women's players and in each women's competition per median training/playing time (A, C, E) and per hour (B, D, F). Position is represented by the absence (forwards) or presence (backs) of dots. NB: No HAEs for PLA > 25g and only one HAE > 2000 rad/s² were recorded during Premiership Women's Rugby training, therefore robust confidence intervals could not be calculated.

Comparisons were deemed significantly different if confidence limits did not overlap or did not contain one [18].

3 | Results

3.1 | Incidence of HAEs in Women's Training and Match-Play

The HAE incidence for players was significantly greater in a typical match compared to a typical training week for all magnitude thresholds (Figures 1 and 2; A,C and E). For all HAEs, differences between a typical training week and a typical match, as measured by incidence rate ratios (Table 2), ranged from 2.80 (2.38–3.30; men's forwards) to 4.00 (3.31– 4.84; women's forwards). At higher magnitudes, incidence rate ratios were larger, ranging from 3.64 (2.02–6.55; HAEs > 2000 rad/s²; men's backs) to 11.70 (6.50–21.08); HAEs > 2000 rad/s²; women's forwards depending on the HAE threshold, sex and positional group (Table 2). When normalized per hour, differences between training and match-play were greater than per typical training week/match, ranging from 6.88 (6.01–7.88; all HAEs; men's forwards) to 14.24 (7.62–26.62; PLA > 25g; men's backs). Similar trends were seen in individual competitions. Figure 3 depicts all




FIGURE 2 | Head acceleration event incidence for all men's players and in each men's competition per median training/playing time (A, C, E) and per hour (B, D, F). Position is represented by the absence (forwards) or presence (backs) of dots.

head kinematic values for training and match-play for women's and men's players.

4 | Discussion

The aim of the present study was to quantify and compare the incidence of HAEs in training and match-play in women's and men's players competing at the highest level of domestic rugby union, across a range of competitions globally. The HAE incidence for players was significantly greater in a typical match compared to a typical training week for all magnitude thresholds, with incidence rate ratios ranging from 2.80 (2.38–3.30; all HAEs; men's forwards) to 11.70 (6.50–21.08); HAEs > 2000 rad/s²; women's backs depending on the HAE threshold, sex and positional group. These ratios increased when training and match-play were normalized per hour. Similar trends were observed in individual competitions. Where feasible, mitigation strategies designed to reduce players' HAE exposure may have greater scope to do so if targeted at match-play (e.g., rules changes and player match/HAE limits), particularly where higher HAE magnitudes are of primary concern.

In both women's and men's players, HAE incidence for forwards and backs was significantly greater in a typical match than in a typical training week (Figures 1 and 2; A,C and E), despite

TABLE 2 | Incidence rate ratios (confidence intervals) for training compared to match-play per median training/playing times and per hour. Training is used as the reference value; for example, HAEs exceeding 25 g were 5.25 times more numerous in a typical match than in a typical training week in Premiership Rugby forwards, while the incidence of HAEs per hour was 9.03 higher during match-play than training in that competition.

| | Per typical weekly training/playing time | | Per hour training/match-play | |
|-------------------------------|--|--------------------|------------------------------|--------------------|
| | Forwards | Backs | Forwards | Backs |
| Sex | | | | |
| Women's | | | | |
| All HAEs | 4.00 (3.31–4.84) | 3.03 (2.37–3.87) | 7.41 (6.31–8.70) | 8.76 (6.91–11.11) |
| PLA > 25 g | 8.20 (4.87–13.82) | 9.04 (4.18–19.55) | 13.39 (8.28–21.67) | 13.24 (6.28–27.90) |
| PAA > 2000 rad/s ² | 11.70 (6.50–21.08) | 6.65 (3.42–12.93) | 12.49 (7.80–20.00) | 9.88 (5.17–18.88) |
| Men's | | | | |
| All HAEs | 2.80 (2.38–3.30) | 3.40 (2.56–4.51) | 6.88 (6.01–7.88) | 7.02 (5.65–8.71) |
| PLA > 25 g | 4.98 (3.68–6.73) | 4.64 (2.43–8.87) | 11.60 (8.77–15.35) | 14.24 (7.62–26.62) |
| PAA > 2000 rad/s ² | 6.60 (4.65–9.36) | 3.64 (2.02–6.55) | 8.98 (6.78–11.89) | 7.34 (4.54–11.86) |
| Competition | | | | |
| Premiership Women's Rugby (W) | | | | |
| All HAEs | 5.30 (3.63–7.76) | 2.43 (1.18–4.99) | 6.37 (4.84–8.39) | 10.58 (6.54–17.13) |
| PLA > 25 g | 8.92 (3.13–25.41) | NA | 10.59 (4.98–22.52) | NA |
| PAA > 2000 rad/s ² | 8.87 (3.20–24.55) | NA | 7.72 (4.06–14.68) | NA |
| Farah Palmer Cup (W) | | | | |
| All HAEs | 3.42 (2.76–4.24) | 2.82 (2.09–3.80) | 7.77 (6.33–9.53) | 8.05 (6.06–10.70) |
| PLA > 25 g | 7.07 (3.98–12.54) | 11.13 (4.19–29.55) | 14.92 (7.98–27.91) | 10.54 (4.65–23.90) |
| PAA > 2000 rad/s ² | 8.84 (4.61–16.96) | 6.87 (2.98–15.86) | 17.46 (8.55–35.65) | 7.81 (3.71–16.43) |
| Premiership Rugby (M) | | | | |
| All HAEs | 2.58 (2.11–3.14) | 2.69 (1.95–3.70) | 6.08 (5.16–7.15) | 6.76 (5.23–8.74) |
| PLA > 25 g | 5.25 (3.64–7.58) | 3.83 (1.90–7.69) | 9.03 (6.62–12.34) | 13.07 (6.46–26.45) |
| PAA > 2000 rad/s ² | 6.53 (4.31–9.92) | 2.91 (1.53–5.52) | 7.48 (5.44–10.27) | 5.76 (3.43–9.70) |
| Currie Cup (M) | | | | |
| All HAEs | 2.52 (1.84–3.44) | 6.31 (3.90–10.20) | 8.93 (7.06–11.29) | 7.01 (4.71–10.42) |
| PLA > 25 g | 2.95 (1.62–5.38) | 12.65 (3.13–51.07) | 26.09 (14.02–48.53) | 17.70 (4.74–66.13) |
| PAA > 2000 rad/s ² | 4.87 (2.31–10.23) | 8.43 (2.79–25.49) | 15.99 (8.66–29.50) | 19.31 (5.07–73.56) |

Note: No HAEs for PLA > 25 g and only one HAE > 2000 rad/s² were recorded during Premiership Women's Rugby training. Therefore, incidence rate ratios were not calculated for this competition at these thresholds.

Abbreviations: M, men's competition; W, women's competition.

players spending approximately 1.75–2.5 times more time training (Table 1). When normalized per hour (e.g., approximately half of the median training times, Table 1), HAE incidence rate ratios ranged from 7.41 (6.31–8.70; all HAEs in forwards) to 13.39 (8.28–21.67; PLA > 25 g in forwards) in women's players, and from 6.88 (6.01–7.88; all HAEs in forwards) to 14.24 (7.62–26.62; PLA > 25 g in backs) in men's players. In practical terms, this means that players in the present study would have to train between approximately 7–14 h to experience a similar number of HAEs in training as in an hour of match-play, depending on their sex and position. This is far greater than the median training times of the teams in the current study (97–149 mins, Table 2).

At higher HAE magnitude thresholds (PLA > 25 g; PAA > 2000 rad/s²), the difference between training and match-play increased (Table 2). However, it is important to note that the absolute number of HAEs experienced by players decreased (Figure 1), confirming previous findings in elite-level match-play [8, 9]. For example, although HAE incidence for women's forwards was 8.20 times greater at PLA > 25 g in a typical match than a typical training week (Table 2), in absolute terms, on average forwards had only 0.13 HAEs in a typical training week (median of 129 min training time) and 1.04 HAEs during a typical match (median of 58 min of match-play) at this magnitude threshold. Nevertheless, across a competitive season or a playing




FIGURE 3 | Head kinematic values during training (black dots) and match-play (red dots) for men's and women's players.

career, these differences would have a significant impact on player exposure. The clinical relevance of exposure to HAEs at different magnitude threshold ranges needs to be determined before conclusions can be made regarding the pertinence of these findings to player safety.

In the present study, for most competitions and playing positions, significantly more data (2–9-fold more) were collected from match-play than from training (Table 1). In a recent study investigating iMG usage in professional rugby, iMG managers (appointed practitioners responsible for the collection and analysis of iMG data in sports teams) reported that players did not habitually wear mouthguards during training (instrumented or otherwise) [19]. This represented a major barrier to players wearing iMGs, and player and staff engagement with iMG data. Furthermore, in the present study, the quality of the training video data provided by teams in certain competitions was not always of a standard that could be used. Due to the limited data, incidence rate ratios could not be calculated for Premiership Women's Rugby at higher thresholds as there was only 1 HAE > 2000 rad/s² and no HAE > 25 g, while the incidence rate ratio confidence limits calculated for Currie Cup backs were very wide (Table 2). Further research is therefore required to understand if the HAE incidence estimations in the present study accurately represent these two playing populations.

The differences between training and match-play identified in this study have important implications for player welfare policy. In other collision sports, for example, American Football, research has demonstrated significant associations between retrospective estimations of cumulative head impacts throughout players' careers and chronic traumatic encephalopathy [3] and neurocognitive impairments in later life [20]. Thus, interventions

to reduce player exposure to HAEs are of high priority. In contact sports such as American Football and ice hockey, observational studies have identified training activities (e.g., full-contact [21]) and match actions (e.g., longer closing distances [22], contact technique [23], and illegal contacts [24]) that result in higher magnitude HAEs. Recently in rugby union, Roe et al. [8] observed that tackling, followed by ball carrying, had the highest probability of resulting in a HAE regardless of magnitude. To date, however, the evaluation of policy changes to enhance player safety with respect to head injury has primarily focused on concussion. For example, rule changes such as the "targeting rule" in the NFL [25] and the outlawing of body checking in youth ice hockey [26], have reduced concussion incidence, while in rugby union, changes to the tackle height in semi-professional [27] and amateur [28] players did not. It has yet to be ascertained whether such interventions would have had a significant impact on HAE incidence in these populations.

In intervention design, an important initial step is to develop an understanding of the problem requiring intervention and identify the factors that are most amenable to change [29]. In the present study, on average, both women's and men's rugby union players experienced a significantly greater number of HAEs during match-play than training. Therefore, interventions targeting match-play, such as reduced maximum match limits for players in a season and/or law change to change the frequency and/or nature of match contact events may have greater scope for reducing long-term player HAE exposure than those targeting training (restricting the time spent on selected training activities). However, to support the design of such interventions additional research is required to determine the most feasible and appropriate ways to do so (e.g., investigating the HAE causal mechanisms on which to intervene [5], assessing the appropriateness and feasibility of rule changes while considering player

and fan viewpoints, optimal match limits [30] and training activity exposure [31]). Moreover, the acute and chronic clinical significance of HAE exposure has not been prospectively determined. These are important considerations for whether policy changes should target activities that result in HAEs of specific magnitudes, or HAEs overall.

4.1 | Limitations

Although this study provides novel insights, it has some limitations. The first of these, consistent with similar studies [7–9], is sampling bias. This is present in the form of non-random sampling (a convenience sample from each competition was used) and volunteer bias (only players who volunteered were included). The generalizability of the results in the present study is conditional on the sample of players being representative of the population more broadly. For example, those that volunteered to wear an iMG may have done so because they are involved in more contact events, which could bias the findings by overestimating HAE incidence. Given that the wearing of iMGs during matches and contact training has recently been mandated, future research should aim to include a broad range of players, playing positions, ages and other potentially important factors. Additionally, the number of training sessions in the data set was much lower than the number of matches, particularly in Premiership Women's Rugby and Currie Cup men's backs. Therefore, HAE incidence estimates in training may not be fully representative of these playing populations. Furthermore, due to players not habitually wearing iMGs during certain training sessions (match-day + 2, match-day–1), only data from the main training sessions of the week were available. Although contact training is not typically undertaken in sessions other than those quantified in the present study, it is possible that players may have experienced some HAEs during these other training sessions. Moreover, the sub-components of training were not investigated in this study, nor were the mechanisms of HAE. It is likely that certain activities within training (e.g., full-contact training) contribute more to HAE incidence than others. Thus, it is important that future research investigates the HAEs associated with different training activities to determine if mitigation strategies targeting training may also be appropriate and feasible. However, careful consideration is warranted to ensure that strategies to reduce HAEs in training do not interfere with the preparation of players, particularly given the importance of technical proficiency for safe tackling and ball-carrying in match-play [32]. Finally, attempts were made to model higher HAE magnitudes thresholds in accordance with other rugby union literature [8]. However, due to the lack of data in training versus matches for both men (e.g., >40g; 1685 HAEs in match-play and 23 in training) and women (e.g., >40g; 399 in matches and 9 in training), models did not converge. Future research may require larger data sets and advanced statistical models to cater for the rarity of these events in training.

4.2 | Perspective

This is the first study to quantify the differences in HAE incidence between training and match-play in women's and men's

players competing at the highest level of domestic competition. The HAE incidences for women's and men's players were significantly greater in match-play compared to training for all magnitude thresholds in both forwards and backs, despite players spending longer (approximately 1.75–2.5 times more) in training than in matches. Differences in HAE incidence ranged from 2.80 to 11.70 times greater in matches than in training, depending on the HAE threshold, sex and positional group; and were even larger when training and matches were compared per hour. Given that long-term exposure to HAEs in rugby union players is a health concern, mitigation strategies targeting match-play may have greater scope to reduce HAE exposure, although future research is required to determine the most feasible and appropriate ways to do so. Furthermore, the number of HAEs associated with different types of training drills also needs to be explored to understand if HAEs can be reduced in training, alongside optimizing match performance (e.g., enhancing contact technique).

Author Contributions

G.R., L.S., M.C., K.S., and B.J. conceptualized the research. G.R., D.S., J.T., and S.Hu. collected the data. J.T., T.S., G.R., S.Hu., R.W., and L.M. analyzed the data while all authors were involved in the interpretation of the results. G.R. drafted the research paper and all authors contributed equally to the writing beyond this point.

Conflicts of Interest

T.S. role is part-funded by Premiership Rugby. G.R. role is part-funded by World Rugby. J.T. role is part-funded by the Rugby Football League, Premiership Rugby and World Rugby. C.O. is part-funded by the Rugby Football League. B.J. is employed by Premiership Rugby and Rugby Football League as a consultant and has received funding from Prevent Biometrics and World Rugby. M.C. is employed by Premiership Rugby. L.S., É.F. and D.S. are employed by World Rugby. C.R. is employed by the South African Rugby Union. K.R. is employed by New Zealand Rugby. S.H. is funded by the Rugby Football Union. K.S. and S.K. are employed by the Rugby Football Union. R.T. is employed by World Rugby as a consultant. S.H., R.W., and L.M. have no conflicts of interest.

Data Availability Statement

Data is available upon reasonable request from the corresponding author, Gregory Roe.

References

- O. Heyward, S. Emmonds, G. Roe, S. Scantlebury, K. Stokes, and B. Jones, "Applied Sports Science and Sports Medicine in Women's Rugby: Systematic Scoping Review and Delphi Study to Establish Future Research Priorities," *BMJ Open Sport & Exercise Medicine* 8, no. 3 (2022): e001287.
- G. L. Iverson, R. J. Castellani, J. D. Cassidy, et al., "Examining Later-in-Life Health Risks Associated With Sport-Related Concussion and Repetitive Head Impacts: A Systematic Review of Case-Control and Cohort Studies," *British Journal of Sports Medicine* 57, no. 12 (2023): 810–824.
- D. H. Daneshvar, E. S. Nair, Z. H. Baucom, et al., "Leveraging Football Accelerometer Data to Quantify Associations Between Repetitive Head Impacts and Chronic Traumatic Encephalopathy in Males," *Nature Communications* 14, no. 1 (2023): 3470.
- C. Finch, "A New Framework for Research Leading to Sports Injury Prevention," *Journal of Science and Medicine in Sport* 9, no. 1–2 (2006): 3–9, discussion 10.

5. D. Wight, E. Wimbush, R. Jepson, and L. Doi, "Six Steps in Quality Intervention Development (6SQuID)," *Journal of Epidemiology and Community Health* 70, no. 5 (2016): 520–525.
6. J. Tooby, K. Till, A. Gardner, et al., "When to Pull the Trigger: Conceptual Considerations for Approximating Head Acceleration Events Using Instrumented Mouthguards," *Sports Medicine* 54, no. 6 (2024): 1361–1369.
7. M. D. Bussey, D. Salmon, J. Romanchuk, et al., "Head Acceleration Events in Male Community Rugby Players: An Observational Cohort Study Across Four Playing Grades, From Under-13 to Senior Men," *Sports Medicine* 54, no. 2 (2024): 517–530.
8. G. Roe, T. Sawczuk, C. Owen, et al., "Head Acceleration Events During Tackle, Ball-Carry, and Ruck Events in Professional Southern Hemisphere Men's Rugby Union Matches: A Study Using Instrumented Mouthguards," *Scandinavian Journal of Medicine & Science in Sports* 34, no. 6 (2024): e14676.
9. J. Tooby, J. Woodward, R. Tucker, et al., "Instrumented Mouthguards in Elite-Level Men's and Women's Rugby Union: The Incidence and Propensity of Head Acceleration Events in Matches," *Sports Medicine* 54, no. 5 (2023): 1327–1338.
10. D. J. Smart, W. G. Hopkins, and N. D. Gill, "Differences and Changes in the Physical Characteristics of Professional and Amateur Rugby Union Players," *Journal of Strength & Conditioning Research* 27, no. 11 (2013): 3033–3044.
11. J. Al-Attar, H. Croft, K. Bloxham, and M. R. Blair, "Training Load and Its Progression From Amateur to Semi-Professional Rugby," *Journal of Australian Strength and Conditioning* 28, no. 7 (2020): 6–15.
12. P. Tierney, C. Blake, and E. Delahunt, "Physical Characteristics of Different Professional Rugby Union Competition Levels," *Journal of Science and Medicine in Sport* 24, no. 12 (2021): 1267–1271.
13. S. Williams, C. Robertson, L. Starling, et al., "Injuries in Elite Men's Rugby Union: An Updated (2012–2020) Meta-Analysis of 11 620 Match and Training Injuries," *Sports Medicine* 52, no. 5 (2022): 1127–1140.
14. B. Jones, J. Tooby, D. Weaving, et al., "Ready for Impact? A Validity and Feasibility Study of Instrumented Mouthguards (iMGs)," *British Journal of Sports Medicine* 56, no. 20 (2022): 1171–1179.
15. J. Tooby, J. Woodward, and G. Tierney, "Quantifying and Characterising Head Kinematics From Non-Contact Events Using Instrumented Mouthguards," *IRCOBI Conference Proceedings* (Conference proceedings International Research Council on the Biomechanics of Injury, IRCOBI, 2022), 624–625.
16. M. E. Brooks, K. Kristensen, K. J. Va Benthem, et al., "GlmTMB Balances Speed and Flexibility Among Packages for Zero-Inflated Generalized Linear Mixed Modeling," *R Journal* 9, no. 2 (2017): 378–400.
17. R. Lenth, "Emmeans: Estimated Marginal Means, aka Least-Squares Means," R package version 1.8.7 (2023), <https://CRAN.R-project.org/package=emmeans>.
18. K. Noguchi and F. Marmolejo-Ramos, "Assessing Equality of Means Using the Overlap of Range-Preserving Confidence Intervals," *American Statistician* 70, no. 4 (2016): 325–334.
19. G. Roe, S. Whitehead, L. Starling, et al., "Embracing the Impact From Instrumented Mouthguards (IMGs): A Survey of IMG Managers' Perceptions of Staff and Player Interest Into the Technology, Data and Barriers to Use," *European Journal of Sport Science* 24, no. 6 (2024): 670–681.
20. P. H. Montenegro, M. L. Alosco, B. M. Martin, et al., "Cumulative Head Impact Exposure Predicts Later-Life Depression, Apathy, Executive Dysfunction, and Cognitive Impairment in Former High School and College Football Players," *Journal of Neurotrauma* 34, no. 2 (2017): 328–340.
21. S. P. Broglio, D. Martini, L. Kasper, J. T. Eckner, and J. S. Kutcher, "Estimation of Head Impact Exposure in High School Football: Implications for Regulating Contact Practices," *American Journal of Sports Medicine* 41, no. 12 (2013): 2877–2884.
22. K. E. Ocwieja, J. P. Mihalik, S. W. Marshall, J. D. Schmidt, S. C. Trulock, and K. M. Guskiewicz, "The Effect of Play Type and Collision Closing Distance on Head Impact Biomechanics," *Annals of Biomedical Engineering* 40, no. 1 (2012): 90–96.
23. J. Alois, S. Bellamkonda, E. T. Campolettano, et al., "Do American Youth Football Players Intentionally Use Their Heads for High-Magnitude Impacts?," *American Journal of Sports Medicine* 47, no. 14 (2019): 3498–3504.
24. J. P. Mihalik, R. M. Greenwald, J. T. Blackburn, R. C. Cantu, S. W. Marshall, and K. M. Guskiewicz, "Effect of Infraction Type on Head Impact Severity in Youth Ice Hockey," *Medicine and Science in Sports and Exercise* 42, no. 8 (2010): 1431–1438.
25. J. M. May, H. S. Angileri, D. E. McLoughlin, M. M. Owen, M. Terry, and V. Tjong, "Decreased Concussion Incidence Following the Implementation of the Targeting Rules: An Updated Epidemiology of National Football League Concussions From 2017 to 2022," *Cureus* 15, no. 12 (2023): e50997.
26. C. A. Emery, P. Eliason, J.-M. Galarneau, et al., "Body Checking in Non-Elite Adolescent Ice Hockey Leagues: It Is Never Too Late for Policy Change Aiming to Protect the Health of Adolescents," *British Journal of Sports Medicine* 56, no. 1 (2022): 12–18.
27. K. A. Stokes, D. Locke, S. Roberts, et al., "Does Reducing the Height of the Tackle Through Law Change in Elite Men's Rugby Union (The Championship, England) Reduce the Incidence of Concussion? A Controlled Study in 126 Games," *British Journal of Sports Medicine* 55, no. 4 (2021): 220–225.
28. T. Rv, L. Starling, S. Surmon, et al., "Tackling Sport-Related Concussion: Effectiveness of Lowering the Maximum Legal Height of the Tackle in Amateur Male Rugby—A Cross-Sectional Analytical Study," *Injury Prevention* 29, no. 1 (2023): 56–61.
29. A. O' Cathain, L. Croot, E. Duncan, et al., "Guidance on How to Develop Complex Interventions to Improve Health and Healthcare," *BMJ Open* 9, no. 8 (2019): e029954.
30. T. Sawczuk, M. Cross, C. Owen, et al., "The Application of Match-Event and Instrumented Mouthguard (IMG) Data to Inform Match Limits: An Example Using Rugby Union Premiership and Rugby League Super League Data From England," *European Journal of Sport Science*. Published ahead of print, September 21, 2024. <https://doi.org/10.1002/ejsc.12188>.
31. L. T. Starling, R. Tucker, K. Quarrie, et al., "The World Rugby and International Rugby Players Contact Load Guidelines: From Conception to Implementation and the Future," *South African Journal of Sports Medicine* 35, no. 1 (2023): v35i1a16376.
32. V. Meintjes, P. Forshaw, S. den Hollander, et al., "Tackler and Ball-Carrier Technique During Moderate and Severe Injuries (≥ 8 Days Lost) Compared With Player-Matched and Team-Matched Injury-Free Controls in Elite Rugby Union," *British Journal of Sports Medicine* 55, no. 24 (2021): 1411–1419.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.