



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Peart, DJ and Graham, M and Lunn, DE and Burn, N and Franklin, JD and Arnold, JB (2025) Editorial: Remote assessment, measurement, and delivery in sport, physical activity and health. *Frontiers in Sports and Active Living*, 7. pp. 1-3. ISSN 2624-9367 DOI: <https://doi.org/10.3389/fspor.2025.1584714>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/11938/>

Document Version:

Article (Published Version)

---

Creative Commons: Attribution 4.0

© 2025 Peart, Graham, Lunn, Burn, Franklin and Arnold

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.



## OPEN ACCESS

EDITED AND REVIEWED BY  
Kamiar Aminian,  
Swiss Federal Institute of Technology  
Lausanne, Switzerland

## \*CORRESPONDENCE

Daniel J. Peart  
✉ daniel.peart@northumbria.ac.uk  
John D. Franklin  
✉ j.franklin@tees.ac.uk

RECEIVED 27 February 2025

ACCEPTED 04 March 2025

PUBLISHED 14 March 2025

## CITATION

Peart DJ, Graham M, Lunn DE, Burn N,  
Franklin JD and Arnold JB (2025) Editorial:  
Remote assessment, measurement, and  
delivery in sport, physical activity and health.  
*Front. Sports Act. Living* 7:1584714.  
doi: 10.3389/fspor.2025.1584714

## COPYRIGHT

© 2025 Peart, Graham, Lunn, Burn, Franklin  
and Arnold. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Editorial: Remote assessment, measurement, and delivery in sport, physical activity and health

Daniel J. Peart<sup>1\*</sup>, Michael Graham<sup>2</sup>, David E. Lunn<sup>3</sup>, Naomi Burn<sup>4</sup>,  
John D. Franklin<sup>2\*</sup> and John B. Arnold<sup>4</sup>

<sup>1</sup>Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle upon Tyne, United Kingdom, <sup>2</sup>School of Health and Life Sciences, Teesside University, Middlesbrough, United Kingdom, <sup>3</sup>Carnegie School of Sport, Leeds Beckett University, Leeds, United Kingdom, <sup>4</sup>Alliance for Research in Exercise, Nutrition and Activity (ARENA), University of South Australia, Adelaide, SA, Australia

## KEYWORDS

remote assessment, measurement, sport, physical activity, health

## Editorial on the Research Topic

**Remote assessment, measurement, and delivery in sport, physical activity and health**

## Introduction

Technology has had an increasing influence on our lives for some time, and this was accelerated by the enforced remote way of working five years ago. The first paper in this special topic from [Chahin-Inostroza et al.](#) articulates some of the challenges faced during this time as countries across the world enforced distancing measures and remote working to control the spread of COVID-19. Sport was not exempt, and both national and global events were put on standby. However, this not only affected elite athletes, but amateur athletes too had fewer opportunities to socialise with peers and receive feedback and advice. In their cross-section of amateur Chilean athletes, [Chahin-Inostroza et al.](#) report how the use of some technologies and training software changed during periods of quarantine, and remained elevated when quarantine ended.

The increase in remote technology acceptance and usage in sport, health and exercise could present an opportunity to move interventions and data collection out of the laboratory or any other fixed location to improve accessibility. However, researchers, practitioners and participants need to be confident that the data collected, processed and analysed in this way is reliable, valid and trustworthy, and that these aspects are not sacrificed as we strive to harness the opportunities and convenience offered by such technology. Therefore, it is important that there is an evolving evidence base as the technology itself evolves.

The goal of this collection was to share evidence regarding remote data collection and intervention in sport, health and exercise sciences to support academics, practitioners and teachers to make evidence informed decisions about the implementation of such technology.

This editorial summarises three areas of interest that are informed by the papers within this special edition.

## Validity of information integration based on subjective and physiological data from a real sports condition: application to the judgment of fatigue in sport

The measurement and assessment of fatigue in both sporting and health care contexts remains challenging (1, 2) however, researchers, practitioners and athletes are becoming increasingly aware of the impact of fatigue on performance and injury risk (3). If researchers can develop a valid and reliable method to effectively measure this construct remotely this could have real life application for developing training sessions with athletes.

The paper by Legall et al. highlights that current subjective measures of fatigue do not allow for the investigation into the possible cognitive processes that may be involved in how an individual generates and perceives fatigue. This paper attempts to explore this by assessing if exercise duration and intensity are used to create a judgement of fatigue. In this study 20 healthy, experienced participants conducted a laboratory session where they provided subjective self-reported fatigue scores based on six different cycling scenarios (15 or 30 min at 30, 50 or 70% maximum intensity). They then undertook each of the six scenarios in a sports hall to see if the perceived fatigue score correlated with the actual fatigue score, as well as collecting objective measures of fatigue. This study shows that all three measures of fatigue correlated with each other, which provides evidence of a correlation between physiological and psychological measures of fatigue. Legall et al. argue that this could be used as a judgement indicator that may allow for more precise and individualised training programmes, although they highlighted that athletes may underestimate their fatigue levels at higher workloads.

## Quantifying lumbar sagittal plane kinematics using a wrist-worn inertial measurement unit

Measuring human movement is important in many different contexts including health, sport and physical activity. Until recently, accurate measurement of human movement has been confined to well controlled laboratory environments. This limits the environments where data is collected and limits the ecological validity of many measurements. However, recent advancements in remote monitoring have enabled human movement measurement outside the laboratory. Examples of remote monitoring in the context of human movement include markerless motion capture (4) and wearable sensors (5).

Wearable sensors, such as inertial measurement units (IMUs), are often small devices attached to the human

body and often consist of gyroscopes and accelerometers allowing for a range of biomechanical variables to be calculated. However, the reliability and validity of these devices is still being assessed (6), particularly when movement data can only be measured using devices worn on the area of interest.

To understand the transferability of information from lumbar worn sensors to wrist worn sensors, the paper by Liew et al. in this special issue tested the feasibility of using a wrist-worn IMU to infer lumbar sagittal plane kinematics as a substitute for a lumbar-worn IMU. Eighteen healthy participants performed spinal flexion and extension movements while wearing IMUs on the wrist and lumbar spine. The results showed that flexion range of motion (RoM) was the only variable with a statistically significant difference between sensor locations, with a mean difference of 4.54° (95% CI = 1.82°–7.27°). However, the maximal difference across all outcomes was less than 8°, suggesting that wrist-worn IMUs may provide a practical solution for remote lumbar mobility monitoring, potentially leading to more ergonomically acceptable methods of collecting lumbar spine data. Despite this potential, challenges remain. The accuracy of wrist-worn IMUs may be influenced by individual anthropometric differences and movement speeds. Further research is needed to validate this method across different populations and movement conditions. Nonetheless, wrist-worn IMUs may present an alternative approach for remote lumbar mobility assessment, particularly in rehabilitation contexts where self-monitoring is essential.

## Promoting physical activity and a healthy active lifestyle in community-dwelling older adults: a design thinking approach for the development of a mobile health application

Mobile health (mHealth) applications have become an integral part of modern healthcare, leveraging the ubiquity of smartphones and wearable devices to improve health outcomes, and place a growing emphasis on more accessible, personalised healthcare. These applications encompass a broad range of functions, from monitoring chronic conditions and offering mental health support to providing medication reminders and tracking physical activity (7).

The paper from Daniels et al. in this issue focusses on the public health challenge of engaging older adults (>65 years old) in PA and provides a qualitative synthesis of barriers and facilitators to PA in older adults. Outputs from one-to-one interviews (phase 1) directly informed the co-creation of an mHealth app prototype (phase 2) to promote an active lifestyle to community-dwelling older adults. The authors employ co-creation sessions with older adults and topic experts, underpinned by design thinking methodology (8). The co-produced mHealth prototype was then tested in a population of community-dwelling older adults (phase 3). This study highlights the diverse perceptions of PA engagement in this population,

with key considerations for both their own co-created mHealth app and future studies interested in the development of remote PA promotion in older adults.

Challenges persist regarding data privacy and app efficacy (9), along with the need for further evaluation, particularly among those with limited digital access. As mHealth applications continue to develop, it holds the potential to transform how individuals manage their health and interact with healthcare systems globally.

## Summary

Technology in sport and health has accelerated the inclusion of remote data collection and mobile applications. Current research reports on the increased use of technology in this way, the validity and feasibility of these measures, and the development of mobile health apps for promoting physical activity. Despite promising advancements, challenges like data reliability, privacy, and accessibility remain.

## Author contributions

DP: Conceptualization, Writing – original draft, Writing – review & editing. MG: Conceptualization, Writing – original draft, Writing – review & editing. DL: Conceptualization,

Writing – original draft, Writing – review & editing. NB: Writing – original draft, Writing – review & editing. JF: Conceptualization, Writing – original draft, Writing – review & editing. JA: Writing – original draft, Writing – review & editing.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

1. Billones R, Liwang JK, Butler K, Graves L, Saligan LN. Dissecting the fatigue experience: a scoping review of fatigue definitions, dimensions, and measures in non-oncologic medical conditions. *Brain Behav Immunity Health*. (2021) 15:100266. doi: 10.1016/j.bbhih.2021.100266
2. Fitzpatrick JF, Hicks KM, Russell M, Hayes PR. The reliability of potential fatigue-monitoring measures in elite youth soccer players. *J Strength Cond Res*. (2021) 35(12):3448–52. doi: 10.1519/JSC.0000000000003317
3. Díaz-García J, González-Ponce I, Ponce-Bordón JC, López-Gajardo MÁ, Ramírez-Bravo I, Rubio-Morales A, et al. Mental load and fatigue assessment instruments: a systematic review. *Int J Environ Res Public Health*. (2021) 19(1):419. doi: 10.3390/ijerph19010419
4. Lam WW, Tang YM, Fong KN. A systematic review of the applications of markerless motion capture (mmc) technology for clinical measurement in rehabilitation. *J Neuroeng Rehabil*. (2023) 20(1):57. doi: 10.1186/s12984-023-01186-9
5. Picerno P, Iosa M, D'Souza C, Benedetti MG, Paolucci S, Morone G. Wearable inertial sensors for human movement analysis: a five-year update. *Expert Rev Med Devices*. (2021) 18(sup1):79–94. doi: 10.1080/17434440.2021.1988849
6. Ghattas J, Jarvis DN. Validity of inertial measurement units for tracking human motion: a systematic review. *Sports Biomech*. (2024) 23(11):1853–66. doi: 10.1080/14763141.2021.1990383
7. Abaza H, Marschollek M. Mhealth application areas and technology combinations. *Methods Inf Med*. (2017) 56(S 01):e105–e22. doi: 10.3414/ME17-05-0003
8. Roberts JP, Fisher TR, Trowbridge MJ, Bent C. A design thinking framework for healthcare management and innovation. *Healthcare*. (2016) 4(1):11–14. doi: 10.1016/j.hjdsi.2015.12.002
9. Messner E-M, Probst T, O'Rourke T, Stoyanov S, Baumeister H. mHealth applications: potentials, limitations, current quality and future directions. In: Baumeister H, Montag C, editors. *Digital Phenotyping and Mobile Sensing. Studies in Neuroscience, Psychology and Behavioral Economics*. Cham: Springer (2019). doi: 10.1007/978-3-030-31620-4\_15