



Article

“I Don’t Approve of a Fat Person...”: A Cross-Sectional Survey Exploring the Perceptions of Health, Weight and Obesity

Jordan D. Beaumont ^{1,*}, Tina Reimann ^{1,†}, Rosie Wyld ^{1,†} and Beverley O’Hara ²¹ Sheffield Business School, Sheffield Hallam University, Sheffield S1 1WB, UK² School of Health, Leeds Beckett University, Leeds LS1 3HE, UK; b.a.ohara@leedsbeckett.ac.uk

* Correspondence: j.beaumont@shu.ac.uk

† These authors contributed equally to this work.

Abstract

Despite being acknowledged as a complex and multi-faceted condition, the prevailing view within society is that obesity is the result of individual choices and can be reversed simply by “eating less and moving more”. This is oversimplistic and leads to the view that obesity is the individual’s fault and is therefore their responsibility to remedy. These views are grounded in individuals’ beliefs around health and contribute to weight bias and stigma. In the present study, participants (n = 143) completed a cross-sectional survey which explored views around weight and health and whether weight bias or stigma differed based on demographic characteristics, weight status, and prior experience of weight stigma. Results indicate differences in the way individuals living with overweight and obesity are viewed in comparison with those of a healthy weight, with the former viewed in a more negative light. Interestingly, while women presented with higher weight bias scores ($p = 0.036$), men scored higher for externalised weight stigma ($p = 0.001$). Weight status was seen as an important factor contributing to overall health. These results demonstrate that weight bias and stigma are prevalent and highlight the need for further measures to reduce stigmatising views of people living with overweight and obesity.



Academic Editor: Andrew John Hill

Received: 27 April 2025

Revised: 5 June 2025

Accepted: 17 June 2025

Published: 20 June 2025

Citation: Beaumont, J.D.; Reimann, T.; Wyld, R.; O’Hara, B. “I Don’t Approve of a Fat Person...”: A Cross-Sectional Survey Exploring the Perceptions of Health, Weight and Obesity. *Obesities* **2025**, *5*, 49. <https://doi.org/10.3390/obesities5030049>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: weight bias; weight stigma; thin ideal; muscular ideal

1. Introduction

Officially recognised as a major public health challenge and a global epidemic in 1997 [1], the prevalence of obesity has since continued to rise with global estimates of overweight and obesity circa 2.5 billion adults [2]. Despite obesity being acknowledged and well documented as a complex and multi-faceted condition [3,4], the prevailing view within society is that obesity is the result of individual choices and can be reversed through decisions to reduce dietary intake and increase physical activity [5]. This oversimplifies the factors contributing to increased weight status and leads to the view that obesity is the individual’s fault and is therefore their responsibility to remedy [6]. These assumptions impact societal views of individuals living with obesity, who are widely perceived as lacking willpower, being lazy, greedy or selfish, and draining healthcare resources [7–9].

These views are grounded in individuals’ beliefs about the impact of individual behaviours on health [10]. Health is a relative state which is often perceived differently between individuals depending on their personal associations with and expectations of what contributes towards being healthy [11]. Societal perception of health status is often closely linked to physical appearance and weight status [12]. While there are links with higher weight

status and disease prevalence, Ortega et al. [13] argue that weight should not be used as the sole criterion for establishing health status and that, instead, cardiorespiratory fitness is a more valid metric. Still, despite not being included in any definition of health [14–16], being ‘healthy’ is associated with adherence to the thin and muscular ideals [17,18]. This focus on physical appearance has given rise to body dissatisfaction and body image disorders [19,20], and individuals who do not adhere with societal views of health (i.e., the thin and muscular ideals) often experience weight-based bias and stigmatisation [21,22].

Weight bias refers to the negative attitudes, beliefs, assumptions and judgements held about those living with obesity and can take the form of explicit (i.e., overtly negative attitudes towards people in larger bodies), implicit (i.e., unconscious negative attitudes towards people in larger bodies), and internalised bias (i.e., extent to which negative beliefs are endorsed by individuals) [5]. Weight stigma is the manifestation of weight bias through harmful social stereotypes associated with the ‘ideal’ body shape and/or size. The stigmatisation of individuals living with obesity is pervasive in current society [5,23]. Indeed, individuals living with obesity are nearly nine times more likely to suffer from weight-based discrimination compared with those who are classed as a healthy weight [24].

Experiences of weight bias and stigma are prevalent across the lifecycle and in a range of settings, including within the home, in social settings, in the workplace, and when accessing healthcare [23,25–27]. In a recent survey by Puhl et al. [27], 58% of participants reported experiencing weight stigma at some point during their life, and more than half reported being teased about their weight. Internalisation of weight stigma, where societal negative stereotypes are applied to oneself contributing to mistreatment or devaluation of self [28], has a mediating role on an individuals’ body image, health-related behaviour (e.g., physical activity), and engagement with and experience of healthcare [29].

It has been proposed that weight bias and stigma are routes through which to motivate weight loss [30]; i.e., if it were suitably distasteful to live with obesity, then individuals would be motivated to alter their behaviours and lose weight [31]. However, there is no empirical evidence to support this, and instead, this appears to create the opposite reaction [32]. Experiencing weight-based stigma can negatively impact body image and mental health and wellbeing, with internalised weight bias associated with weight gain, weight cycling, perceived stress, and eating to cope [33–36]. Interestingly, experiencing stigma has been shown to predict weight gain, regardless of age, baseline weight status, race, ethnicity, and socioeconomic factors [5,27,35,37]. What is particularly important to note is that stigmatising views and assumptions around weight and obesity mislead public health policies, confuse messages in popular media, undermine access to evidence-based treatments, and compromise advances in research [5].

Exploring views and perceptions of weight and health across populations is important for identifying those who are more likely to express stigmatising views in order to develop targeted approaches to reduce stigmatising views and bias. While prior work has established clear links between stigmatising views and a wide range of impacts on the individual, this is often through validated questionnaires or qualitative methodology. Given that weight-based stigma is often fuelled by visual cues (e.g., viewing an individual living with obesity, comparing them to the thin ideal, depictions in news articles) [5,38,39], it is important to additionally understand whether views based on such visual cues differ from those captured through validated measures which require more abstract thinking. This project looked to further explore the perceptions of and links between health and weight based on both questionnaire- and image-based measures and assess whether the perception of weight, and particularly weight bias or stigma, differed based on demographic characteristics, weight status, and prior experience of weight bias and/or stigma. Specifically, this project looked to address the following questions: (i) How do individuals

perceive ‘health’, and what factors contribute to being ‘healthy’? (ii) Are there specific demographic characteristics associated with weight-based stigmatising views? (iii) Does prior experience of weight bias/stigma affect internalised/externalised stigmatising views?

2. Materials and Methods

2.1. Study Design

This study involved a cross-sectional online survey completed through Qualtrics. Participants were self-identified in response to study advertisements. The study was shared via social media and word-of-mouth, with interested individuals directed to the online survey where they could review a participant information sheet. All individuals provided informed consent prior to their participation. Procedures were independently reviewed and approved by the Sheffield Hallam University Research Ethics Committee (project ID: ER42086514, approved 12 April 2022). Given the sensitive nature of the survey, participants were directed to self-help resources should they be worried about their weight or wider health and wellbeing.

This article is an expanded version of a paper entitled ‘Exploring the perceptions of health, weight, and obesity’, which was presented at the UK Congress on Obesity at Queen’s University Belfast on 14 September 2023 [40].

2.2. Participants

The survey was open to adults (18 years of age or older) who were free-living (i.e., non-clinical populations). The study aimed to recruit a minimum target sample size of 133 individuals, in line with other similar studies [41–43].

2.3. Materials

2.3.1. Participant Demographics

Participants self-reported their age, gender, ethnicity, highest level of attained education, current student status, and perceived weight status (underweight, a healthy weight, overweight, or obese). Body image was determined using the scale produced by Stunkard et al. [44]; participants selected a silhouette drawing ranging from one (leanest silhouette) to nine (largest silhouette) that best matched how they viewed their body size. The level of physical activity was measured using the International Physical Activity Questionnaire (IPAQ) [45], capturing levels of activity over the previous seven days and providing an indication of the habitual level of activity, and categorised as low, moderate, or high.

2.3.2. Weight Bias

Weight bias was measured using the Attitudes Towards Obese Persons Scale (ATOPS) and the Beliefs About Obese Persons Scale (BAOPS) [46]. The ATOPS is a 20-item questionnaire measuring attitudes and perceptions about individuals living with obesity. Participants respond to statements over a six-point Likert scale, with scores ranging from −3 (I strongly disagree) to +3 (I strongly agree). Scores are totalled and summed with 60, with a total score ranging from 0 to 120; scores below 60 indicate more negative attitudes towards individuals living with obesity, whereas scores greater than 60 correspond with more positive attitudes towards individuals living with obesity. Similarly, the BAOPS is an eight-item scale assessing beliefs about the causes of obesity and is completed over the same six-point Likert scale. Item scores are totalled and summed with 24 to produce a total score. Scores less than 24 highlight the belief that obesity is under the individual’s control, whereas scores greater than 24 highlight the belief that obesity is not controlled by the individual. Both the ATOPS (Cronbach’s $\alpha = 0.80$ to 0.84) and BAOPS (Cronbach’s $\alpha = 0.65$ to 0.82) have good internal consistency [46].

2.3.3. Weight Stigma

Internalised weight stigma was measured using the 12-item Weight Self-Stigma Questionnaire (WSSQ) [47]. This questionnaire views weight stigma as a construct with two distinct factors—self-devaluation and fear of enacted stigma. Participants respond to statements over a five-point Likert scale (completely disagree to completely agree), and responses are summed, ranging from 12 to 60 for the total scale and 6 to 30 for individual factors. Higher scores indicate that participants experience greater shame related to their weight or body shape. The WSSQ has good internal consistency for the total scale (Cronbach's $\alpha = 0.88$) and for the self-devaluation (Cronbach's $\alpha = 0.81$) and fear of enacted stigma (Cronbach's $\alpha = 0.87$) sub-scales [47].

The Universal Measures of Bias-Fat (UMB-FAT) [48] was used to measure externalised weight stigma (Cronbach's $\alpha = 0.87$). Participants respond to 20 items over a seven-point Likert scale, ranging from strongly agree through strongly disagree. Scores are totalled and provide individual item scores for Adverse Judgement, Social Distance, Attraction, and Equal Rights. Individual item scores range from 5 to 35, with higher scores indicating greater stigmatising views of those living with obesity.

Experience of weight-based stigma was measured using the Stigmatising Situations Inventory-Brief (SSI-B) [34], a 10-item questionnaire where participants report the frequency (from never to daily) with which they experience stigma relating to their weight from various sources (e.g., family members, healthcare professionals, members of the public). Scores are totalled and range from 0 to 90; higher scores demonstrate a greater experience of weight-based stigma. The SSI-B has high internal consistency (Cronbach's $\alpha = 0.95$) [34].

2.3.4. Perceptions of Health and Weight

Participants were shown eight images depicting 'healthy' /non-stigmatising (eating a balanced meal, taking part in physical activity) and 'unhealthy' /stigmatising (eating junk food, sedentarism) behaviours conducted by individuals of a healthy weight and individuals living with obesity. The content of each image was matched across weight status groups (e.g., showing the same physical activity in the same environment). Non-stigmatising images of individuals living with obesity were selected from Obesity Canada's image bank; other images were found via an internet image search. The participants were shown each image in a random order and asked to describe what the image depicted. Finally, the participants were asked to list the individual behaviours, characteristics, traits, or other factors that they believe contribute to good and poor health as well as the societal or environmental factors that they believe contribute to good and poor health; the participants were also asked to describe an individual they would consider to be in good and poor health.

2.4. Data Analysis

Questionnaires were scored in line with instructions provided in the validation papers (as described above). Normality of data was determined using a Shapiro–Wilk test. For normally distributed data, means were compared using independent sample t-tests or one-way analysis of variance (ANOVA) as appropriate for the comparison. Where data were not normally distributed, the Mann–Whitney U test and the Kruskal–Wallis test were used. Spearman's correlation coefficient was used to determine the correlation between measures of weight bias and stigma. As the WSSQ is only validated for use in individuals with overweight and obesity, analyses were restricted to those participants self-reporting as being overweight or obese. Analyses were completed using the Statistical Package for the Social Sciences (SPSS) version 29 (IBM, New York, NY, USA). Data are presented as mean and standard deviation (SD) or as median and interquartile range (IQR), as appropriate to an alpha level of 0.05.

Qualitative data on factors contributing to good and poor health were analysed through inductive thematic analysis [49] by two authors independently (J.D.B. and B.O.). Initial codes were developed and used to inform themes, which were then defined and agreed upon. The themes are presented below with illustrative quotes. Qualitative data in response to images were analysed through sentiment analysis and coded as ‘positive’, ‘neutral’, or ‘negative’ by two authors (J.D.B. and B.O.). Prior to coding, authors agreed upon the criteria for coding, which differed based on image type (i.e., whether the picture depicted eating or exercise). To improve the clarity of reporting, where appropriate, quantitative and qualitative data will be discussed concurrently. Data and associated study files are available via the Open Science Framework: <https://doi.org/10.17605/OSF.IO/KZ4VJ> (accessed on 18 June 2025).

3. Results

A total of 143 participants completed the survey. The participants were between 21 and 70 years of age (35.1 ± 12.0 years; $n = 105$) and were predominantly white, women, non-students, and well educated, with 79% being educated to an undergraduate degree level or higher (Table 1). Of the participants, 76.0% ($n = 109$) self-reported having a healthy weight, and 22.4% ($n = 32$) self-reported living with overweight or obesity. Body image scores ranged from 1 to 7 (median 4.0 arbitrary units [AU], IQR 2.0 AU). Only 16.1% ($n = 23$) of the participants reported that they did not use any method to help manage their weight (Table 2). There were no differences across characteristics when comparing by gender ($p \geq 0.213$) or level of education ($p \geq 0.146$). Physical activity level differed when comparing across ethnicity, with those of Asian or Asian British ethnicity having lower levels of activity ($\chi^2_{(3)} = 20.082$, $p < 0.001$). Similarly, those with higher body image scores reported lower levels of physical activity ($\chi^2_{(2)} = 13.938$, $p < 0.001$). Both higher age ($\chi^2_{(2)} = 6.899$, $p = 0.032$) and body image scores ($\chi^2_{(3)} = 56.060$, $p < 0.001$) were associated with higher weight status, whereas individuals self-identifying as living with overweight or obesity had lower levels of physical activity compared with those with a healthy weight ($\chi^2_{(3)} = 8.576$, $p = 0.035$). Interestingly, a perceived struggle to maintain a healthy weight was associated with lower weight status ($U = 1224.500$, $p < 0.001$), lower body image scores ($U = 1153.50$, $p < 0.001$), and lower age ($U = 859.00$, $p = 0.010$).

Table 1. Participant characteristics ($n = 143$).

		n	%
Gender	Women (including transgender women)	104	72.7
	Men (including transgender men)	38	26.6
	Prefer not to state	1	0.7
Ethnicity	White	121	84.6
	Asian or Asian British	13	9.1
	Mixed or multiple ethnic group	5	3.5
	Black, African, Caribbean, or Black British	4	2.8
Education	Undergraduate degree	64	44.8
	Postgraduate taught degree	36	25.2
	Further education (e.g., A-level)	25	17.5
	Doctorate or other postgraduate research degree	13	9.2
	Secondary education (e.g., GCSE)	4	2.8
	Prefer not to state	1	0.7

Table 1. *Cont.*

		n	%
Student status	No	101	70.6
	Yes	42	29.4
Perceived weight status	Healthy weight	109	76.2
	Overweight	31	21.7
	Underweight	2	1.4
	Obese	1	0.7
Struggle to maintain a healthy weight	No	93	65.0
	Yes	50	35.0
Physical activity level	High	123	86.0
	Moderate	16	11.2
	Low	4	2.8

Table 2. Methods used to support weight management (n = 143).

	n	%
Weight-loss diet	15	10.5
Skip meals	28	19.6
Avoid certain foods/food groups	47	32.9
Avoid certain eating practices (e.g., snacking)	42	29.4
Meal replacement products (e.g., shakes, bars)	7	4.9
Meal planning	1	0.7
Calorie tracking apps	25	17.5
Weight-loss clubs or groups	5	3.5
Exercise/Physical activity	99	69.2
Slimming or diet products (e.g., pills, medications)	2	1.4
I do not use any method to manage my weight	23	16.1

3.1. Weight Bias

All participants reported more negative attitudes towards individuals living with obesity (39.3 ± 9.0 AU). While there appeared to be no difference in views based on age ($r_{(106)} = -0.014$, $p = 0.885$), women (38.4 ± 9.1 AU) held more negative attitudes than men (41.9 ± 8.3 AU) ($t_{(141)} = 2.114$, $p = 0.036$). Median ATOPS scores did not differ based on perceived weight status ($\chi^2_{(3)} = 4.868$, $p = 0.182$), ethnicity ($\chi^2_{(3)} = 2.223$, $p = 0.527$), level of education ($\chi^2_{(6)} = 4.560$, $p = 0.601$), physical activity level ($\chi^2_{(2)} = 1.275$, $p = 0.529$), or use of weight management techniques ($z = -0.388$, $p = 0.698$).

ATOPS scores were positively correlated with BAOPS scores ($r_{s(144)} = 0.282$, $p < 0.001$) (Table 3). This suggests that while views were negative, participants still appreciated that obesity was beyond an individual's control. In line with this, median BAOPS scores were 26.0 AU (IQR 6.0 AU). There was no difference in BAOPS scores when comparing age ($r_{s(106)} = 0.014$, $p = 0.883$), gender (women: median 26.0 AU, IQR 7.0 AU; men: median 26.0 AU, IQR 5.0 AU; $z = -0.476$, $p = 0.634$), ethnicity ($\chi^2_{(3)} = 1.092$, $p = 0.779$), perceived weight status (healthy weight: median 26.0 AU, IQR 6.0 AU; overweight or obesity: median 26.0 AU, IQR 6.0 AU; $z = -0.31$, $p = 0.756$), level of education ($\chi^2_{(6)} = 5.279$, $p = 0.509$), physical activity level ($\chi^2_{(2)} = 5.850$, $p = 0.054$), or use of weight management techniques (techniques used: median 26.0 AU, IQR 6.0 AU; techniques not used: median 26.0 AU, IQR 6.0 AU; $z = -0.745$, $p = 0.456$).

Table 3. Spearman’s correlation coefficient for measures of weight bias and stigma (n = 143).

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.
1. ATOPS		0.282 ‡	0.021	0.265 ‡	0.340 ‡	0.146	0.230 †	0.246 †	−0.094	−0.112	−0.063
2. BAOPS	0.282 ‡		0.048	0.352 ‡	0.264 ‡	0.171 *	0.439 ‡	0.185 *	−0.085	−0.073	−0.099
3. SSI-B	0.021	0.048		−0.007	−0.009	−0.083	−0.078	−0.074	−0.360 ‡	−0.324 ‡	−0.310 ‡
4. UMB-FAT (total)	0.265 ‡	0.352 ‡	−0.007		0.550 ‡	0.657 ‡	0.744 ‡	0.824 ‡	0.083	0.060	0.088
5. UMB-FAT (adverse judgement)	0.340 ‡	0.264 ‡	−0.009	0.550 ‡		0.325 ‡	0.489 ‡	0.516 ‡	0.026	−0.006	0.041
6. UMB-FAT (social distance)	0.146	0.171 *	−0.083	0.657 ‡	0.325 ‡		0.318 ‡	0.498 ‡	0.094	0.059	0.149
7. UMB-FAT (attraction)	0.230 †	0.439 ‡	−0.078	0.744 ‡	0.489 ‡	0.318 ‡		0.387 ‡	0.029	0.056	−0.026
8. UMB-FAT (equal rights)	0.246 †	0.185 *	0.074	0.824 ‡	0.516 ‡	0.498 ‡	0.387 ‡		0.019	−0.029	0.068
9. WSSQ (total)	−0.094	−0.085	−0.360 ‡	0.083	0.026	0.094	0.029	0.019		0.876 ‡	0.856 ‡
10. WSSQ (self-devaluation)	−0.112	−0.073	−0.324 ‡	0.060	−0.006	0.059	0.056	−0.029	0.876 ‡		0.567 ‡
11. WSSQ (fear of enacted stigma)	−0.063	−0.099	−0.310 ‡	0.088	0.041	0.149	−0.026	0.068	0.856 ‡	0.567 ‡	

* $p \leq 0.05$, † $p \leq 0.01$, ‡ $p \leq 0.001$.

3.2. Weight Stigma

Individuals who self-reported as being overweight or obese experienced greater stigma (4.0 AU, IQR 9.0 AU) than healthy or underweight participants (1.0 AU, IQR 6.0 AU) ($U = 1124.0, p < 0.001$). Externalised stigma was more prevalent in men (3.8 AU, IQR 0.5 AU) than women (3.6 AU, IQR 0.3 AU) ($t = 3.32, p = 0.001$). Specifically, men enacted more negative judgement of individuals living with obesity ($z = -2.90, p = 0.004$) and viewed individuals living with obesity as less attractive (5.2 AU, IQR 2.0 AU), as compared with women (4.1 AU, IQR 2.0 AU) ($z = -2.90, p = 0.004$). Support for equal rights for people living with obesity was also greater among men (median 2.2 AU, IQR 2.8 AU) than women (1.6 AU, IQR 1.4 AU) ($z = -2.69, p = 0.007$). There was no difference in reported internalised stigma (men: 36 AU, 7 AU; women: 36 AU, 9 AU; $z = -0.23, p = 0.824$) or experience of weight stigma between men (11.0 AU, IQR 6.0 AU) and women (12.0 AU, IQR 8.0 AU) ($U = 1891.5, p = 0.523$).

3.3. Perceptions of Health and Weight

When asked to describe an individual of good health, participants commented primarily on the need for physical activity ($n = 67, 46.9\%$), a healthy diet ($n = 54, 37.8\%$), low weight status ($n = 28, 19.6\%$), and good mental health ($n = 26, 18.2\%$). In comparison, descriptions of an individual of poor health focussed on poor diet ($n = 55, 38.5\%$), lack of physical activity ($n = 51, 35.7\%$), high weight status ($n = 39, 27.3\%$), and personal attributes ($n = 29, 20.3\%$). This is reflected in the individual, societal, and environmental factors participants associated with good and poor health (Table 4).

Table 4. Codes identified during thematic analysis of responses outlining the individual, societal, and environmental factors associated with good and poor health (n [%]).

Factors Contributing to Good Health				Factors Contributing to Poor Health			
Individual		Societal and Environmental		Individual		Societal and Environmental	
Physical activity	78 (54.5)	Social health	51 (35.7)	Diet	58 (40.6)	Resource	55 (38.5)
Diet	73 (51.0)	Environment	44 (30.8)	Personal attributes	54 (37.8)	Environment	50 (35.0)
Personal attributes	60 (42.0)	Resource	43 (30.1)	Physical activity	47 (32.9)	Social health	33 (23.1)
Mental health	20 (14.0)	Education	26 (18.2)	Mental health	41 (28.7)	Education	19 (13.3)
Education	17 (11.9)	Physical activity	20 (14.0)	Resource	17 (11.9)	Mental health	16 (11.2)
Sleep	14 (9.8)	Personal attributes	16 (11.2)	Substances	16 (11.2)	Employment	13 (9.1)
Social health	14 (9.8)	Diet	13 (9.1)	Education	14 (9.8)	Personal attributes	13 (9.1)
Environment	7 (4.9)	Mental health	10 (7.0)	Social health	14 (9.8)	Diet	9 (6.3)
Resource	7 (4.9)	Employment	7 (4.9)	Sleep	10 (7.0)	Marketing	9 (6.3)
Genetics	5 (3.5)	Healthcare	6 (4.2)	Upbringing	9 (6.3)	Healthcare	7 (4.9)
Substances	4 (2.8)	Marketing	6 (4.2)	Environment	7 (4.9)	Upbringing	7 (4.9)
Physical health	3 (2.1)	Upbringing	6 (4.2)	Genetics	7 (4.9)	Work/life balance	6 (4.2)
Work/life balance	3 (2.1)	Work/life balance	5 (3.5)	Physical health	6 (4.2)	Physical activity	5 (3.5)
Hobbies	2 (1.4)	Lifestyle	3 (2.1)	Work/life balance	4 (2.8)	Lifestyle	4 (2.8)
Lifestyle	2 (1.4)	Genetics	2 (1.4)	Lifestyle	3 (2.1)	Social media	3 (2.1)
Economy	1 (0.7)	Government	2 (1.4)	Social media	2 (1.4)	Substances	2 (1.4)
Employment	1 (0.7)	Social media	2 (1.4)	Marketing	2 (1.4)	Demographics	1 (0.7)
Support	1 (0.7)	Society	2 (1.4)	Economy	1 (0.7)	Physical health	1 (0.7)
Upbringing	1 (0.7)	Psychology	1 (0.7)	Personal attributes	1 (0.7)	Society	1 (0.7)
		Substances	1 (0.7)				

Qualitative comments in response to images were consistently more negative if an image depicted an individual living with obesity versus a healthy weight, regardless of the

activity type or whether the image was stigmatising or non-stigmatising (Table 5). When exploring the comments, there is a clear distinction in participants' focus. For images depicting activities involving food/eating, those images that included individuals with a healthy weight often focussed on the activity and setting (e.g., *"This is a group of friends enjoying some pizza and socialising with each other. They all look happy and like they're enjoying themselves."*; *"Friends out for dinner having fun"*). In comparison, where these images included an individual living with obesity, comments focussed primarily on the individuals' physical appearance (e.g., *"First thought was 'ew' looks sort of slobbish and unappealing"*; *"Sedentary and overweight"*; *"I'm very sad that a young guy in his prime of life has let himself get like that"*) or assumed individual behaviours, characteristics, or background (e.g., *"Lazy, poorly motivated, lacking in healthy role models"*; *"Out of control, lazy, negative feeling"*; *"Lazy inactive person, not aspiring for good things"*).

Table 5. Positive, neutral, and negative comments (n [%]) in response to images depicting stigmatising and non-stigmatising activities of individuals with a healthy weight or with obesity (n = 143).

Weight Status	Activity	Positive	Neutral	Negative	
Obesity	Consuming unhealthy food (stigmatising)	2 (1.4)	70 (49.0)	71 (49.7)	$z = -8.618$, $p < 0.001$
Healthy	Consuming unhealthy food (stigmatising)	69 (48.3)	64 (44.8)	10 (7.0)	
Obesity	Consuming balanced meal (non-stigmatising)	52 (36.4)	60 (42.0)	31 (21.7)	$z = -7.119$, $p < 0.001$
Healthy	Consuming balanced meal (non-stigmatising)	89 (62.2)	52 (36.4)	2 (1.4)	
Obesity	Being sedentary (stigmatising)	3 (2.1)	40 (28.0)	100 (69.9)	$z = -5.851$, $p < 0.001$
Healthy	Being sedentary (stigmatising)	5 (3.5)	105 (73.4)	33 (23.1)	
Obesity	Being physically active (non-stigmatising)	39 (27.8)	87 (60.8)	17 (11.9)	$z = -5.601$, $p < 0.001$
Healthy	Being physically active (non-stigmatising)	88 (61.5)	54 (37.8)	1 (0.7)	

Similarly, descriptions of images depicting sedentary behaviour versus physical activity were generally negative and judgemental (e.g., *"Overweight person"*; *"I can't tell where the person starts and the sofa ends. Clearly the bloke is eating more than he needs. . ."*), focussing on appearance (e.g., *"Looks sort of slobbish and unappealing"*), or their personal attributes (e.g., *"Lazy poorly motivated lacking in healthy role models"*; *"Unhappy, unhealthy, lonely"*). In comparison, images of those with a healthy weight focussed more on rest and relaxation for sedentary behaviours (e.g., *"Looks like he's earned a nice nap, although should probably have gone to bed a little earlier"*; *"He looks exhausted and tired out, maybe he is having a relaxed day or has just completed a task. He looks quite relaxed and that he is happy"*) or health/aspirational goals for physical activity (e.g., *"Looks healthy, actively trying to better herself"*; *"A woman exercising and trying to live a healthy lifestyle—she may be doing this to lose weight or get in better shape, but if she isn't, that's fine"*).

4. Discussion

This work looked to explore the perceptions held by participants around health, weight, and obesity, particularly where weight-based bias and stigma were prevalent. The findings suggest that there are differences in the way individuals living with overweight and obesity are viewed in comparison with those of a healthy weight, with the former viewed in a more negative light. Weight status was seen as an important factor contributing

to both good health (i.e., maintaining a 'healthy' weight status) and poor health (i.e., living with overweight or obesity).

While weight may be associated with health, health is complex, and a 'one size fits all' approach cannot be applied to the concept of 'being healthy' [50,51]. Historically, being healthy was considered as being free of illness or disease [14]. Today health is seen more as a holistic concept, incorporating physical, mental, and social wellbeing and not just the absence of disease or infirmity, with some definitions associating health primarily with good fitness [52]. Our findings support this connection, with comments in response to those of a healthy weight focussing on physical, mental, and social wellbeing. However, this was not observed for those living with obesity, who were instead described with more stigmatising language. Individuals with overweight or obesity are at increased risk of experiencing weight-based stigmatisation compared with their healthy-weight counterparts [24,53].

The comments made by participants in the present work align with those observed in other studies, that is, perceiving individuals of higher weight status as 'lazy' or 'weak-willed' and making assumptions about behaviours and lifestyle (e.g., lack of participation in physical activity) [8,54,55]. The overwhelmingly negative views held about those living with obesity in the present sample may be surprising given their higher educational status. Education is generally associated with greater tolerance and decreased prejudice [56]. However, more recent research has shown that where views are more ideological (e.g., aligned with political beliefs), this tolerance flips; people with higher educational status exhibit greater intolerance and higher prejudice [57]. This could suggest that beliefs about those living with obesity may be linked to ideology. Replication of the present work in a less educated sample may provide further information on this.

Weight stigma is considered the last acceptable form of discrimination [58–60]. The prevalence of stigmatising views around weight has given rise to approaches such as Health at Every Size® [61] and the Fat but Fit Paradox [13]. These acknowledge that weight status is not an appropriate proxy for health status, with disease being prevalent across weight categories. Indeed, weight status should not be used as the sole determinant of health, as it provides no conclusive indication of health status given that individuals can be metabolically unhealthy at a healthy weight and metabolically healthy at a higher weight [62–64]. An individual's metabolic health is associated with disease development and progression [64–66]. While the risk for disease development cannot exclude body weight and adipose distribution completely, it is important to consider other factors such as the social and commercial determinants of health [67–69]. Recognising this will help address weight-based stereotypes, prejudice, discrimination, and unfair treatment experienced by individuals living with obesity across multiple facets of everyday life [36,70] and as observed in the present work.

The present work observed significantly higher weight bias in women, compared with men. Given the increased societal pressures placed on appearance in women [71,72] and the need to adhere with the 'thin ideal' [17,18], it is unsurprising that such a bias exists, and it is in fact observed in the present sample. In addition, women are more vulnerable to weight stigma [73–75], as are those with higher weight status [24,53,76]. Individuals who more frequently experience weight stigma are more likely to internalise such stigma, endorsing the negative stereotypes and attributing them to themselves [29,77]. This internalisation, however, was not observed in the present sample, as there were no significant differences in the internalised weight stigma scores. Interestingly, externalised weight stigma was higher in men than women. This suggests that while women appear to hold more negative views, men are more likely to enact harmful social stereotypes associated with weight.

Both diet and physical activity were deemed as important contributors to health—contributing to both good and poor health. Dryer and Ware [78] observed 'eating more food

than you need', 'eating more food than you need', and 'not getting enough physical activity' as factors contributing to weight gain. This, again, demonstrates the parallels drawn between weight and health. The experience of weight-based stigma is associated with weight gain, weight cycling, perceived stress, and eating to cope [33–36]. Importantly, a higher incidence of weight stigma is a precursor of psychological disorders (e.g., disordered eating, anxiety, depression, suicidal ideation) [33,79,80]. Stigma is experienced throughout a range of societal settings, for example, healthcare [81], education [21], employment [82], and the media [38], contributing to significant psychological harm, discrimination, and social inequalities.

While this study has identified a number of interesting findings in relation to weight bias and stigmatising views and has captured these through validated questionnaires and qualitative questions, the study is not without limitations. While the study has captured views around health and weight, it was outside the aim of this project to capture the factors contributing to these views. In addition, the cross-sectional nature of the project did not allow for an in-depth exploration of these views beyond simple qualitative questions. Such a detailed exploration would have provided an opportunity to explore nuances in perceptions and understanding of weight and health. Finally, the sample size was modest in comparison with other published work. However, a posteriori power calculations suggest that while some analyses have low achieved power, many demonstrate satisfactory achieved power (>0.816 , e.g., UMB-FAT total, UMB-FAT attraction). The sample was not representative, and therefore, the views expressed here are likely limited to a white, female, well-educated, and healthy-weight demographic.

5. Conclusions

This study identified prevalent stigmatising view of those living with overweight and obesity across validated measures and in response to images depicting individuals with a healthy weight or with obesity. These findings agree with the prior literature and demonstrate that weight bias and stigma are prevalent in a sample of highly educated adults. These findings highlight the need for further measures to reduce stigmatising views.

Author Contributions: Conceptualization, J.D.B., T.R., and R.W.; methodology, J.D.B., T.R., and R.W.; validation, J.D.B. and B.O.; formal analysis, J.D.B. and B.O.; investigation, J.D.B., T.R., and R.W.; resources, J.D.B.; data curation, J.D.B.; writing—original draft preparation, J.D.B. and B.O.; writing—review and editing, J.D.B., T.R., R.W., and B.O.; visualization, J.D.B.; supervision, J.D.B.; project administration, J.D.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Sheffield Hallam University Research Ethics Committee (project ID: ER42086514, approved 12 April 2022).

Informed Consent Statement: Informed consent was obtained from all participants involved in the study.

Data Availability Statement: Data and associated study files are available via the Open Science Framework: <https://doi.org/10.17605/OSF.IO/KZ4VJ>, accessed on 18 June 2025.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

ANOVA	Analysis of variance
ATOPS	Attitudes Towards Obese Persons Scale
AU	Arbitrary units
BAOPS	Beliefs About Obese Persons Scale
IPAQ	International Physical Activity Questionnaire
IQR	Interquartile range
SD	Standard deviation
SPSS	Statistical Package for the Social Sciences
SSI-B	Stigmatising Situations Inventory-Brief
UMB-FAT	Universal Measures of Bias-Fat
WSSQ	Weight Self-Stigma Questionnaire

References

1. World Health Organisation. *Obesity: Preventing and Managing the Global Epidemic*; World Health Organisation: Geneva, Switzerland, 2000.
2. World Health Organisation. Obesity and Overweight. Available online: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed on 27 April 2025).
3. Hruby, A.; Hu, F.B. The Epidemiology of Obesity: A Big Picture. *PharmacoEconomics* **2015**, *33*, 673–689. [CrossRef] [PubMed]
4. Vandenbroeck, P.; Goossens, J.; Clemens, M. Tackling Obesities: Future Choices—Building the Obesity System Map. *Foresight* **2007**, *80*, 101037.
5. Rubino, F.; Puhl, R.M.; Cummings, D.E.; Eckel, R.H.; Ryan, D.H.; Mechanick, J.I.; Nadglowski, J.; Ramos Salas, X.; Schauer, P.R.; Twenefour, D.; et al. Joint international consensus statement for ending stigma of obesity. *Nat. Med.* **2020**, *26*, 485–497. [CrossRef] [PubMed]
6. Luck-Sikorski, C.; Riedel-Heller, S.G.; Phelan, J.C. Changing attitudes towards obesity—Results from a survey experiment. *BMC Public Health* **2017**, *17*, 373. [CrossRef]
7. Mankar, M.; Joshi, R.S.; Belsare, P.V.; Jog, M.M.; Watve, M.G. Obesity as a Perceived Social Signal. *PLoS ONE* **2008**, *3*, e3187. [CrossRef]
8. Sikorski, C.; Riedel, C.; Lupp, M.; Schulze, B.; Werner, P.; König, H.-H.; Riedel-Heller, S.G. Perception of overweight and obesity from different angles: A qualitative study. *Scand. J. Public Health* **2012**, *40*, 271–277. [CrossRef]
9. Spahlholz, J.; Baer, N.; König, H.-H.; Riedel-Heller, S.G.; Luck-Sikorski, C. Obesity and discrimination—a systematic review and meta-analysis of observational studies. *Obes. Rev.* **2016**, *17*, 43–55. [CrossRef]
10. Elwell-Sutton, T.; Marshall, L.; Bibby, J.; Volmert, A. *Reframing the Conversation on the Social Determinants of Health*; The Health Foundation: London, UK, 2019.
11. Maniscalco, L.; Miceli, S.; Bono, F.; Matranga, D. Self-Perceived Health, Objective Health, and Quality of Life among People Aged 50 and Over: Interrelationship among Health Indicators in Italy, Spain, and Greece. *Int. J. Environ. Res. Public Health* **2020**, *17*, 2414. [CrossRef]
12. Idema, C.L.; Roth, S.E.; Upchurch, D.M. Weight perception and perceived attractiveness associated with self-rated health in young adults. *Prev. Med.* **2019**, *120*, 34–41. [CrossRef]
13. Ortega, F.B.; Ruiz, J.R.; Labayen, I.; Lavie, C.J.; Blair, S.N. The Fat but Fit paradox: What we know and don't know about it. *Br. J. Sports Med.* **2018**, *52*, 151–153. [CrossRef]
14. Brüssow, H. What is health? *Microb. Biotechnol.* **2013**, *6*, 341–348. [CrossRef] [PubMed]
15. Ereshefsky, M. Defining 'health' and 'disease'. *Stud. Hist. Philos. Sci. Part C Stud. Hist. Philos. Biol. Biomed. Sci.* **2009**, *40*, 221–227. [CrossRef] [PubMed]
16. Huber, M.; Knottnerus, J.A.; Green, L.; Horst, H.v.d.; Jadad, A.R.; Kromhout, D.; Leonard, B.; Lorig, K.; Loureiro, M.I.; Meer, J.W.M.v.d.; et al. How should we define health? *BMJ* **2011**, *343*, d4163. [CrossRef]
17. Jackson, A.M.; Busig, J.; Lanigan, J. "Thin and muscular": A cross-sectional mixed methods study of people's descriptions of health. *Stigma Health* **2022**, *7*, 389–395. [CrossRef]
18. Stephen, I.D.; Perera, A.T.-M. Judging the Difference between Attractiveness and Health: Does Exposure to Model Images Influence the Judgments Made by Men and Women? *PLoS ONE* **2014**, *9*, e86302. [CrossRef] [PubMed]

19. Jiotsa, B.; Naccache, B.; Duval, M.; Rocher, B.; Grall-Bronnec, M. Social Media Use and Body Image Disorders: Association between Frequency of Comparing One's Own Physical Appearance to That of People Being Followed on Social Media and Body Dissatisfaction and Drive for Thinness. *Int. J. Environ. Res. Public Health* **2021**, *18*, 2880. [CrossRef]
20. Merino, M.; Tornero-Aguilera, J.F.; Rubio-Zarapuz, A.; Villanueva-Tobaldo, C.V.; Martín-Rodríguez, A.; Clemente-Suárez, V.J. Body Perceptions and Psychological Well-Being: A Review of the Impact of Social Media and Physical Measurements on Self-Esteem and Mental Health with a Focus on Body Image Satisfaction and Its Relationship with Cultural and Gender Factors. *Healthcare* **2024**, *12*, 1396. [CrossRef]
21. Nutter, S.; Ireland, A.; Alberga, A.S.; Brun, I.; Lefebvre, D.; Hayden, K.A.; Russell-Mayhew, S. Weight Bias in Educational Settings: A Systematic Review. *Curr. Obes. Rep.* **2019**, *8*, 185–200. [CrossRef]
22. Nutter, S.; Russell-Mayhew, S.; Saunders, J.F. Towards a sociocultural model of weight stigma. *Eat. Weight. Disord.—Stud. Anorex. Bulim. Obes.* **2021**, *26*, 999–1005. [CrossRef]
23. Brown, A.; Flint, S.W.; Batterham, R.L. Pervasiveness, impact and implications of weight stigma. *EClinicalMedicine* **2022**, *47*, 101408. [CrossRef]
24. Hill, B.; Bergmeier, H.; Incollingo Rodriguez, A.C.; Barlow, F.K.; Chung, A.; Ramachandran, D.; Savaglio, M.; Skouteris, H. Weight stigma and obesity-related policies: A systematic review of the state of the literature. *Obes. Rev.* **2021**, *22*, e13333. [CrossRef] [PubMed]
25. Puhl, R.M. Weight Stigma and Barriers to Effective Obesity Care. *Gastroenterol. Clin. N. Am.* **2023**, *52*, 417–428. [CrossRef]
26. Puhl, R.M.; Lessard, L.M. Weight Stigma in Youth: Prevalence, Consequences, and Considerations for Clinical Practice. *Curr. Obes. Rep.* **2020**, *9*, 402–411. [CrossRef] [PubMed]
27. Puhl, R.M.; Lessard, L.M.; Pearl, R.L.; Himmelstein, M.S.; Foster, G.D. International comparisons of weight stigma: Addressing a void in the field. *Int. J. Obes.* **2021**, *45*, 1976–1985. [CrossRef] [PubMed]
28. Durso, L.E.; Latner, J.D. Understanding Self-directed Stigma: Development of the Weight Bias Internalization Scale. *Obesity* **2008**, *16*, S80–S86. [CrossRef]
29. Bidstrup, H.; Brennan, L.; Kaufmann, L.; de la Piedad Garcia, X. Internalised weight stigma as a mediator of the relationship between experienced/perceived weight stigma and biopsychosocial outcomes: A systematic review. *Int. J. Obes.* **2022**, *46*, 1–9. [CrossRef]
30. Callahan, D. Obesity: Chasing an Elusive Epidemic. *Hastings Cent. Rep.* **2013**, *43*, 34–40. [CrossRef]
31. Vartanian, L.R.; Pinkus, R.T.; Smyth, J.M. The phenomenology of weight stigma in everyday life. *J. Context. Behav. Sci.* **2014**, *3*, 196–202. [CrossRef]
32. Vartanian, L.R.; Smyth, J.M. Primum Non Nocere: Obesity Stigma and Public Health. *J. Bioethical Inq.* **2013**, *10*, 49–57. [CrossRef]
33. Alimoradi, Z.; Golboni, F.; Griffiths, M.D.; Broström, A.; Lin, C.-Y.; Pakpour, A.H. Weight-related stigma and psychological distress: A systematic review and meta-analysis. *Clin. Nutr.* **2020**, *39*, 2001–2013. [CrossRef]
34. Myers, A.; Rosen, J.C. Obesity stigmatization and coping: Relation to mental health symptoms, body image, and self-esteem. *Int. J. Obes.* **1999**, *23*, 221–230. [CrossRef] [PubMed]
35. Pearl, R.L.; Puhl, R.M.; Himmelstein, M.S.; Pinto, A.M.; Foster, G.D. Weight Stigma and Weight-Related Health: Associations of Self-Report Measures Among Adults in Weight Management. *Ann. Behav. Med.* **2020**, *54*, 904–914. [CrossRef]
36. Phelan, S.M.; Burgess, D.J.; Puhl, R.; Dyrbye, L.N.; Dovidio, J.F.; Yeazel, M.; Ridgeway, J.L.; Nelson, D.; Perry, S.; Przedworski, J.M.; et al. The Adverse Effect of Weight Stigma on the Well-Being of Medical Students with Overweight or Obesity: Findings from a National Survey. *J. Gen. Intern. Med.* **2015**, *30*, 1251–1258. [CrossRef]
37. Vartanian, L.R.; Pinkus, R.T.; Smyth, J.M. Experiences of weight stigma in everyday life: Implications for health motivation. *Stigma Health* **2018**, *3*, 85–92. [CrossRef]
38. Flint, S.W.; Hudson, J.; Lavalley, D. The portrayal of obesity in U.K. national newspapers. *Stigma Health* **2016**, *1*, 16–28. [CrossRef]
39. Rao, A.; Puhl, R.; Farrar, K. Weight Stigma in Online News Images: A Visual Content Analysis of Stigma Communication in the Depictions of Individuals with Obesity in U.S. and U.K. News. *J. Health Commun.* **2024**, *29*, 95–106. [CrossRef]
40. Beaumont, J.D.; Wyld, R.; Reimann, T.; O'Hara, B. Exploring the perceptions of health, weight and obesity. *Int. J. Obes.* **2024**, *48*, 36. [CrossRef]
41. Bally, E.; Boniface, E. Weight Stigma in Leeds. Available online: <https://observatory.leeds.gov.uk/health-and-wellbeing/ph-documents/> (accessed on 26 February 2025).
42. Forbes, Y.; Donovan, C. The role of internalised weight stigma and self-compassion in the psychological well-being of overweight and obese women. *Aust. Psychol.* **2019**, *54*, 471–482. [CrossRef]
43. Remmert, J.E.; Convertino, A.D.; Roberts, S.R.; Godfrey, K.M.; Butryn, M.L. Stigmatizing weight experiences in health care: Associations with BMI and eating behaviours. *Obes. Sci. Pract.* **2019**, *5*, 555–563. [CrossRef]
44. Stunkard, A.J.; Sørensen, T.; Schulsinger, F. Use of the Danish Adoption Register for the study of obesity and thinness. *Res. Publ.—Assoc. Res. Nerv. Ment. Dis.* **1983**, *60*, 115–120.

45. Craig, C.L.; Marshall, A.L.; Sjöström, M.; Bauman, A.E.; Booth, M.L.; Ainsworth, B.E.; Pratt, M.; Ekelund, U.L.F.; Yngve, A.; Sallis, J.F.; et al. International Physical Activity Questionnaire: 12-Country Reliability and Validity. *Med. Sci. Sports Exerc.* **2003**, *35*, 1381–1395. [[CrossRef](#)] [[PubMed](#)]
46. Allison, D.B.; Basile, V.C.; Yunker, H.E. The measurement of attitudes toward and beliefs about obese persons. *Int. J. Eat. Disord.* **1991**, *10*, 599–607. [[CrossRef](#)]
47. Lillis, J.; Luoma, J.B.; Levin, M.E.; Hayes, S.C. Measuring Weight Self-stigma: The Weight Self-stigma Questionnaire. *Obesity* **2010**, *18*, 971–976. [[CrossRef](#)] [[PubMed](#)]
48. Latner, J.D.; O'Brien, K.S.; Durso, L.E.; Brinkman, L.A.; MacDonald, T. Weighing obesity stigma: The relative strength of different forms of bias. *Int. J. Obes.* **2008**, *32*, 1145–1152. [[CrossRef](#)]
49. Braun, V.; Clarke, V. *Thematic Analysis: A Practical Guide*; SAGE: Los Angeles, CA, USA, 2022.
50. Field, A.E.; Camargo, C.A., Jr.; Ogino, S. The Merits of Subtyping Obesity: One Size Does Not Fit All. *JAMA* **2013**, *310*, 2147–2148. [[CrossRef](#)] [[PubMed](#)]
51. Tinetti, M.E.; Hladek, M.d.; Ejem, D. One Size Fits All—An Underappreciated Health Inequity. *JAMA Intern. Med.* **2024**, *184*, 7–8. [[CrossRef](#)]
52. Svalastog, A.L.; Donev, D.; Jähren Kristoffersen, N.; Gajović, S. Concepts and definitions of health and health-related values in the knowledge landscapes of the digital society. *Croat. Med. J.* **2017**, *58*, 431–435. [[CrossRef](#)]
53. Andreyeva, T.; Puhl, R.M.; Brownell, K.D. Changes in Perceived Weight Discrimination Among Americans, 1995–1996 Through 2004–2006. *Obesity* **2008**, *16*, 1129–1134. [[CrossRef](#)]
54. Chaput, J.P.; Ferraro, Z.M.; Prud'homme, D.; Sharma, A.M. Widespread misconceptions about obesity. *Can. Fam. Physician* **2014**, *60*, 973–975.
55. Puhl, R.M.; Heuer, C.A. The Stigma of Obesity: A Review and Update. *Obesity* **2009**, *17*, 941–964. [[CrossRef](#)]
56. Allport, G.W. *The Nature of Prejudice*; Addison-Wesley: Oxford, UK, 1954.
57. Henry, P.J.; Napier, J.L. Education is Related to Greater Ideological Prejudice. *Public Opin. Q.* **2017**, *81*, 930–942. [[CrossRef](#)]
58. Bevan, S. Obesity is one of the last acceptable forms of discrimination—But should it be a protected characteristic in the workplace? *The Independent*, 14 January 2019.
59. Kahan, S. *The Perils of Obesity Prejudices*; Huffpost: New York, NY, USA, 2015.
60. Sampath, S. Weight bias—The last accepted form of discrimination. *Br. Columbia Med. J.* **2019**, *69*, 349.
61. Bacon, L. *Health at Every Size: The Surprising Truth About Your Weight, Rev. and Upd*; BenBella Books: Dallas, TX, USA, 2010; p. 374.
62. Ler, P.; Ojalehto, E.; Zhan, Y.; Finkel, D.; Dahl Aslan, A.K.; Karlsson, I.K. Conversions between metabolically unhealthy and healthy obesity from midlife to late-life. *Int. J. Obes.* **2024**, *48*, 433–436. [[CrossRef](#)]
63. Mathew, H.; Farr, O.M.; Mantzoros, C.S. Metabolic health and weight: Understanding metabolically unhealthy normal weight or metabolically healthy obese patients. *Metabolism* **2016**, *65*, 73–80. [[CrossRef](#)]
64. Stefan, N. Metabolically Healthy and Unhealthy Normal Weight and Obesity. *Endocrinol. Metab.* **2020**, *35*, 487–493. [[CrossRef](#)]
65. Moore, L.L.; Chadid, S.; Singer, M.R.; Kreger, B.E.; Denis, G.V. Metabolic Health Reduces Risk of Obesity-Related Cancer in Framingham Study Adults. *Cancer Epidemiol. Biomark. Prev.* **2014**, *23*, 2057–2065. [[CrossRef](#)]
66. Park, Y.-M.M.; White, A.J.; Nichols, H.B.; O'Brien, K.M.; Weinberg, C.R.; Sandler, D.P. The association between metabolic health, obesity phenotype and the risk of breast cancer. *Int. J. Cancer* **2017**, *140*, 2657–2666. [[CrossRef](#)] [[PubMed](#)]
67. Baez, A.S.; Ortiz-Whittingham, L.R.; Tarfa, H.; Osei Baah, F.; Thompson, K.; Baumer, Y.; Powell-Wiley, T.M. Social determinants of health, health disparities, and adiposity. *Prog. Cardiovasc. Dis.* **2023**, *78*, 17–26. [[CrossRef](#)]
68. Kalra, S.; Verma, M.; Kapoor, N. Commercial determinants of health: A critical component of the obesogenic environment. *Clin. Epidemiol. Glob. Health* **2023**, *23*, 101367. [[CrossRef](#)]
69. Williams, M.S.; McKinney, S.J.; Cheskin, L.J. Social and Structural Determinants of Health and Social Injustices Contributing to Obesity Disparities. *Curr. Obes. Rep.* **2024**, *13*, 617–625. [[CrossRef](#)]
70. Pearl, R.L. Weight Bias and Stigma: Public Health Implications and Structural Solutions. *Soc. Issues Policy Rev.* **2018**, *12*, 146–182. [[CrossRef](#)]
71. Åberg, E.; Koivula, A.; Kukkonen, I. A feminine burden of perfection? Appearance-related pressures on social networking sites. *Telemat. Inform.* **2020**, *46*, 101319. [[CrossRef](#)]
72. Schaefer, L.M.; Burke, N.L.; Anderson, L.M.; Thompson, J.K.; Heinberg, L.J.; Bardone-Cone, A.M.; Neyland, M.K.H.; Frederick, D.A.; Anderson, D.A.; Schaumberg, K.; et al. Comparing internalization of appearance ideals and appearance-related pressures among women from the United States, Italy, England, and Australia. *Eat. Weight. Disord.—Stud. Anorex. Bulim. Obes.* **2019**, *24*, 947–951. [[CrossRef](#)] [[PubMed](#)]
73. Falkner, N.H.; French, S.A.; Jeffery, R.W.; Neumark-Sztainer, D.; Sherwood, N.E.; Morton, N. Mistreatment Due to Weight: Prevalence and Sources of Perceived Mistreatment in Women and Men. *Obes. Res.* **1999**, *7*, 572–576. [[CrossRef](#)] [[PubMed](#)]
74. Puhl, R.M.; Andreyeva, T.; Brownell, K.D. Perceptions of weight discrimination: Prevalence and comparison to race and gender discrimination in America. *Int. J. Obes.* **2008**, *32*, 992–1000. [[CrossRef](#)] [[PubMed](#)]

75. Puhl, R.M.; Brownell, K.D. Confronting and Coping with Weight Stigma: An Investigation of Overweight and Obese Adults. *Obesity* **2006**, *14*, 1802–1815. [[CrossRef](#)] [[PubMed](#)]
76. Jackson, S. Obesity, Weight Stigma and Discrimination. *J. Obes. Eat. Disord.* **2016**, *2*, 6. [[CrossRef](#)]
77. Myre, M.; Berry, T.R.; Ball, G.D.C.; Hussey, B. Motivated, Fit, and Strong—Using Counter-Stereotypical Images to Reduce Weight Stigma Internalisation in Women with Obesity. *Appl. Psychol. Health Well-Being* **2020**, *12*, 335–356. [[CrossRef](#)]
78. Dryer, R.; Ware, N. Beliefs about causes of weight gain, effective weight gain prevention strategies, and barriers to weight management in the Australian population. *Health Psychol. Behav. Med.* **2014**, *2*, 66–81. [[CrossRef](#)]
79. Vartanian, L.R.; Porter, A.M. Weight stigma and eating behavior: A review of the literature. *Appetite* **2016**, *102*, 3–14. [[CrossRef](#)]
80. Vartanian, L.R.; Shaprow, J.G. Effects of Weight Stigma on Exercise Motivation and Behavior: A Preliminary Investigation among College-aged Females. *J. Health Psychol.* **2008**, *13*, 131–138. [[CrossRef](#)] [[PubMed](#)]
81. Alberga, A.S.; Edache, I.Y.; Forhan, M.; Russell-Mayhew, S. Weight bias and health care utilization: A scoping review. *Prim. Health Care Res. Dev.* **2019**, *20*, e116. [[CrossRef](#)] [[PubMed](#)]
82. Rudolph, C.W.; Wells, C.L.; Weller, M.D.; Baltes, B.B. A meta-analysis of empirical studies of weight-based bias in the workplace. *J. Vocat. Behav.* **2009**, *74*, 1–10. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.