



LEEDS
BECKETT
UNIVERSITY

Citation:

Boduszek, D and Dhingra, K and Hyland, P and Debowska, A (2016) A Bifactorial Solution to the Psychopathy Checklist: Screening Version in a Sample of Civil Psychiatric Patients. *Criminal Behaviour and Mental Health*, 26 (3). pp. 174-185. ISSN 1471-2857 DOI: <https://doi.org/10.1002/cbm.1956>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/1654/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

**A Bifactorial Solution to the Psychopathy Checklist Screening Version in a
Sample of Civil Psychiatric Patients**

Daniel Boduszek¹, Katie Dhingra², Philip Hyland³, & Agata Debowska⁴

¹University of Huddersfield, Huddersfield, United Kingdom

²Manchester Metropolitan University Manchester, United Kingdom

³National College of Ireland, Dublin, Ireland

⁴University of Chester, Chester, United Kingdom

Version accepted for publication in *Criminal Behaviour and Mental Health*

Correspondence to:

Dr Katie Dhingra

Manchester Metropolitan University

Department of Psychology

Manchester, M15 6GX

United Kingdom

Tel: +44 (0)1612472593

Email: K.Dhingra@mmu.ac.uk

ABSTRACT

Background: There is considerable debate about the underlying factor structure of the Psychopathy Checklist: Screening Version (PCL: SV). An established view is that it reflects a unitary construct underpinned by two correlated factors. More recent research has, however, undermined this conceptualisation.

Aims: Our aim was to compare 10 competing models of the PCL: SV in a sample of civil psychiatric patients.

Method: Ten distinct factor models were specified and tested using conventional confirmatory factor analytic techniques, along with confirmatory bifactor modelling.

Results: A bifactor model, including two general factors (interpersonal-affective and antisocial-lifestyle), and four subordinate factors (interpersonal, affective, antisocial, and lifestyle) provided the best fit to the data. The reliability of the conceptualisation was supported through the use of composite reliability, and the differential relationships exhibited between the general factors and measures of personality, impulsivity, and mental health.

Conclusions: The results suggest that two general factors should be taken into account when interpreting the PCL:SV for clinical purposes.

Keywords: PCL: SV; Bifactorial Modelling; Confirmatory Factor Analysis; Psychopathy.

INTRODUCTION

The Psychopathy Checklist: Screening Version (PCL: SV; Hart et al., 1995) is a 12-item version of the Psychopathy Checklist-Revised (PCL-R; Hare, 1991), created for use as a stand-alone instrument for research with non-offenders, or as a screen for psychopathy in offender populations. It was developed to assess a unitary psychopathy construct formed by two correlated factors ($r \sim 0.50$ Hare, 1991). Factor 1 reflects interpersonal and affective traits, whereas Factor 2 reflects a socially deviant lifestyle. Although there is some evidence for this two-factor structure (e.g., Harpur et al., 1989; Hart et al., 1995; Skeem & Mulvey, 2001), there is some against it (e.g., Dolan & Anderson, 2003; Forth et al., 1996).

More recent research suggests that either a three (Cooke & Michie, 2001) or four-factor model (Hare, 2003) may better represent the structure of the PCL: SV than the two-factor approach. Cooke and Michie's (2001) 3-factor model divided the original Factor 1 into an interpersonal factor (*arrogant and deceitful interpersonal style*) and an affective factor (*deceitful affective experience*) and combined the original Factor 2 items (except for the antisocial behaviour items) to form an *impulsive and irresponsible behaviour style* factor. In Hare's (2003) four-factor model, Factor 1 of the original two-factor model is divided into 'interpersonal' and 'affective' facets and Factor 2 is into a 'lifestyle' facet and an 'antisocial' facet. Studies that are even more recent have found a good fit for this 4-factor model (e.g., Forth et al., 2003; Neumann & Hare, 2008; Vitacco et al., 2006; Žukauskienė et al., 2010). Association between the four-factors is, however, generally high – for example from $r = 0.45$ between interpersonal and antisocial latent variables to $r = 0.99$ between lifestyle and antisocial latent variables (Žukauskienė et al., 2010), implying the presence of a higher-order psychopathy factor or a general factor with four method-factors.

The inconsistent and unsatisfactory model fit reported in the literature suggests that traditional confirmatory factor analysis (CFA) methods are not sufficient to explain the dimensionality of the PCL: SV. Thus, some researchers (e.g., Flores-Mendoza et al., 2008; Patrick et al., 2007) have used an alternative model structure to the PCL-R which may yield a theoretically and statistically satisfactory solution. This involves the application of bifactor modelling (see Reise et al., 2010). Within a bifactorial modelling approach, covariation among items is presumed to be explained by both ‘general factors’ (the source of common variance running through all measure items) and separate uncorrelated grouping factors that reflect the unique coherency among particular subgroups of items. Thus, the bifactor approach differs from the higher-order model approach in that subfactors are not subsumed by the general factor(s) but are, instead, uncorrelated and distinct. Consequently, if a bifactor model is found to provide a statistically superior fit to the data than alternative models tested, this indicates that: (a) the domain being modelled is saturated by one or more broad factor(s) that reflects the common variance running among all scale items, and (b) specific scales in the domain are also saturated by other specific (i.e. residual) uncorrelated factors that reflect additional common variance among clusters of items, typically, with highly similar content. Thus, in a bifactor model, each scale is a measure of the general factor(s), but some scales also index other more specific constructs that are not accounted for by the general factor(s).

Patrick et al. (2007) found that a bifactor model including a single general “psychopathy” factor and three subfactors factors (interpersonal, affective, and impulsivity) provided a better fit to the data than alternative models tested. Similarly, Flores-Mendoza et al. (2008) reported that the bifactorial solution was a better representation of the data than any other tested model in a sample of 124 male prisoners. The later had, however, removed non-

significant loadings from their model, which may suggest that their solution was a poor approximation of their data.

Our aim was to provide a more accurate determination of the optimal number of factors necessary to explain the dimensionality of the PCL: SV. We therefore investigated a series of theoretically plausible models of the underlying structure of the PCL: SV (see Cooke et al., 1999; Debowska et al., 2014; Forth et al., 1996; Flores-Mendoza et al., 2008; Hare, 2003; Harpur et al., 1989; Neuman & Hare, 2008; Patrick et al., 2007), including bifactor models which have not previously been empirically tested but are in-line with theoretical formulations.

Method

Sample

Participants were 1,136 civil psychiatric patients sampled from one of three acute inpatient hospitals as part of the MacArthur Violence Risk Assessment Study. Inclusion criteria were age 18–40, having English as a first language, having been hospitalised for less than 21 days and having a records diagnosis of schizophrenia, schizophreniform disorder, schizoaffective disorder, major depression, dysthymia, mania, brief reactive psychosis, delusional disorder, alcohol or other drug abuse or dependence or personality disorder. A total of 1,695 patients met the inclusion criteria, but just 71% agreed to participate (for more information on the data collection method and sample characteristics see Monahan et al., 2001).

For our study, we used data from baseline and two follow-up interviews. After excluding data from participants who had not been administered PCL: SV, we were left with data for analysis from 871 patients (502 men, 369 women).

Measures

Psychopathy was assessed by trained raters using the 12-item PCL: SV (Hart et al., 1995), using semi-structured interviews supplemented by records data. Each item is rated on a 3-point scale (0 *does not apply*, 1 *applies to a certain extent*, 2 *applies*). The PCL: SV has good reliability and validity and is strongly related to the PCL-R (Cooke et al., 1999; Guy & Douglas, 2006).

Criminal Behaviour. Three items were used as an indication of criminal behaviour. The first was arrests for crimes against persons and the second crimes against property, both coded dichotomously from official criminal records. The third reflected several categories of violent behaviour in the 10 weeks prior to baseline interview (for details, see Monahan et al., 2001).

The NEO-Five Factor Inventory, Short Form (NEO-PI-SF) (Costa & McCrae, 1992) is a 60-item inventory which taps the "big 5" dimensions of personality: neuroticism, extraversion, openness, agreeableness, and conscientiousness. Items that composed these scales had acceptable levels of internal consistency.

The Wechsler Abbreviated Scale of Intelligence (WASI) (Wechsler, 1999) produces an estimate of general intelligence, with higher scores indicating greater intellectual ability.

Psychiatric symptoms. The Brief Symptom Inventory (BSI; Derogatis, 1993) is a 16-item rating scale used to assess psychiatric symptoms along a 7-point scale at the time of interview. Its subscales are: activation, thought disturbance, hostile-suspiciousness, anergia, and anxiety-depression.

Analysis

The dimensionality of the PCL: SV was investigated through the use of conventional confirmatory factor analytic (CFA) techniques, along with confirmatory bifactor modelling (see Reise, et al., 2010). Ten alternative models of the latent factor structure of the PCL: SV

were specified and estimated using Mplus version 6.12 (Muthen & Muthen, 1998 – 2010) with maximum likelihood (ML) estimation. Six of these models were ‘traditional’ in CFA terms. Within these models, items were restricted to load only onto a single factor; while in the bifactor models, each item was allowed to load onto one or two general factors and four subordinate factors (interpersonal, affective, lifestyle, antisocial), as recommended by Reise, et al (2010). Thus, in bifactor models, each item is a measure of the general factor, as well as more specific constructs that are not correlated with the general factor. In all cases measurement error terms remained uncorrelated, as suggested in previous research (Boduszek et al., 2013; Boduszek et al., 2012; Bollen, 1989; Brown, 2006).


Model 1 is a one-factor solution in which the twelve items of the PCL: SV load onto a single latent variable of psychopathy. Model 2 is a correlated two-factor model in which the two latent variables are represented by psychopathy Factor 1 (interpersonal-affective; items 1, 2, 3, 4, 5, and 6) and psychopathy Factor 2 (antisocial-lifestyle; items 7, 8, 9, 10, 11, and 12). Model 3 is a correlated three-factor model in which the three latent variables of psychopathy are represented by an interpersonal factor (items 1, 2, and 3), an affective factor (items 4, 5, and 6), and a lifestyle factor (items 7, 9, and 10). Model 4 is a correlated four-factor model in which the four latent variables of psychopathy are represented by an interpersonal factor (items 1, 2, and 3), an affective factor (items 4, 5, and 6), a lifestyle factor (items 7, 9, and 10), and an antisocial factor (items 8, 11, and 12). Models 5 and 6 are higher-order models with one and two superordinate latent factors respectively. Model 7 is a bifactor conceptualisation containing five latent factors; a single general factor of psychopathy and four subordinate factors represented by interpersonal, affective, antisocial, and lifestyle latent variables (items loading as above). Model 8 (see Figure 1) is also a bifactor conceptualisation containing six latent variables: two general factors (Factor 1 – items 1, 2, 3, 4, 5, and 6;

Factor 2 - items 7, 8, 9, 10, 11, and 12) and four subordinate factors (items as in Model 4).

Model 9 is a bifactor conceptualisation containing four latent factors; a single general factor of psychopathy and three subordinate factors represented by interpersonal, affective, and lifestyle latent variables (items loading as above). Model 9 is another bifactor conceptualisation containing two general factors (Factor 1 – items 1, 2, 3, 4, 5, and 6; Factor 2 - items 7, 8, 9, 10, 11, and 12) and three subordinate factors (items as in Model 3). Within a bifactor model the grouping factors are restricted to be uncorrelated with each other and uncorrelated with the general factors. For the purposes of model identification, the variance of each factor is set to 1.0.

Figure 1

Model 8: Bifactor model of the PCL:SV



NB: PCL1 = general factor 1 (Interpersonal-Affective), PCL2 = general factor 2 (Antisocial-Lifestyle), F1 = Interpersonal, F2 = Affective, F3 = Antisocial, F4 = Lifestyle.

The overall fit of each model and the relative fit between models were assessed using a range of goodness-of-fit statistics. The chi-square (χ^2) statistic assesses the sample and implied covariance matrix; a model with good fit is indicated by a non-significant result. As the chi-square statistic is strongly associated with sample size, however, good models tend to be over-rejected. Therefore, Tanaka (1987) suggested that a model should not be rejected simply on the basis of a significant chi-square result. The Comparative Fit Index (CFI; Cronbach, 1990) and the Tucker Lewis Index (TLI; Tucker & Lewis, 1973) are measures of how much better the model fits the data compared to a baseline model where all variables are uncorrelated. For these indices, values above 0.95 indicate good model fit (Bentler, 1990; Hu & Bentler, 1999). In addition, two more absolute indices are presented: the standardised root mean-square residual (SRMR; Joreskog & Sorbom, 1981) and the root mean-square error of approximation (RMSEA; Steiger, 1990). Ideally, these indices should be less than 0.05 to suggest good fit (Bentler, 1990; Hu & Bentler, 1999; Joreskog & Sorbom, 1993). Furthermore, the Akaike Information Criterion (AIC; Akaike, 1974) was used to evaluate the alternative models, with the smaller value indicating the best fitting model.

Results

General characteristics of the sample.

Participants in this sample were between the ages of 18–40 years ($M = 29.86$, $SD = 6.20$).

The mean total PCL:SV score is 8.52 ($SD = 5.61$, Median = 8, Min = 0, Max = 24), and 72 participants are above the cut-off score of 14. The mean general factor 1 score is 3.11 ($SD = 2.99$, Median = 2, Min = 0, Max = 12); while the mean general factor 2 score is 5.41 ($SD =$

3.30, Median = 5, Min = 0, Max = 12). Cronbach's alpha coefficients for general factor 1 and general factor 2 are 0.84 and 0.82, respectively.

Model Results

Table 1 shows the fit indices and comparative fit indices of the ten alternative models of the PCL: SV. Based on these findings, Models 1 to 6 and 9 were rejected as poor approximations of the data. Models 7, 8, and 10 were found to provide good representations, with Model 8 providing the best fit to the data. This model (Model 8), which includes two general factors of psychopathy and four subordinate factors, was determined to be a good approximation of the covariation matrix in the obtained data based upon all fit indices. In an analysis including only the women in the sample, the bifactorial solution was again statistically superior to the alternative models tested ($\chi^2 = 60.06$, $p = 0.03$; RMSEA 0.04 [90% CI 0.02-0.05]; SRMR 0.03; CFI 0.99; TLI 0.98).

The adequacy of bifactor model (total sample) can also be determined in relation to its parameter estimates. As shown in Table 2, all items displayed statistically significant ($p < 0.001$) factor loadings on the two general psychopathy factors. Factor loading were all in the expected direction and all items displayed factor loading above 0.4. Further inspection of the factor loadings for the four subordinate factors provides critical information regarding the appropriateness of including these factors in the scoring of the PCL: SV. Reise et al. (2010) advise that when items load strongly onto a general factor (or factors), and less strongly on each of the subordinate factors, this demonstrates the superiority of the general factors over the grouping factors in the conceptualisation of the factor structure of the scale, and thus its related scoring scheme. Alternatively, when items load as strongly (or more strongly) onto

each of the respective subordinate factors than onto the general factor, creation of subscales based on these factors can be considered appropriate.

As outlined in Table 2, factor loadings for each subordinate factor were poorer than those on the two general factors. These parameter estimate results provide strong support for the supremacy of a model containing two general factors, and the presence of four substantively meaningful subordinate factors. The two general factors were moderately correlated ($r = 0.45$). The four grouping factors were kept uncorrelated as suggested by Reise et al. (2010).

Further analysis examined the relationships between the PCL: SV factors and external variables (Table 3). Results indicate that general factor 2 but not general factor 1 is significantly, if weakly, associated with neuroticism ($r = 0.10$), openness ($r = -0.11$), conscientiousness ($r = -0.19$), BIS cognitive ($r = 0.22$), and anxiety-depression ($r = 0.09$).

Reliability Analysis

Composite reliability calculations indicate that the general factor 1 ($\rho_c = 0.84$) and general factor 2 ($\rho_c = 0.86$) of the PCL: SV possesses satisfactory composite reliability.

Table 1

Fit Indices for Ten Alternative Models of the PCL: SV

	χ^2	<i>df</i>	CFI	TLI	RMSEA	SRMR	AIC
<i>Models</i>							
1factor	967.98*	54	.76	.72	.14	.08	19418.97
Correlated 2 factors	514.18*	53	.89	.86	.10	.05	18967.17
Correlated 3 factors	230.72*	24	.93	.89	.10	.05	13937.92 ⁺
Correlated 4 factors	357.61*	48	.92	.90	.09	.05	18820.60
1 higher-order factor, 4 correlated factors	486.20*	50	.89	.86	.10	.06	18945.19
2 higher-order factors, 4 correlated factors	357.72*	49	.92	.90	.09	.05	18818.71
Bifactorial with 1 general + 4 subordinate factors	245.51*	42	.95	.92	.08	.05	18720.49
Bifactorial with 2 general + 4 subordinate factors	104.84*	41	.98	.98	.04	.02	18581.83
Bifactorial with 1 general + 3 subordinate factors	374.27*	45	.92	.88	.09	.06	18843.26
Bifactorial with 2 general + 3 subordinate factors	165.59*	44	.97	.96	.06	.04	18636.57

Note. $N = 871$; χ^2 = chi square goodness of fit statistic; *df* = degrees of freedom; RMSEA = Root-Mean-Square Error of Approximation; CI = Confidence Interval; AIC = Akaike Information Criterion; CFI = Comparative Fit Index; TLI = Tucker Lewis Index; SRMR = Standardized Square Root Mean Residual. * Indicates χ^2 are statistically significant ($p < .05$). ⁺ As the 3-factor model is based on a different set of items and, therefore, a different covariance matrix, direct statistical comparison of this model with the alternative models tested is not possible (Brown, 2006; Kline, 1998).

Table 2

Standardized Factor Loadings for the four Subordinate (F1- F4) and two General Factors (PCL1 and PCL2) of the PCL: SV

Item	F1	F2	F3	F4	PCL1	PCL2
1. Superficial	.72***				.61***	
2. Grandiose	.19				.48***	
3. Deceitful	.11				.68***	
4. Lacks remorse		.47***			.78***	
5. Lacks empathy		.43***			.61***	
6. Doesn't accept responsibility		.17**			.79***	
7. Impulsive			.11*			.71***
9. Lack goals			.56***			.51***
10. Irresponsibility			.35***			.70***
8. Poor behavioural controls				.15*		.62***
11. Adolescent antisocial behaviour				.90**		.48***
12. Adult antisocial behaviour				.22**		.73***

Note. Factor loadings are statistically significant at *** $p < .001$, ** $p < .01$, * $p < .05$

Table 3

Relationships between PCL: SV factors and external variables

Variable	F1	F2	F3	F4	PCL1	PCL2
NEO-Neuroticism	-.06	.01	.11**	.07	-.02	.10**
NEO-Extraversion	.12**	-.02	-.03	.02	.05	.01
NEO-Openness	.06	-.10**	-.12**	-.08*	-.02	-.11**
NEO-Agreeableness	-.24**	-.34**	-.29**	-.37**	-.32**	-.36**
NEO-Conscientiousness	.02	-.06	-.19**	-.14**	-.02	-.19**
WAIS-R	-.08*	-.22**	-.29**	-.20**	-.17**	-.28**
BIS Motor	.10**	.17**	.24**	.27**	.15**	.29**
BIS non-planning	.10**	.15**	.33**	.31**	.14**	.36**
BIS cognitive	.01	.09**	.19**	.20**	.06	.22**
BPRS Activation	.12**	.17**	.17**	.15**	.16**	.18**
BPRS thought disturbance	.10**	.09*	.07*	.08*	.11**	.08*
BPRS hostile-suspiciousness	.09*	.17**	.09**	.21**	.15**	.17**
BPRS Anergia	-.10**	.06	.03	-.04	-.02	-.01
BPRS anxiety-depression	-.01	.01	.08*	.08*	.01	.09*
Number of violent acts	.12**	.21**	.18**	.24**	.18**	.23**
Crime against people	.05	.07*	.06	.07*	.07	.07
Crime against property	.16**	.17**	.14**	.14**	.18**	.17**

Note. ** $p < .01$, * $p < .05$

Discussion

Crucial to the study of any psychological construct is the clear delineation of its underlying structure. This is necessary not only for the interpretation of scores on a measure, but also because dimensions of a construct may differentially relate to external variables (Reise, 1999) and because inaccurate factor conceptualisations may result in unstable estimates of reliability (see Shevlin et al., 2000). Our study was carried out to provide a methodologically rigorous investigation of the dimensionality of the PCL: SV (Hart et al., 1995), a frequently used measure of personality traits, which has been at the centre of much debate with respect to the appropriate latent structure of the scale. Based on the inappropriateness of including correlated measurement errors in factorial models (see Boduszek et al., 2012, 2013), we examined the 10 conceptualisations of the structure of the PCL: SV, including four bifactorial solutions.

Fit indices indicated that the bifactor model with two general factors (*interpersonal-affective* and *antisocial-lifestyle*) and four independent subordinate factors (interpersonal, affective, antisocial, lifestyle), each of which account for unique variance in their respective set of items over and above the variance accounted for by the general factors, was a superior representation of the underlying factor structure of the PCL: SV than the alternative factor solutions tested. This suggests that the PCL: SV represents a two-dimensional construct with four distinct domains of item content (subordinate factors) that, among general psychiatric patients, vary independently from the general factors. This conceptualisation is theoretically satisfying as it is consistent with Hare's (1991) earlier 2-factor characterisation of psychopathy. Furthermore, it is consistent with the notion that psychopathy is not a unitary construct, but is instead a combination of two continuous independent, however statistically correlated, latent factors. To the extent that the bifactorial solution is more parsimonious than

other tested models, the results of this study suggest supporting the utility of two general factors model of psychopathy for clinical interpretive purposes. This is important because delineating the latent variables that define a construct is necessary to inform risk assessment and treatment options. Although the present findings suggest that antisocial tendencies (general factor 2) are important features of the psychopathy construct, further research is warranted in this area. For instance, research is needed to determine whether antisocial tendencies are simply a consequence of other psychopathic traits, or whether antisocial features influence the nature and development of other psychopathic features (general factor 1).

The appropriateness of this factorial solution was supported by the differential relationship between the two general factors and measures of personality, impulsivity, and mental health. As suggested by Carmines and Zeller (1979), if factors measure substantially different dimensions, they should differentially relate to external variables. We found that only general Factor 2 was significantly related to neuroticism, openness, conscientiousness, BIS cognitive, and anxiety-depression.

Parameter estimates also highlight the need to consider two meaningful general factors when applying the PCL: SV in research contexts. Failing to control for systematic error variance which arises due to the presence of the subordinate factors could not only prevent identification of a theoretically consistent and logical factor structure, but ultimately lead to inaccurate interpretations about the relationships between the scale factors and various external variables. Future research should, therefore, seek to compare the predictive effect of PCL: SV modelled as a four-factor structure, as is currently standard practice, and the predictive effect of the two general factors when the four subordinate factors are modelled

and controlled for within the bifactorial conceptualisation. The two general factors also showed good reliability, as assessed using composite reliability (Novick & Lewis, 1967).

It is important to note that our analysis was based on data from adult general psychiatric patients who had completed very short inpatient admissions after a mental health crisis; our findings may not be generalisable to any other group. Future research might apply the same statistical approaches to data from people who never require hospital admission for mental disorder, people with primary personality disorder, offender patients and/or adolescents and, in particular, check whether such approaches actually provide any improvement in capacity to aid assessment of risk of maladaptive or criminal behaviour.

In conclusion, we rigorously investigated the factor structure of the PCL: SV with one large group of recently discharged psychiatric patients. We found that the PCL: SV has two general factors, on which the items of the original two-factor model load (six items on each factor), along with four separate subordinate factors on which the items of the four-factor model load. We also found that the two general factors are differentially associated with external variables, and provided empirical support for the value of adopting a bifactor modelling approach when assessing the dimensionality of this measure.

References

- Akaike H (1974) A new look at the statistical model identification. *IEEE Transactions on Automatic Control* **19**:716–723.
- Bentler PM (1990) Comparative fit indices in structural models. *Psychological Bulletin* **107**: 238–246.
- Boduszek D, Hyland P, Dhingra K, Mallett J (2013) The Factor Structure and Composite Reliability of the Rosenberg Self-Esteem Scale among Ex-Prisoners. *Personality and Individual Differences* **55**: 877-881.
- Boduszek D, Shevlin M, Mallett J, Hyland, P, O’Kane D (2012) Dimensionality and construct validity of the Rosenberg self-esteem scale within a sample of recidivistic prisoners. *Journal of Criminal Psychology* **2**:19-25.
- Bollen KA (1989) *Structural equations with latent variables*. New York: Wiley.
- Brown TA (2006) *Confirmatory factor analysis for applied research*. New York: The Guilford Press.
- Carmines EG, & Zeller RA (1979) *Reliability and validity assessment*. Beverly Hills, CA: Sage.
- Cooke DJ, & Michie C (2001) Refining the construct of psychopathy: towards a hierarchical model. *Psychological assessment* **13**:171-188.
- Cooke DJ, Michie C, Hart SD, Hare RD (1999) Evaluating the Screening Version of the Hare Psychopathy Checklist—Revised (PCL: SV): An item response theory analysis. *Psychological Assessment* **11**:3-13.
- Costa, T, & McCrae, RR. (1992). Normal personality assessment in clinical practice: The NEO Personality Inventory. *Psychological assessment* **4**: 5-13.
- Cronbach LJ (1951) Coefficient alpha and the internal structure of tests. *Psychometrika*, **16**:

297-334.

Debowska A, Boduszek D, Kola S, & Hyland P (2014) A Bifactor Model of the Polish Version of the Hare Self-Report Psychopathy Scale. *Personality and Individual Differences*. **69**: 231-237.

Derogatis LR, & Spencer P M. (1993). *Brief symptom inventory: BSI*. Upper Saddle River, NJ: Pearson.

Dolan MC, Anderson, IM (2003) The relationship between serotonergic function and the Psychopathy Checklist: Screening Version. *Journal of Psychopharmacology*, **17**:216-222.

Dolan M, Davies G (2006) Psychopathy and institutional outcome in patients with schizophrenia in forensic settings in the UK. *Schizophrenia research* **81**:277-281.

Flores-Mendoza C. E., Alvarenga M. A. S., Herrero Ó., & Abad F. J. (2008). Factor structure and behavioural correlates of the Psychopathy Checklist-Revised [PCL-R] in a Brazilian prisoner sample. *Personality and Individual Differences* **45**: 584-590.

Forth AE, Brown, SL, Hart, SD, Hare, RD (1996) The assessment of psychopathy in male and female noncriminals: Reliability and validity. *Personality and Individual Differences* **20**: 531-543.

Forth AE, Kosson, DS, Hare RD (2003) *Hare psychopathy checklist: Youth version (PCL: YV)*. Toronto: Multi-Health Systems.

Guy LS, Douglas KS (2006) Examining the utility of the PCL: SV as a screening measure using competing factor models of psychopathy. *Psychological assessment* **18**: 225-230.

Hare RD (1991) *The Hare Psychopathy Checklist-Revised: PLC-R*. MHS, Multi-Health Systems.

- Hare RD (2003) *Manual for the Revised Psychopathy Checklist* (2nd ed.). Toronto, Ontario, Canada: Multi-Health Systems.
- Harpur TJ, Hare RD, Hakstian AR (1989) Two-factor conceptualization of psychopathy: Construct validity and assessment implications. *Psychological Assessment: A Journal of consulting and clinical Psychology* **1**: 6-17.
- Hart S, Cox D, Hare R (1995) *Manual for the psychopathy checklist: Screening version (PCL: SV)*. Toronto: Multi-Health Systems.
- Hu L, Bentler P (1998) Fit indices in covariance structural modeling: Sensitivity to underparameterized model misspecification. *Psychological Methods*, **3**: 424- 453.
- Hu L, Bentler PM (1999) Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling* **6**: 1–55.
- Joreskog K, Sorbom D (1981) *LISREL V: Analysis of linear structural relationships by the method of maximum likelihood*. Chicago: National Educational Resources.
- Jöreskog K, Sörbom D (2004) *Interactive Lisrel (8,72)*. Scientific Software International. Inc.
- Kendler K S, Aggen SH, & Patrick CJ. (2012). A multivariate twin study of the DSM-IV criteria for antisocial personality disorder. *Biological psychiatry*, **71**: 247-253.
- Monahan J, Steadman H, Silver E, Appelbaum P, Robbins P, Mulvey E., . . . Banks S (2001) *Rethinking violence risk assessment: Mental disorder and the law*: New York: Oxford University Press.
- Muthen LK, Muthen BO (1998 - 2010) *Mplus – Statistical analysis with latent variables. User's guide* (6th Ed). Muthen and Muthen: Los Angeles.
- Neumann CS, Hare RD (2008) Psychopathic traits in a large community sample:

- links to violence, alcohol use, and intelligence. *Journal of Consulting and Clinical Psychology* **76**: 893-899.
- Novick MR, & Lewis C. (1967). Coefficient alpha and the reliability of composite measurements. *Psychometrika*, **32**: 1-13.
- Patrick C J, Fowles DC, & Krueger R F. (2009). Triarchic conceptualization of psychopathy: Developmental origins of disinhibition, boldness, and meanness. *Development and psychopathology* **21**:913-938.
- Patrick CJ, Hicks BM, Nichol PE, & Krueger RF. (2007). A bifactor approach to modeling the structure of the psychopathy checklist-revised. *Journal of personality disorders* **21**: 118-220.
- Reise SP, Moore TM, Haviland MG (2010) Bifactor models and rotations: Exploring the extent to which multidimensional data yield univocal scale scores. *Journal of Personality Assessment* **92**, 544–559.
- Shevlin ME, Miles, JNV, Davies MNO, Walker S (2000) Coefficient alpha: A useful indicator of reliability? *Personality and Individual Differences* **28**, 2, 229–238.
- Skeem JL, & Mulvey EP (2001) Psychopathy and community violence among civil psychiatric patients: results from the MacArthur Violence Risk Assessment Study. *Journal of consulting and clinical psychology* **69**(3), 358-374.
- Steiger JH (1990) Structural model evaluation and modification: An interval estimation approach. *Multivariate Behavioural Research* **25**, 173-180.
- Tanaka JS (1987) How big is big enough? Sample size and goodness of fit in structural equation models with latent variables. *Child Development* **58**, 134–146.
- Tucker LR, Lewis C (1973) The reliability coefficient for maximum likelihood factor analysis. *Psychometrika* **38**, 1-10.

Vitacco MJ, Neumann CS, Caldwell MF, Leistico AM, Van Rybroek GJ

(2006) Testing factor models of the Psychopathy Checklist: Youth Version and their association with instrumental aggression. *Journal of Personality Assessment* **87**(1), 74-83.

Wechsler, D. (1999). *Wechsler abbreviated scale of intelligence*. Psychological Corporation.

Žukauskienė R, Laurinavičius A, Čėsniėnė I (2010) Testing factorial structure and validity of the PCL: SV in Lithuanian prison population. *Journal of Psychopathology and Behavioral Assessment* **32**(3), 363-372.