



LEEDS
BECKETT
UNIVERSITY

Citation:

Pringle, A and Zwolinsky, S and McKenna, J and Daly-Smith, A and Robertson, S and White, A (2014) Reducing CVD Risk Factors for Men/Hard-to-Reach Men Using English Premier League Soccer Clubs. *MEDICINE AND SCIENCE IN SPORTS AND EXERCISE*, 46 (5). 595 - 595. ISSN 0195-9131 DOI: <https://doi.org/10.1249/01.mss.0000495256.74315.cb>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/1868/>

Document Version:

Article (Accepted Version)

Thematic poster

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Control/Tracking Number: 14-SA-2595-ACSM
Activity: Scientific Abstract
Current Date/Time: 4/23/2014 4:11:02 AM

Reducing CVD Risk Factors for Men/Hard-to-Reach Men Using English Premier League Soccer Clubs

Author Block Andy Pringle, Stephen Zwolinsky, James McKenna, Andy Daly-Smith, Steve Robertson, Alan White. *Leeds Metropolitan University, LEEDS, United Kingdom.*

Abstract:

Over two thirds of men in England UK, have two or more risk factors for cardio-vascular disease (CVD). When combined with comparatively low levels of engagement with primary health care (PC), innovative approaches and alternative strategies/places are required to connect these hard-to-reach (HTR) men with health improvement programmes.

PURPOSE: To investigate the effect of Premier League Health (PLH), a national programme of CVD risk factor reduction/prevention for men/HTR men delivered in/by English Premier League (EPL) soccer clubs.

METHODS: Health interventions were delivered to males with heightened risk of CVD and not using PC services. Interventions were underpinned by physical activity (PA) (e.g. soccer/gym/walking) and delivered alongside lifestyle educational sessions (e.g. smoking/diet) led by professional health trainers employed by the clubs. Sessions took place at 16 EPL soccer clubs. Validated self-reported data for demographics and CVD risk factors were collected at pre-intervention and 12-week follow-up.

RESULTS: Predominantly, White British men (70.4%, n=2669/3788), aged 18-44 years (80.2%, n=3032/3779) in employment (60.7%, n=1907/3143) responded. Over 85%, (n=1428/1667) presented with ≥ 2 simultaneous risk-factors for CVD. The majority of men (80%, n=1984/2481), did not perceive their health to be 'at risk'. A third, (n=860/2580) 'never' used PC and were more likely to report 3-4 CVD risk factors when compared to those who did use PC $\chi^2 [1]= 4.17, p<.05$). Table 1 shows level of change in CVD risk factors at pre vs. post-intervention.

Variable	N	Pre (SD)	Post (SD)	df	t	r
Physical Activity (Sessions/Week)	2917	2.34 (1.52)	2.49 (1.53)	2916	-13.75**	0.26
Diet: Fruit and vegetables (Daily portions)	2673	2.34 (1.38)	2.44 (1.38)	2672	-8.39**	0.16
Sitting (Hours/Day)	2223	5.63 (2.70)	5.26 (2.67)	2222	13.09**	0.27
BMI (kg/m ²)	2074	26.72 (5.50)	26.43 (5.39)	2073	10.05**	0.21
Alcohol (Units/Week)	1727	16.35 (18.67)	15.68 (18.23)	1726	7.84**	0.18

Note: **p<0.001, df= degrees of freedom, t= t-statistic, r= effect size.

CONCLUSION: PLH engaged HTR men not using PC services and showed positive change in CVD profiles for participating males.