



LEEDS
BECKETT
UNIVERSITY

Citation:

Paley, C and Johnson, MI (2016) Chronic pain in the obese population: is exercise the key? *Pain Management*, 6 (2). 121 - 123. ISSN 1758-1869 DOI: <https://doi.org/10.2217/pmt.15.64>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/2394/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Journal: Pain Management

Article type: Editorial/Review

Title:

Chronic pain in the obese population: is exercise the key?

Authors:

Carole Paley [1,2,3]

Mark I. Johnson [1,2]

Affiliations

1. Airedale NHS Foundation Trust, Skipton Road, Steeton, Keighley, West Yorkshire, BD20 6TD, UK

2. Faculty of Health and Social Sciences, Leeds Beckett University, Leeds, U.K.

2. Leeds Pallium Research Group, http://www.st-gemma.co.uk/?page_id=1910

Leeds, U.K.

Address for correspondence

Dr Carole A. Paley, Airedale NHS Foundation Trust, Skipton Road, Steeton, Keighley, West Yorkshire,
BD20 6TD, UK

Email: carole.paley@anhst.nhs.uk

INTRODUCTION

There are over 500 million adults worldwide classified as obese and a further 1.5 billion individuals classified as overweight [1]. Individuals with a high body mass index (BMI) are more likely to report chronic pain than their leaner counterparts and are more sensitive to pain [2]. Recently in this journal, Narouze and Souzdalnitski [3] reviewed pain management techniques and interventional approaches for weight reduction for individuals who are obese. They recommended that medication, rehabilitation, and interventional management strategies should be used to manage individuals with chronic pain and obesity and that this should be possible within the present infrastructure for pain management. In this article, we extend their argument by considering the role of exercise in the management of chronic pain for individuals who are overweight or obese by exploring the inter-relationships between pain, obesity, low-grade systemic inflammation and exercise.

HOW ARE CHRONIC PAIN AND OBESITY LINKED?

Evidence suggests that the positive association between chronic pain and obesity is mediated in part by abnormal loading on weight-bearing joints such as the knees, hips and spine leading to postural changes, muscle weakness and mechanical strain on connective tissues [4]. Hence, weight loss is a common goal in the management of pain in overweight and obese individuals. Weight loss through dietary modification is notoriously difficult to achieve and maintain and may be accompanied by loss of fat-free mass which can lower metabolic rate making maintenance of lower weight more difficult [5]. Exercise targets excess body fat and promotes improvements in cardiovascular risk factors even if no weight is lost [6]. Exercise also improves the sense of well-being by reducing depression and improving sleep which in turn improves the quality of life and lessens the psychological impact of chronic pain. A recent systematic review provided evidence that pain and function improved in individuals with osteoarthritis receiving exercise therapy and that these effects were mediated by increased upper leg strength, decreased extension impairments and improvements in proprioception [7]. A meta-analysis found that effect sizes of hypoalgesic outcomes associated with exercise for individuals with chronic pain were variable and dependent on diagnosis and on the intensity of the exercise [8]. However, the meta-analysis provided strong evidence that acute isometric, aerobic, and dynamic resistance exercise reduced experimentally-induced pain in healthy individuals. This reduction in pain sensitivity response in the short-term is likely to be mediated in part through activation of descending pain inhibitory systems and the release of endorphins [9]. It is also plausible that therapeutic exercise may influence systemic concentrations of pro-inflammatory

biomarkers associated with obesity and a number of studies have demonstrated this relationship (for review see [10]).

THE ROLE OF EXERCISE

The emergence of evidence that individuals who are obese may exhibit a low grade pro-inflammatory state offers new insights to the role of therapeutic exercise for these individuals. Excessive body weight contributes to adiposopathy ('sick fat') which results in elevated concentrations of pro-inflammatory biomarkers in the systemic circulation which are secreted by adipose tissue. These include prostaglandins, C-reactive protein (CRP), and cytokines such as interleukins (e.g. interleukin-6), tumor necrosis factor alpha (TNF-alpha), and leptin, all of which sensitise nociceptors resulting in an amplified nociceptive input to the central nervous system [10]. Hence, adiposopathy may contribute to non-resolving systemic inflammation and states of persistent pain. Furthermore, un-resolving systemic inflammation may contribute to continued sensitisation of the nociceptive system. Consumption of foods rich in carbohydrate that result in postprandial elevation in blood glucose cause repetitive acute inflammation that can worsen systemic inflammation [11]. Research suggests that weight loss reduces the levels of pro-inflammatory markers in fibromyalgia [12], low back pain [13] and osteoarthritis of the knee [4]. It has also been claimed that exercise itself has a prophylactic effect on the development of some pain conditions such as fibromyalgia syndrome, and neck, shoulder and back pain [14]. Evidence from large population cohort studies suggests that exercise reduces systemic inflammation independent of weight loss [15]. Reduced adipose tissue hypoxia from increased tissue perfusion associated with exercise may be a potential mechanism [16]. Thus, exercise has the potential to reduce low grade systemic inflammation, decrease mechanical loading and improve psychological state in addition to having other health benefits such as improved cardiovascular risk status [17].

THE CHALLENGE

There are a variety of challenges implementing exercise therapy into clinical practice including adherence, dosage and optimising exercise and recovery between treatments [18]. Adherence to regular exercise is challenging for individuals who are in pain and obese, especially as they may well be habitually sedentary. Attrition rates can be as high as 50% for adults engaging in exercise programmes so perhaps the way forward is to try and integrate more exercise into the normal daily routine. In their meta-analysis of the hypoalgesic effects of exercise, Naugle et al. were unable to determine the optimal dose of exercise needed to produce hypoalgesia [8]. Moreover, there are no guidelines for exercise prescription to reduce systemic inflammation, so advice is taken from

investigations of physical activity on body fat. Studies by Tudor-Locke et al. suggest that as little as 30 minutes walking activity per day, equating to approximately 3,000 extra steps (equivalent to approximately 30 minutes of activity), will be sufficient for improving general health including cardiovascular risk factors and weight control [19]. This can be achieved by integrating exercise into activities of daily living and therefore does not need to be prescriptive, formal or unpleasant. Several short bouts of exercise are as effective as fewer longer sessions, so increasing exercise can be as simple as encouraging less use of the car, taking the stairs more often or taking the dog for a walk for an extra 10 minutes [20]. For this to happen it is necessary to create the right social conditions that result in health for all. This would include the creation of environments supportive of exercise such as for example improved public transport, cycle lanes, pedestrian safety, and subsidised swimming and leisure facilities. Workplace exercise promotion such as cycle to work schemes or lunchtime exercise groups are also effective. Health promotion can play a major part in developing a 'social model of health' as we have discussed in a previous Editorial in this journal [21].

THE WAY FORWARD

In these stringent times we have in our pain management toolbox a potentially cost-effective way of dealing with the increasing prevalence of chronic pain whilst simultaneously helping us tackle some of the other health problems caused by obesity. Factors contributing to an increased prevalence and severity of chronic pain in individuals who are obese include systemic inflammation, mechanical overload, and autonomic dysfunction. Reductions in abdominal fat decreases circulating levels of pro-inflammatory biomarkers that may contribute to chronic pain. Further research is necessary to reveal precisely how much exercise is effective in reducing systemic inflammation and decreasing pain with or without weight loss. There is a need for evidence-based guidelines which should provide direction on appropriate types and amount of exercise required for the reduction of pain.

Nevertheless, there is already sufficient available evidence to meet the challenge head-on and start encouraging our obese patients to take more walking steps daily as a first-line treatment for chronic pain.

KEYWORDS

Chronic pain, Obesity, Exercise, Physical Activity, Systemic Inflammation, Cytokines, Pain Management

REFERENCES

1. Ng M, Fleming T, Robinson M *et al.* Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 384(9945), 766-781 (2014).
2. Bonakdar RA. Targeting systemic inflammation in patients with obesity-related pain: Obesity- related pain: time for a new approach that targets systemic inflammation. *The Journal of Family Practice* 62(9 Suppl CHPP), S22-S29 (2013).
3. Narouze S, SouzdaInitski D. Obesity and chronic pain: opportunities for better patient care. *Pain Manag* 5(4), 217-219 (2015).
4. Messier SP, Mihalko SL, Legault C *et al.* Effects of intensive diet and exercise on knee joint loads, inflammation, and clinical outcomes among overweight and obese adults with knee osteoarthritis: the IDEA randomized clinical trial. *JAMA* 310(12), 1263-1273 (2013).
5. Stiegler P, Cunliffe A. The role of diet and exercise for the maintenance of fat-free mass and resting metabolic rate during weight loss. *Sports Medicine (Auckland, N.Z.)* 36(3), 239-262 (2006).
6. Shaw K, Gennat H, O'Rourke P, Del Mar C. Exercise for overweight or obesity. *The Cochrane Database Of Systematic Reviews* (4), CD003817 (2009).
7. Runhaar J, Luijsterburg P, Dekker J, Bierma-Zeinstra SM. Identifying potential working mechanisms behind the positive effects of exercise therapy on pain and function in osteoarthritis; a systematic review. *Osteoarthritis Cartilage* 23(7), 1071-1082 (2015).
8. Naugle KM, Fillingim RB, Riley JL. A meta-analytic review of the hypoalgesic effects of exercise. *J Pain* 13(12), 1139-1150 (2012).
9. Nijs J, Kosek E, Van Oosterwijck J, Meeus M. Dysfunctional endogenous analgesia during exercise in patients with chronic pain: to exercise or not to exercise? *Pain Physician* 15(3 Suppl), ES205-213 (2012).
10. Mcvinnie DS. Obesity and pain. *British Journal of Pain* 7(4), 163-170 (2013).
11. Esposito K, Nappo F, Marfella R *et al.* Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans - Role of oxidative stress. *Circulation* 106(16), 2067-2072 (2002).
12. Senna MK, Sallam Ra-ER, Ashour HS, Elarman M. Effect of weight reduction on the quality of life in obese patients with fibromyalgia syndrome: a randomized controlled trial. *Clinical Rheumatology* 31(11), 1591-1597 (2012).
13. Roffey DM, Ashdown LC, Dornan HD *et al.* Pilot evaluation of a multidisciplinary, medically supervised, nonsurgical weight loss program on the severity of low back pain in obese adults. *The Spine Journal: Official Journal of The North American Spine Society* 11(3), 197-204 (2011).
14. Mork PJ, Vasseljen O, Nilsen TIL. Association between physical exercise, body mass index, and risk of fibromyalgia: longitudinal data from the Norwegian Nord-Trøndelag Health Study. *Arthritis Care & Research* 62(5), 611-617 (2010).
15. Bergstrom G, Behre CJ, Schmidt C. Moderate intensities of leisure-time physical activity are associated with lower levels of high-sensitivity C-reactive protein in healthy middle-aged men. *Angiology* 63(6), 412-415 (2012).
16. You T, Arsenis NC, Disanzo BL, Lamonte MJ. Effects of exercise training on chronic inflammation in obesity : current evidence and potential mechanisms. *Sports Medicine (Auckland, N.Z.)* 43(4), 243-256 (2013).
17. Zdziarski LA, Wasser JG, Vincent HK. Chronic pain management in the obese patient: a focused review of key challenges and potential exercise solutions. *J Pain Res* 8 63-77 (2015).
18. Daenen L, Varkey E, Kellmann M, Nijs J. Exercise, not to exercise, or how to exercise in patients with chronic pain? Applying science to practice. *Clin J Pain* 31(2), 108-114 (2015).
19. Tudor-Locke C, Craig CL, Brown WJ *et al.* How many steps/day are enough? For adults. *The International Journal Of Behavioral Nutrition And Physical Activity* 8 79-79 (2011).

20. Marshall SJ, Levy SS, Tudor-Locke CE *et al.* Translating Physical Activity Recommendations into a Pedometer-Based Step Goal: 3000 Steps in 30 Minutes. *American Journal Of Preventive Medicine* 36(5), 410-415 (2009).
21. Johnson MI, Briggs M, Dixey R. Should health promotion be on the pain agenda? *Pain Manag* 4(6), 385-388 (2014).