



LEEDS
BECKETT
UNIVERSITY

Citation:

Boos, C and O'Hara, J and Mellor, A and Hodkinson, PD and Tsakirides, C and Reeve, N and Gallagher, L and Green, ND and Woods, DR (2016) A Four-Way Comparison of Cardiac Function with Normobaric Normoxia, Normobaric Hypoxia, Hypobaric Hypoxia and Genuine High Altitude. PLoS One, 11 (4). -. ISSN 1932-6203 DOI: <https://doi.org/10.1371/journal.pone.0152868>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/2595/>

Document Version:

Article (Published Version)

Creative Commons: Attribution 3.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

RESEARCH ARTICLE

A Four-Way Comparison of Cardiac Function with Normobaric Normoxia, Normobaric Hypoxia, Hypobaric Hypoxia and Genuine High Altitude

Christopher John Boos^{1,2,3*}, John Paul O'Hara³, Adrian Mellor^{3,4,5}, Peter David Hodgkinson^{5,6,7}, Costas Tsakirides³, Nicola Reeve³, Liam Gallagher³, Nicholas Donald Charles Green^{5,6}, David Richard Woods^{3,5,8,9}

1 Department of Cardiology, Poole Hospital NHS Foundation trust, Poole, United Kingdom, **2** Dept of Postgraduate Medical Education, Bournemouth University, Bournemouth, United Kingdom, **3** Research Institute, for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, United Kingdom, **4** James Cook University Hospital, Middlesbrough, TS4 3BW, United Kingdom, **5** Defence Medical Services, Lichfield, United Kingdom, **6** RAF Centre of Aviation Medicine, RAF Henlow, Beds, SG16 6DN, United Kingdom, **7** Division of Anaesthesia, University of Cambridge, Box 93, Addenbrooke's Hospital, Hills Road, Cambridge CB2 2QQ, United Kingdom, **8** Northumbria and Newcastle NHS Trusts, Wansbeck General and Royal Victoria Infirmary, Newcastle, United Kingdom, **9** University of Newcastle, Newcastle upon Tyne, United Kingdom

* christopherboos@hotmail.com



OPEN ACCESS

Citation: Boos CJ, O'Hara JP, Mellor A, Hodgkinson PD, Tsakirides C, Reeve N, et al. (2016) A Four-Way Comparison of Cardiac Function with Normobaric Normoxia, Normobaric Hypoxia, Hypobaric Hypoxia and Genuine High Altitude. PLoS ONE 11(4): e0152868. doi:10.1371/journal.pone.0152868

Editor: Christopher Torrens, University of Southampton, UNITED KINGDOM

Received: October 22, 2015

Accepted: March 21, 2016

Published: April 21, 2016

Copyright: © 2016 Boos et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Research reported in this study was supported by the Surgeon General, UK and the Drummond Foundation and Leeds Beckett University. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Defence Medical Services.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Background

There has been considerable debate as to whether different modalities of simulated hypoxia induce similar cardiac responses.

Materials and Methods

This was a prospective observational study of 14 healthy subjects aged 22–35 years. Echocardiography was performed at rest and at 15 and 120 minutes following two hours exercise under normobaric normoxia (NN) and under similar PiO₂ following genuine high altitude (GHA) at 3,375m, normobaric hypoxia (NH) and hypobaric hypoxia (HH) to simulate the equivalent hypoxic stimulus to GHA.

Results

All 14 subjects completed the experiment at GHA, 11 at NN, 12 under NH, and 6 under HH. The four groups were similar in age, sex and baseline demographics. At baseline rest right ventricular (RV) systolic pressure (RVSP, $p = 0.0002$), pulmonary vascular resistance ($p = 0.0002$) and acute mountain sickness (AMS) scores were higher and the SpO₂ lower ($p < 0.0001$) among all three hypoxic groups (GHA, NH and HH) compared with NN. At both 15 minutes and 120 minutes post exercise, AMS scores, Cardiac output, septal S', lateral S', tricuspid S' and A' velocities and RVSP were higher and SpO₂ lower with all forms of

hypoxia compared with NN. On post-test analysis, among the three hypoxia groups, SpO₂ was lower at baseline and 15 minutes post exercise with GHA (89.3±3.4% and 89.3±2.2%) and HH (89.0±3.1 and (89.8±5.0) compared with NH (92.9±1.7 and 93.6±2.5%). The RV Myocardial Performance (Tei) Index and RVSP were significantly higher with HH than NH at 15 and 120 minutes post exercise respectively and tricuspid A' was higher with GHA compared with NH at 15 minutes post exercise.

Conclusions

GHA, NH and HH produce similar cardiac adaptations over short duration rest despite lower SpO₂ levels with GHA and HH compared with NH. Notable differences emerge following exercise in SpO₂, RVSP and RV cardiac function.

Introduction

Hypoxic exposure has a number of important clinical applications. These include pre-acclimatization training for athletes, the investigation of high altitude (HA) illnesses such as acute mountain sickness (AMS) and clinical diseases complicated by tissue hypoxia [1,2]. In order to improve the understanding of the clinical effects of genuine HA, hypoxia has been experimentally reproduced typically using normobaric hypoxia (NH) or hypobaric hypoxia (HH). NH lowers the partial pressure of inspired oxygen (PiO₂) by reducing the fraction of inspired oxygen (FiO₂) through addition of exogenous nitrogen (N₂) without altering the barometric pressure whereas HH lowers the PiO₂ by reduction of barometric pressure [2].

There has been considerable and ongoing debate in the medical literature as to whether these differing methods of hypoxic challenge are meaningfully different or clinically important and most importantly whether they are effective surrogate for real life HA [2–8]. A large number of the comparative studies have been in animal models and the human studies have been predominantly two-way comparisons of NH with HH exposure and have not included either a genuine 'real-world' terrestrial HA (GHA) or a normobaric normoxia control (NN) group, akin to normal sea level, limiting the clinical impact of their findings [2]. It has become increasingly appreciated that the physiological responses to GHA at a given altitude are influenced by variations in the ambient barometric pressure due to the differences in latitude, time of year and prevailing weather conditions which may be an important factor in comparative studies [3,9]. Hence, it is crucial that the ambient pressure for a given field altitude is documented in to allow more reliable comparison both between GHA experiments and with HH chamber studies [3,9]. Most of the previously published studies to compare differing hypoxia modalities have been undermined by their relatively short periods of hypoxic exposure (<30 minutes), the use of a separate but matched populations for the differing hypoxic challenge groups and the use of only brief wash out periods between each exposure increasing the risk of acclimatization bias [2]. Furthermore, the important stimulus of exercise which is a crucial factor in the majority of HA ventures in real life has been frequently overlooked.

Acute hypoxia leads to a number of recognized cardiopulmonary responses, which notably includes pulmonary vasoconstriction and an associated increase in pulmonary vascular resistance [1, 10–13]. Whether HA and the associated hypoxia leads to deleterious effects on cardiac function remains still remains a controversial issue [2,8]. Published studies have consistently shown that acute hypoxia leads to an increase in resting cardiac output and preservation of long axis systolic and radial systolic function [2, 9–12]. However, more concerning, there is a

blunted stroke volume response and variable effects on left ventricular diastolic filling and right ventricular systolic function have been observed [9,11,13]. Right ventricular (RV) diastolic function has been barely explored [2,10]. Evidence to suggest the potential deleterious effects of HA on cardiac performance include the observed increase in brain natriuretic peptide (BNP) levels at HA compared with sea level and their link to AMS and its severity [10,14,15]. It has also been shown that sustained hypoxia can lead to a decline in cardiac energetics, which is linked to adverse changes in left ventricular diastolic function despite preservation of systolic function [16].

There have been only two studies to date that have tried to compare potential changes in cardiac function during exercise following differing modes of hypoxic challenge and in both echocardiographic assessments of biventricular performance and/or right ventricular systolic pressure were not assessed [17,18]. A four-way comparison of NN, genuine HA (GHA), NH and HH on cardiac function has never been performed. Furthermore, none of the cross comparison studies to date utilized recent advances in echocardiography allowing much detailed assessment of biventricular systolic and diastolic function. Consequently, in this study we sought to investigate, for the first time, the effects of acute and sustained hypoxia on cardiac function at rest and following exercise under NN, NH and HH and GHA.

Materials and Methods

Study population

This was a prospective observational study of 14 healthy British military servicemen aged 22–35 years. In addition to completing a detailed health questionnaire all subjects were required to have a normal baseline ECG and echocardiogram to confirm suitability for inclusion. Baseline health status was undertaken via a history, clinical examination, bloods, electrocardiogram and transthoracic echocardiogram.

Study protocol

All participants completed a standard maximal incremental cycle test to volitional exhaustion at sea level (absolute altitude ~113m) under normobaric normoxia (NN) to determine maximal oxygen uptake and maximal workload (W_{max} [watts]) [19]. This was followed by a maximal incremental test to volitional exhaustion >24 hours later under NH (an FiO_2 equivalent to 3375m/11078ft (PiO_2 ~95 mmHg) in order to establish and ensure equivalent workloads for the hypoxic experimental trials [19].

Participants were then required to complete physiological assessments prior to and during exercise and rest under four different conditions. They were then assessed at GHA (3375m/11078ft, 'real' altitude, barometric pressure 506.4 ± 1.7 mmHg), followed in order with assessments at NN, NH (TISS, Alton, UK and Sporting Edge, Sherfield on Loddon, UK) and HH (Centre for Aviation Medicine, RAF Henlow, Henlow, UK) ensuring a minimum washout period of >7 days between each experimental condition. This sequence ensured the PiO_2 experienced breathing ambient air during GHA ($PiO_2 = 96.3 \pm 0.4$ mmHg) could be replicated for each individual during subsequent NH and HH exposures.

For NH the FiO_2 ($13.9 \pm 0.2\%$) was manipulated to equate to each individual's PiO_2 established at terrestrial GHA using the following equation, which considers fluctuations in sea level barometric pressure [3,8]:

$$FiO_2 = PiO_2 \text{ (mmHg)} / \text{local PB (mmHg)} - PH_{20} \text{ (47mmHg)}$$

For the measurements in the HH chamber participants underwent a decompression period (0.33 mmHg s^{-1} equivalent to 5 m.s^{-1} (ascent rate) ~ 10 minutes) to the target altitude, recreating each participants PiO_2 , and on completion of the exposure a recompression period (0.33 mmHg s^{-1}) to the ambient pressure. The chamber was continuously flushed with medical quality gas to maintain the inspired fractions of O_2 and CO_2 at 20.9% and 0.03%, respectively, with nitrogen balance. During the HH exposure participants were continuously monitored by a chief medical officer, who was present in the chamber breathing through an O_2 diluter demand mask. They were in constant contact with the chamber operators and additional medical staff.

The GHA challenge involved rapid ascent by cable car to 3375m after the subjects were driven in a minibus from sea level to 1400m. The NN and NH experiments were undertaken within a NH chamber at Leeds Beckett University and the HH chamber at the Royal Air Force Centre of Aviation Medicine, Henlow.

Each experiment was performed following a 12-hour overnight fast. All subjects underwent 30 minutes of altitude acclimatization followed by a complete 120 minutes of cycling exercise (a progressive intensity warm-up for 15 minutes, followed by 105 minutes at 55% W_{max} based on the NH maximal exercise test). All exercising testing occurred on a bicycle affixed to a bicycle trainer (Compu Trainer Pro Lab, Racer Mate, USA). The cycle ergometer was calibrated following the manufacturer's instructions. The load generator ensured the relative workloads between conditions for each participant were accurately maintained, taking into consideration an individual's natural torque fluctuations with each pedal stroke. The manufacturer's reports an accuracy of 2.5% and repeatability of 1%. Each experimental trial involved the ingestion of a carbohydrate solution (glucose-fructose) so that the dietary intake of each participant was standardised across all four studies. Physiological measurement of AMS scores, haemodynamics and cardiac function were assessed at rest, at 15 minutes into the rested acclimatization process in each study condition and then again rested 15 minutes and 120 minutes post two hours of cycling exercise in the hypoxic environment. A consistent temperature range of 18–23°C was maintained for all four study conditions.

Ethics

The study was approved by the Ministry of Defence Research and Medical Ethics Committee and was conducted according to the standards of the declaration of Helsinki and all subjects underwent written informed consent.

Physiological measurements

Resting recordings of oxygen saturations (SpO_2) were performed using a Nellcor N-20P pulse oximeter (Nellcor Puritan Bennett, Coventry, UK) following a 15 second continuous recording using the index finger of the right hand with the most consistent reading being used. Blood pressure were measured using an automated blood pressure cuff with the subject sat upright for >10 minutes at rest M6 (Omron Healthcare, Milton Keynes, UK) and heart rate was measured from a single lead ECG at the time of the echocardiogram. The ambient temperature was recorded for each experimental condition (NN.NH, HH and GHA) using a PCE-THB 40 Barometer (PCE Instruments UK Ltd).

Acute mountain sickness (AMS) scores

HA related symptoms were assessed using the Lake Louis Scoring System (LLS) [20]. The LLS score allocates a score of 0–3 (symptom not present to severe) for symptoms of AMS (headache, gastrointestinal symptoms, fatigue/weakness, dizzy/light-headedness, difficulty sleeping).

A total score of ≥ 3 in the presence of a headache is consistent with AMS and ≥ 6 with severe AMS [15,20].

Echocardiographic assessment

All echocardiographic assessments were undertaken using a portable Vivid I echocardiogram machine (GE Healthcare™, Amersham, Bucks, UK) with a 1.5–3.6 MHz S4 transducer. Pulsed-wave and two dimensional colour images were acquired in the parasternal short axis and apical four-chamber view during a short end-expiration pause with the subject lying in the left lateral position. RVSP was estimated from the maximum velocity of the trans-tricuspid gradient using continuous wave Doppler imaging [16,21]. The pulsed-wave sample volume of the conventional Doppler was placed at the tips of the mitral and tricuspid valve leaflets in order to measure the peak early transvalvular flow velocity (E), and the peak flow velocity (A) of late diastolic filling and the E/A ratios [22]. Pulsed-wave TDI volume samples were recorded at the septal and lateral mitral annulus and over the RV free wall to assess early and late diastolic filling due to left ventricular relaxation (E') and atrial contraction (A') and long axis systolic function (S') (10–12,22). The pulmonary artery vascular resistance (PVR) was calculated using the following equation $PVR = 80 \times TRV/VTI_{RVOT}$ where TRV was the maximal tricuspid regurgitation velocity and velocity time integral of the RV outflow tract velocity measured using pulsed wave doppler at the level of the pulmonary valve in the parasternal short axis view as previously described [10].

Pulsed-wave TDI was used to quantify the respective left and right ventricular isovolumic contraction (ICT) and isovolumic relaxation times (IRT) and the isovolumic contractile velocities (ICV) [3]. Right and left ventricular myocardial performance (Tei) indices (IRT+ICT/ejection time) were performed using TDI [10, 23]. Tricuspid annular plane systolic excursion (TAPSE) was recorded using M-Mode as previously described [24]. The isovolumic contractile velocity was measured at the tricuspid and mitral annulus using PWTDI [21,22]. Stroke volume and cardiac output were calculated using the aortic systolic flow velocity integral, using pulsed-wave profile of aortic blood flow from the apical five chamber view and the cross sectional area of the Left ventricular outflow tract [11,12]

Statistical Methods and power calculations

Data were analysed using SPSS® statistics version 22. The Kolmogorov-Smirnov test and inspection of the data was undertaken to assess normality of all continuous data. All data are presented as mean \pm standard deviations. Between group comparisons of categorical data for three or more groups were compared using the Fisher's Exact Test. Continuous data across the four experimental altitude groups (NN, HA, NH and HH) were assessed using Ordinary ANOVA with Bonferroni post-test for parametric data and with Kruskal-Wallis and Dunn Post-test for non-parametric data when the P value was < 0.05 . Time dependent changes (rest, 15 and 120 minutes post exercise) of continuous data within each group were assessed using Repeated measures ANOVA with Bonferroni post-test for parametric data and using Friedman Test with Dunn Post-test for non-parametric data. Correlation was assessed using Spearman Rank correlation with the 95% confidence interval of R. Further exploratory analyses of the three hypoxia groups only were undertaken using a Two-Way split level 3x3 Repeated Measures ANOVA. The within-subjects main effect of time (before and 15 and 120 min after exercise) and the between-subject main effects mode of hypoxia (GHA, NH and HH) with Bonferroni post-tests and their interactions (and effect size, Eta [n^2]) were assessed. A two tailed P value < 0.05 was considered statistically significant for all comparisons.

Sample size calculations were based on previous studies. In 11 out of the 13 prior comparative experimental hypoxia studies the sample size has been between 7–12 subjects [2]. In another very recent comparative study of six subjects Beidleman et al observed that cycling time trial performance was impaired to a greater degree in HH versus NH at the same ambient PO₂ equivalent to 4,300 m despite similar cardiorespiratory responses [17]. Hence, based on this previously published work it was calculated that a sample size of ≥12 subjects would be sufficient to detect a significant difference in cardiac performance and allow for a minimum group sample size of 6 subjects in the event of any drop outs given the intense and prolonged nature of these four group comparisons.

Results

Fourteen subjects completed the genuine HA phase, 11 the sea level study, 12 with NH and 6 under HH. All subjects completed the exercise task in each group. Non-completion was mainly due to inter-current illness and in the case of HH failure to clear their ears or voluntary withdrawal. There were no significant differences in any of the baseline demographics across the four groups with similar ages, sex, height, and body weight and body mass indices (Table 1; S1 and S2 Tables).

Physiological and haemodynamic indices

The mean peak oxygen consumption (VO₂) at baseline with NN was 46.3±5.7mls/kg/minute. There was no difference in the mean peak VO₂ among the subjects who were subsequently included in the GHA, NH and HH groups respectively (46.7±5.7, 46.2±5.2 and 47.2±4.7 mls/kg/minute; P = 0.98). As expected peak VO₂ was significantly lower under NH versus NN (38.9± vs 46.3±5.7 mls/kg/minute; P = 0.007).

The ambient temperature was marginally but significantly higher with HH versus NH and HH (Table 1). There was a significant reduction in SpO₂ across all three hypoxia environments (GHA, NH and HH) compared with NN (P<0.001), which was sustained at all three time points (rest, 15 minutes and 120 minutes post exercise) (Table 2). On Post-test analysis SpO₂ was higher at baseline and at 15 minutes post exercise with GHA (89.3±3.4% and 89.3±2.2%) and HH (89.0±3.1 and (89.8±5.0) compared with NH (92.9±1.7 and 93.6±2.5%) (Table 2). AMS scores were higher at all time points in the three hypoxic conditions versus NN with no between group differences among the hypoxia groups. Absolute resting heart rates were higher

Table 1. Baseline demographics.

Variable	NN	GHA	NH	HH	P value
Number	11	14	12	6	
Age, years	26.4 ± 4.0	25.9 ± 3.8	26.1 ± 4.1	26.3 ± 3.8	0.96
Range	22–35	21–35	21–35	22–33	
Males (%)	7 (64%)	8 (57%)	8 (67%)	4 (67%)	0.99
Height, m	175.4 ± 9.7	174.4 ± 9.6	175.5 ± 10.0	178.3 ± 9.7	0.88
Weight, kg	72.5 ± 8.7	71.5 ± 9.9	72.9 ± 10.1	72.8 ± 11.4	0.99
BMI, kg/m ²	23.5 ± 1.9	23.4 ± 1.90	23.6 ± 2.0	22.8 ± 1.8	0.90
Current smokers n,%	6 (54.5%)	7 (50%)	5 (35.7%)	2 (33.3%)	0.89
Blood haemoglobin g/dL	14.5 ± 1.6	14.5 ± 1.7	14.7 ± 1.5	15.2 ± 1.4	0.80
Temperature, °C	20.8±1.7	19.1±1.4	19.3±0.5	22.1±2.0	0.008a,b

NN, normobaric hypoxia; GHA, genuine high altitude; NH, normobaric hypoxia; HH, hypobaric hypoxia
 Post-test differences: a, NH vs HH; b, GHA vs HH

doi:10.1371/journal.pone.0152868.t001

Table 2. Changes in acute mountain sickness (AMS) and haemodynamic variables.

Variable	NN	GHA	NH	HH	P Value
Lake Louise Scores					
-Rest	0	1 ± 1.0	1.4 ± 2.3	0.7 ± 1.2	0.02 ^{ab}
-15 minutes post exercise	0	2.7 ± 3.0	2.6 ± 2.5	2.7 ± 3.4	0.004 ^{ab}
-2 hours post exercise	0	2.3 ± 2.2	2.1 ± 2.1	2.7 ± 3.0	0.004 ^{ab}
Heart Rate, minute ⁻¹					
-Rest	60.1 ± 9.9	72.1 ± 9.0	65.7 ± 10.6	65.7 ± 7.5	0.02 ^a
-15 minutes post exercise	79.9 ± 11.1*	89.7 ± 8.6*	86.6 ± 9.9*	88.7 ± 8.7*	0.09
-2 hours post exercise	67.5 ± 9.4**	89.4 ± 10.0**	76.6 ± 15.3**	80.2 ± 11.6**	0.0005 ^{a,d}
Oxygen Saturations, %					
-Rest	98.6 ± 1.4	89.3 ± 3.4	92.9 ± 1.7	89.0 ± 3.1	<0.001 ^{abcd}
-15 minutes post exercise	98.4 ± 1.4	89.3 ± 2.2	93.6 ± 2.5	89.8 ± 5.0	<0.01 ^{abcd}
-2 hours post exercise	98.9 ± 1.5	91.6 ± 3.1**	92.8 ± 4.3	91.8 ± 5.2	<0.01 ^{abc}

NN, normobaric normoxia; GHA, genuine high altitude; NH, normobaric hypoxia; HH, hypobaric hypoxia; LLS, Lake Louis Scores; AMS-C, Acute Mountain Sickness Scores; Post-test differences: a, NN vs GHA; b, NN vs NH; c, NN vs HH; d, GHA vs NH; e, GHA vs HH; f, NH vs HH

* rest vs 15 minutes post exercise

** rest vs 2 hours post exercise

doi:10.1371/journal.pone.0152868.t002

across all three hypoxic group time points compared with NN (Table 2). At 120 minutes post exercise, resting heart rates were significantly greater with GHA than NH.

Echo parameters of left ventricular function

There were no differences in any of the echo parameters of left ventricular systolic or diastolic function at rest across the four groups. However, at 15 minutes post exercise Cardiac output ($p = 0.01$), septal S' ($p = 0.02$) and lateral S' ($p = 0.03$) and septal A' ($p = 0.003$) velocities were higher with all forms of hypoxia compared with NN with no differences between the hypoxic groups (Table 3). At 120 minutes post exercise cardiac output ($p = 0.006$), septal S' ($p = 0.04$), mitral A ($p = 0.009$) and lateral S' ($p = 0.02$) velocities were higher with acute hypoxia versus NN, with no intergroup differences among the three hypoxia groups (Table 3; Fig 1). Exercise led to an increase in the post exercise left ventricular myocardial performance (Tei) index across all four groups, which was significant at 15 minutes post exercise in the NN, NH and HH groups and only in the HH at 120 minutes post exercise compared with baseline rest (Table 3). Compared with baseline, stroke volume fell at 15 minutes post exercise under all four experimental conditions before increasing again in all except the HH group where the fall in stroke volume was sustained.

Echo parameters of right ventricular function

Resting RVSP ($p = 0.0002$), PVR ($p = 0.0002$) and tricuspid A velocities ($p = 0.01$) were higher with all forms of hypoxia compared with NN with no between group differences among the three hypoxia groups (Table 4). Compared with NN at 15 minutes post exercise RVSP ($p = 0.04$), Tricuspid S' ($p = 0.009$) and A' ($p = 0.0001$) velocities and the right ventricular Tei Index (0.001) were greater with hypoxia. The tricuspid A' velocity was significantly higher with GHA and the Tei index and RVSP with HH on post-test analysis. At 120 minutes post exercise the RVSP ($p = 0.0006$), Tricuspid A ($p = 0.006$), tricuspid S' ($p = 0.03$) and A' ($p = 0.0007$) velocities and the RV Tei index ($p = 0.004$) were higher with hypoxia than NN (Figs 2 and 3). The rise in tricuspid A' velocity, RV Tei index and RVSP were greatest among the GHA and

Table 3. Changes in Echo derived markers of Left ventricular function.

Variable	NN	GHA	NH	HH	P Value
Stroke volume, ml					
-Rest	71.5 ± 11.9	74.1 ± 11.7	73.6 ± 13.6	74.7 ± 8.4	0.85
-15 minutes post exercise	60.6 ± 8.2 *	62.7 ± 8.5*	66.5 ± 11.5 *	69.8 ± 8.5	0.16
-2 hours post exercise	66.1 ± 10.2	66.6 ± 9.3**	70.9 ± 12.6	69.7 ± 9.5	0.58
Cardiac output, L/minute					
-Rest	4.3 ± 0.8	5.3 ± 1.1	4.9 ± 1.3	4.9 ± 1.0	0.24
-15 minutes post exercise	4.8 ± 0.8	5.6 ± 1.0	5.9 ± 1.1*	6.2 ± 0.9*	0.01 ^{b,c}
-2 hours post exercise	4.4 ± 0.5	5.9 ± 1.1	5.4 ± 1.2	5.6 ± 0.9	0.006 ^a
Mitral E velocity, cm/s					
-Rest	88.6 ± 9.6	90.1 ± 12.5	92.5 ± 18.2	95.2 ± 11.7	0.78
-15 minutes post exercise	72.0 ± 16.2*	81.8 ± 13.2	77.4 ± 15.2*	84.5 ± 18.5	0.19
-2 hours post exercise	76.5 ± 15.0	85.4 ± 14.0	79.8 ± 13.3**	76.7 ± 17.6 **	0.48
Mitral A velocity, cm/s					
-Rest	48.5 ± 14.4	51.8 ± 9.2	55.2 ± 11.0	58.2 ± 11.4	0.39
-15 minutes post exercise	56.7 ± 9.5	63.3 ± 17.3	68.5 ± 12.1*	64.3 ± 11.0	0.23
-2 hours post exercise	46.6 ± 12.4	59.9 ± 14.8	67.8 ± 16.7 **	62.0 ± 10.1	0.009 ^b
Mitral E/A ratio					
-Rest	1.71 ± 0.27	1.79 ± 0.42	1.71 ± 0.37	1.67 ± 0.26	0.87
-15 minutes post exercise	1.08 ± 0.25*	1.39 ± 0.45*	1.15 ± 0.26*	1.35 ± 0.36	0.13
-2 hours post exercise	1.25 ± 0.42**	1.53 ± 0.55	1.26 ± 0.42 **	1.27 ± 0.24*	0.32
Septal S' velocity, cm/s					
-Rest	8.9 ± 1.4	10.2 ± 1.4	9.8 ± 1.6	9.0 ± 0.9	0.33
-15 minutes post exercise	8.3 ± 1.2	10.1 ± 1.7	9.4 ± 1.2	9.8 ± 0.8	0.02 ^a
-2 hours post exercise	8.9 ± 1.4	10.9 ± 1.7	9.8 ± 1.2	9.7 ± 1.2	0.04 ^a
Septal E' velocity, cm/s					
-Rest	13.0 ± 1.6	12.9 ± 1.9	13.3 ± 3.3	13.2 ± 1.5	0.97
-15 minutes post exercise	10.8 ± 1.5*	12.9 ± 1.9	12.7 ± 3.0	12.7 ± 1.5	0.11
-2 hours post exercise	11.6 ± 1.6**	12.7 ± 2.5	12.0 ± 2.0	11.0 ± 1.8**	0.36
Septal A' velocity, cm/s					
-Rest	7.9 ± 2.3	8.6 ± 2.2	9.6 ± 2.2	8.3 ± 2.0	0.09
-15 minutes post exercise	8.2 ± 1.2	11.2 ± 3.4*	11.1 ± 2.2*	10.8 ± 2.3*	0.003 ^{abc}
-2 hours post exercise	8.5 ± 1.8	10.7 ± 3.1**	10.9 ± 2.6	10.0 ± 1.1	0.11
Septal ICV, cm/s					
-Rest	6.5 ± 1.8	7.1 ± 1.6	7.0 ± 2.1	6.8 ± 1.3	0.80
-15 minutes post exercise	7.1 ± 2.1	8.4 ± 2.9	7.2 ± 2.2	6.8 ± 2.4	0.55
-2 hours post exercise	7.2 ± 1.9	8.8 ± 3.2	7.3 ± 2.1	7.7 ± 2.2	0.46
Lateral S' velocity, cm/s					
-Rest	11.1 ± 2.2	12.1 ± 2.5	12.6 ± 2.4	10.7 ± 1.7	0.06
-15 minutes post exercise	10.9 ± 1.4	13.8 ± 2.8	12.1 ± 2.4	13.5 ± 2.2*	0.03 ^a
-2 hours post exercise	12.1 ± 2.3	15.1 ± 2.9**	12.9 ± 2.3	12.8 ± 1.6**	0.02 ^a
Lateral E' velocity cm/s					
-Rest	18.3 ± 3.1	18.6 ± 2.1	20.8 ± 2.8	17.8 ± 2.8	0.90
-15 minutes post exercise	16.5 ± 2.8	17.9 ± 3.2	17.2 ± 2.5*	15.5 ± 1.6	0.52
-2 hours post exercise	17.4 ± 2.7	18.0 ± 3.8	18.1 ± 3.2**	16.0 ± 2.3	0.57
Lateral A' velocity cm/s					
-Rest	7.9 ± 2.5	9.4 ± 2.8	9.0 ± 2.6	7.7 ± 2.2	0.30
-15 minutes post exercise	8.4 ± 2.3	10.6 ± 2.4	10.1 ± 2.5	10.5 ± 2.2	0.13

(Continued)

Table 3. (Continued)

Variable	NN	GHA	NH	HH	P Value
-2 hours post exercise	8.9 ± 2.2	10.7 ± 3.5	9.9 ± 3.2	9.8 ± 2.9	0.54
Lateral ICV, cm/s					
-Rest	8.1 ± 2.0	9.1 ± 2.8	7.6 ± 2.0	7.7 ± 1.6	0.34
-15 minutes post exercise	8.6 ± 1.8	9.6 ± 3.0	9.2 ± 2.5	10.7 ± 3.1*	0.52
-2 hours post exercise	9.2 ± 1.8	8.9 ± 3.6	9.5 ± 2.5**	8.8 ± 1.8	0.94
Left ventricular Tei Index					
-Rest	0.28 ± 0.04	0.30 ± 0.03	0.29 ± 0.04	0.29 ± 0.03	0.59
-15 minutes post exercise	0.33 ± 0.04*	0.32 ± 0.04	0.33 ± 0.03*	0.37 ± 0.06*	0.15
-2 hours post exercise	0.32 ± 0.04	0.33 ± 0.03	0.33 ± 0.03	0.36 ± 0.02**	0.13

NN, normobaric normoxia; GHA, genuine high altitude; NH, normobaric hypoxia; HH, hypobaric hypoxia; ICV, isovolumic contractile velocity; Between group post-test differences: a, NN vs GHA; b, NN vs NH; c, NN vs HH; Within groups repeated measures post-test difference

* rest vs 15 minutes post exercise

** rest vs 2 hours post exercise

doi:10.1371/journal.pone.0152868.t003

HH groups respectively on post-test analysis (Table 4). There was a trend for higher PVR with HH than with either NH or GHA at all three sampling time points ($p < 0.05$ for trend). There was a significant correlation between pulmonary vascular resistance and the RV Tei index ($r = 0.54$; 95% CI 0.09–0.81).




Fig 1. Changes in the left ventricular lateral S' velocities (marginal means) with differing experimental conditions and duration of hypoxia (time 1 = baseline rest, time 2 = 15 minutes post exercise and time 3 = 120 minutes post exercise). * demonstrates between group differences on post test.

doi:10.1371/journal.pone.0152868.g001

Table 4. Changes in Echo derived markers of right ventricular function.

	Sea Level	GHA	NH	HH	P Value
Right ventricular systolic pressure, mmHg					
-Rest	21.8 ± 4.0	31.8 ± 3.7	31.5 ± 5.8	31.7 ± 4.0	0.0002 ^{abc}
-15 minutes post exercise	27.6 ± 5.2*	31.9 ± 6.1	31.4 ± 5.7	35.2 ± 5.7	0.04 ^c
-2 hours post exercise	28.2 ± 4.0**	32.0 ± 6.4	32.0 ± 5.6	36.1 ± 8.1	0.0006 ^{cf}
Pulmonary vascular resistance dynes/cm ⁵					
-Rest	86.5 ± 13.8	107.5 ± 12.2	106.1 ± 15.9	118.7 ± 15.1	0.0002 ^{a,b,c}
-15 minutes post exercise	111.5 ± 11.3*	116.8 ± 20.1	118.8 ± 19.3*	127.3 ± 13.8	0.11
-2 hours post exercise	106.7 ± 12.8**	114.5 ± 22.3	117.2 ± 15.6	129.1 ± 12.2	0.04 ^c
TAPSE, cm					
-Rest	2.6 ± 0.4	2.2 ± 0.3	2.4 ± 0.3	2.6 ± 0.3	0.05
-15 minutes post exercise	2.3 ± 0.4*	2.3 ± 0.3	2.4 ± 0.3	2.5 ± 0.3	0.63
-2 hours post exercise	2.3 ± 0.3**	2.5 ± 0.3**	2.4 ± 0.3	2.6 ± 0.3	0.27
Tricuspid E velocity cm/s					
-Rest	56.2 ± 8.3	63.4 ± 18.1	62.2 ± 12.9	66.2 ± 11.5	0.43
-15 minutes post exercise	50.4 ± 9.1	58.2 ± 12.5	50.8 ± 9.6*	57.7 ± 9.0	0.16
-2 hours post exercise	53.9 ± 10.2	58.5 ± 14.0	58.4 ± 10.5	54.3 ± 10.6 **	0.67
Tricuspid A velocity cm/s					
-Rest	33.7 ± 5.2	37.6 ± 1.0	45.9 ± 9.6	43.3 ± 10.0	0.01 ^b
-15 minutes post exercise	39.9 ± 9.0*	45.8 ± 12.3*	51.5 ± 13.1	45.2 ± 13.7	0.16
-2 hours post exercise	35.4 ± 8.4	40.4 ± 7.4	51.7 ± 16.0	40.2 ± 7.1	0.006 ^b
Right ventricular S ¹ velocity, cm/s					
-Rest	14.7 ± 1.3	15.5 ± 1.1	15.4 ± 1.6	14.5 ± 2.8	0.45
-15 minutes post exercise	13.2 ± 1.7	15.4 ± 1.5	13.4 ± 1.8*	15.2 ± 1.9	0.009 ^a
-2 hours post exercise	13.9 ± 1.9	16.4 ± 2.0	14.3 ± 2.3	14.8 ± 2.9	0.03 ^a
Right ventricular E ¹ velocity, cm/s					
-Rest	14.2 ± 2.9	16.4 ± 3.5	14.8 ± 2.8	15.7 ± 2.9	0.30
-15 minutes post exercise	13.3 ± 3.8	16.6 ± 3.8	15.3 ± 4.7	15.3 ± 4.5	0.23
-2 hours post exercise	13.7 ± 3.0	16.4 ± 2.7	15.4 ± 3.3	15.2 ± 5.5	0.23
Right ventricular A ¹ velocity, cm/s					
-Rest	11.1 ± 1.9	11.5 ± 2.1	10.2 ± 1.9	12.2 ± 1.5	0.16
-15 minutes post exercise	10.9 ± 2.5	17.5 ± 3.1*	13.2 ± 4.1*	14.8 ± 1.5	0.0001 ^{ad}
-2 hours post exercise	12.5 ± 2.4	18.1 ± 3.7**	14.9 ± 3.3*	14.8 ± 1.3**	0.0007 ^a
Right ventricular ICV velocity, cm/s					
-Rest	9.5 ± 2.7	10.5 ± 2.3	8.9 ± 1.9	9.1 ± 1.6	0.43
-15 minutes post exercise	10.8 ± 2.7	12.5 ± 3.1	10.2 ± 2.7	11.7 ± 2.1*	0.14
-2 hours post exercise	11.3 ± 3.1	12.2 ± 3.3	10.6 ± 1.9	9.7 ± 1.8	0.24
Right ventricular Tei Index					
-Rest	0.29 ± 0.05	0.32 ± 0.04	0.29 ± 0.03	0.30 ± 0.06	0.29
-15 minutes post exercise	0.33 ± 0.04	0.37 ± 0.05*	0.33 ± 0.03	0.41 ± 0.05*	0.001 ^{cf}
-2 hours post exercise	0.32 ± 0.06	0.31 ± 0.05	0.34 ± 0.07**	0.39 ± 0.05**	0.04 ^e

NN, normobaric normoxia; GHA, genuine high altitude; NH, normobaric hypoxia; HH, hypobaric hypoxia; TAPSE, trans annular plane systolic excursion; ICV, isovolumic contractile velocity; Between group post-test differences: a, NN vs GHA; b, NN vs NH; c, NN vs HH; d, GHA vs NH; e, GHA vs HH; f, NH vs HH. Within groups repeated measures post-test difference

* rest versus 15 minutes post exercise

** rest versus 2 hours post exercise.

doi:10.1371/journal.pone.0152868.t004




Fig 2. Changes in the right ventricular S' velocities (marginal means) with differing experimental conditions and duration of hypoxia (time 1 = baseline rest, time 2 = 15 minutes post exercise and time 3 = 120 minutes post exercise). * demonstrates between group differences on post test.

doi:10.1371/journal.pone.0152868.g002

Two-Way Repeated Measures ANOVA comparing the three hypoxia groups

The main effects of hypoxia and time and the potential interactions (mode of hypoxia x time) and post tests are shown in [Table 5](#) for the three hypoxia groups with NN as a reference. There was a significant main-effect for mode of hypoxia on SpO₂, Tricuspid A, A' and ICV as well as the RV Tei index ([Table 5](#)).

There was a significant main effect for time on heart rate, mitral E and A, septal E' and A', lateral S', E', A' and ICV, left ventricular Tei index and stroke volume, tricuspid E, A' and ICV and RV Tei Index. Within group comparisons of the three hypoxic groups revealed no significant interactions between exercise and time for the majority of echo parameters. However there was a significant mode of hypoxia x time interaction effect for the lateral S' ($F [2, 56] = 3.99$; $p = 0.006$; $n^2 = 0.22$), tricuspid S' ($F [2, 58] = 3.12$; $p = 0.02$; $n^2 = 0.16$) and the right ventricular Tei index ($F [2,58] = 4.1$; $p = 0.006$; $n^2 = 0.23$) (Figs [1–3](#), [Table 5](#)). The marginal means were consistently higher at 15 minutes post exercise with GHA and HH than NH (Figs [1–3](#)).

Discussion

This is the first study to assess the comparative changes in physiological and cardiac responses to exercise under four differing altitude conditions of NN, GHA, NH and HH. The key findings were that whilst all three hypoxic environments (GHA, NH and HH) led to similar cardiac adaptations at rest notable differences emerged following exercise. Compared with NH, the RVSP and RV Tei indices were higher with HH and the tricuspid A' was higher with GHA.




Fig 3. Changes in the right ventricular Tei Index (marginal means) with differing experimental conditions and duration of hypoxia (time 1 = baseline rest, time 2 = 15 minutes post exercise and time 3 = 120 minutes post exercise) * demonstrates between group differences on post test.

doi:10.1371/journal.pone.0152868.g003

The degree of hypoxemia was greater with GHA and HH than with NH at both rest and at 15 minutes post exercise. There were no significant interactions between experimental altitude and time with the exception of the RV performance (Tei index) and the RV (tricuspid) and lateral S' velocities.

There has been considerable debate in the literature as to whether differing modalities of hypoxia challenge are synonymous [2–7,25]. This is an important issue that remains unresolved and has enormous implications for HA research where inferences are often made from sea level chamber or hypoxia studies about potential responses at genuine HA. There have been only two studies, to date, that have assessed the effects of differing hypoxic environments on cardiac function. Miyagawa et al investigated seven young men who cycled for 40 minutes at 50% peak aerobic power in NN, NH and HH equivalent to 3200m in an artificial climate chamber [18]. Hence, this was a smaller sample size but similar altitude to our current study. Cardiac output and stroke volume were the only specific cardiac functional assessments performed beyond heart rate and were undertaken using pulse dye densitometry using Indocyanine Green [18]. However, both the reliability and reproducibility of this method has been challenged and its comparison to

more invasive methods of cardiac output determination have yielded conflicting results [26,27]. In their study, Miyagawa et al did not find any significant effects of experimental condition (trial) on cardiac output or stroke volume but did note there was a significant interactive effect of [trial × time] both cardiac output and stroke volume, during exercise, (suggesting that their responses to exercise were significantly different between the experimental conditions

Table 5. Results of Two-Way Repeated Measures ANOVA comparing the Main Effects of Duration (time) and mode of hypoxia across the three hypoxia groups.

	Mode of hypoxia		Time		Interaction	
	F	P value	F	P value	F	P Value
SpO ₂	6.55	0.004 ^{ac}	2.4	0.10	1.2	0.34
Heart Rate	2.32	0.12	75.47	<0.001 ^{abc}	2.18	0.08
Mitral E velocity	0.17	0.84	11.97	<0.001 ^{ac}	1.2	0.32
Mitral A velocity	1.20	0.33	5.9	0.005 ^{ac}	0.35	0.85
Mitral E/A ratio	1.93	0.16	10.79	<0.0001	0.43	0.79
Septal S' velocity	2.38	0.11	0.99	0.38	0.72	0.58
Septal E' velocity	0.20	0.82	3.37	0.04 ^b	0.77	0.55
Septal A' velocity	0.29	0.75	12.32	<0.001 ^{ab}	0.36	0.83
Septal ICV	1.0	0.40	2.70	0.08	1.0	0.42
Lateral S' velocity	1.40	0.27	10.28	<0.001 ^{ab}	3.99	0.006
Lateral E' velocity	2.20	0.13	7.56	0.001 ^{ab}	1.12	0.36
Lateral A' velocity	0.40	0.68	6.97	0.002 ^{ab}	0.58	0.68
Lateral ICV	0.12	0.89	6.99	0.002 ^a	2.14	0.09
LV Tei Index	2.32	0.12	6.48	0.003 ^a	1.54	0.10
LV stroke volume	0.39	0.68	10.37	0.004 ^{ab}	0.93	0.45
Cardiac output	0.23	0.80	13.8	<0.001 ^{ab}	2.52	0.06
RVSP	0.70	0.51	1.47	0.24	0.88	0.48
Pulmonary vascular resistance	2.10	0.14	5.85	0.005 ^{ab}	0.27	0.76
Tricuspid E velocity	0.55	0.58	5.15	0.009 ^a	0.88	0.48
Tricuspid A velocity	3.90	0.03 ^a	1.9	0.16	0.58	0.68
TAPSE	1.21	0.31	1.1	0.31	2.4	0.06
Tricuspid S' velocity	2.60	0.09	1.3	0.30	3.12	0.02
Tricuspid E' velocity	0.66	0.53	0.01	0.99	0.10	0.98
Tricuspid A' velocity	8.20	0.002 ^a	22.3	<0.001 ^{ab}	1.59	0.19
Tricuspid ICV	3.40	0.048 ^a	6.7	0.002 ^a	0.78	0.54
Right ventricular Tei Index	5.0	0.01 ^c	16.10	<0.001 ^{ab}	4.10	0.006

TAPSE, trans annular plane systolic excursion; RVSP, right ventricular systolic pressure; ICV isovolumic contractile velocity; Results of post hoc tests—time: a baseline versus 15 minutes post exercise, b baseline versus 2h post exercise, c 15 minutes versus 2 hours post exercise, altitude: a GHA vs NH, b GHA vs HH, c NH vs HH

doi:10.1371/journal.pone.0152868.t005

[18]. This related to the finding of a marked increase in cardiac output and stroke volume in both hypoxic groups versus the NN rather than any observed differences between the two hypoxic groups. Hence, as well as looking at a four-way comparison of the NN, GHA, NH and HH we undertook an additional exploratory analyses of the effects of hypoxia duration (time) and mode and their potential interactions across the three hypoxia groups. In another previous study two separate groups of six subjects were compared following approximately two hours of resting exposure to NH and HH at the equivalent of 4400m [17]. The only cardiac functional assessment performed was cardiac output, which was measured non-invasively using finger pulse waveform analysis [17]. No between group differences in cardiac output were observed however cycling time trial performance was worse with HH than NH.

In our study we assessed both right and left ventricular systolic and diastolic performance as well as markers of pulmonary artery haemodynamics (RVSP and the PVR) and global biventricular function using the Tei Index which adds significant novelty. Moreover, we ensured that

the same exercise work and duration was maintained between the groups to reduce the confounding factor of differing exercise burden on any observed results.

One of the most pertinent findings of our study is the observation that it is not only the hypoxic environment but also exercise in this environment, which influences the cardiac and pulmonary vascular responses. Hence, 'resting' comparisons do not adequately reflect the reality of exposure to HA where there is usually an exercise component. This fact is supported in this study by the interaction between experimental conditions and exercise time for the Tei index and lateral left ventricular and right ventricular (tricuspid) S' velocities (Figs 1–3). Whilst no significant between hypoxic group differences were observed over the short duration at rest, several differences emerged following exercise. For example, the RV Tei index and S' velocities and the lateral left ventricular S' velocities were consistently higher at 15 minutes post exercise with GHA and HH than NH. The higher values of RVSP, right ventricular Tei index and PVR and lower SpO₂ post exercise with HH compared with NH are particularly notable in this study. In a very recent systemic review of crossover trials of HH versus NH, Coppel et al (2015) noted that peripheral SpO₂ levels were higher with NH in two out of three short studies involving a hypoxic duration of <30 minutes with no notable differences in studies of ≥ 8 hours [2]. The authors also noted that arterial blood saturations (SaO₂) were lower with HH in all three of the previously published short-term hypoxia-duration studies [2]. Our data is consistent with this limited published literature as we observed significantly lower SpO₂ with HH than with NH. The suggested potential mechanisms to explain these observed differences, include lower minute ventilation, greater intravascular bubble formation and ventilation/perfusion mismatch, increased alveolar dead space as well as differences in alveolar fluid permeability and chemosensitivity with HH versus NH [2, 28].

Another factor, which must be considered is the effect of ambient temperature on SpO₂ readings with finger pulse oximetry. It has been shown that a significant reduction in ambient temperature leads to peripheral vasoconstriction and can lead to a small (≤1.4%) increase in SpO₂ which is thought to be explained by temperature-dependent arteriovenous shunts in the periphery [29,30]. Variations in core temperature can also affect the SaO₂ (and hence SpO₂) by rightward shift of the HbO₂ dissociation curve [31]. In our current study, we tried maintain a similar exercise intensity and ambient temperature across all four experimental conditions. However, the ambient temperature was marginally, albeit significantly higher with HH (+2–3°C) versus NH and GHA and unfortunately, we did not record core temperature. Nevertheless, previously published studies have not identified any significant differences in core temperature and thermoregulation between NH and HH [2]. In either case, we would not expect this small variation in ambient temperature across the differing experimental conditions to lead to meaningful differences in core temperature and SaO₂. Nevertheless the small temperature differences between experimental conditions is still a limitation that should be acknowledged. The fact that SpO₂ was lower and PVR and RVSP were higher with GHA and HH than for NH strengthens our findings, given the well-established reciprocal relationship between SpO₂ and RVSP/PVR, due to hypoxia driven pulmonary arterial vasoconstriction [10,20,32].

The observation of a greater increase in the RV Tei index with HH versus NH is an interesting and novel finding. The myocardial (Tei) index is a marker of global myocardial performance, which includes both systolic and diastolic functional parameters in its assessment and is independent of heart rate and ventricular geometry [23]. It may be a more sensitive marker of myocardial function than many traditional indices of cardiac function with increasing and higher values (>0.40–0.45) indicating worsening cardiac performance [23, 33]. The RV Tei index has been shown to positively correlate with both mean pulmonary artery pressure and PVR in patients with pulmonary artery hypertension and is highly susceptible to the effects of treatment [33,34]. The baseline (0.29–0.32) and post exercise increase in the RV Tei Index (up

to 0.41) in our study is consistent with the published literature [33–35]. Huez et al previously noted that the RV Tei Index increased by approximately 50% (versus up to 33% in our study) following acute but more prolonged exposure to a higher altitude of 3750 among 15 healthy Caucasian adults [36]. More recently Page et al noted that the RV Tei Index increased significantly at genuine field HA (0.32 ± 0.08 at 30 m to 0.43 ± 0.15 at 3450 m and 0.41 ± 0.10 at 4730 m; $P = 0.046$) and was associated with subclinical pulmonary in 13 out of the 14 subjects [37]. This increase in the Tei Index was nearly identical to that in our study that is also in keeping with that noted among patients with treatment responsive pulmonary hypertension (22,33). We noted an interaction between exercise and HA environment on the RV Tei. (Fig 3). It has been previously demonstrated that heavy exercise under hypoxia leads to increased capillary disruption and fluid leak, which may be one of the mechanisms in the development of high altitude pulmonary oedema [38]. We observed that the PVR positively correlated with the RV Tei index and interestingly the HH group had both the highest Tei index and increase in PVR and PASP strengthening our findings. Our data suggests that increased PVR may have a negative effect on RV function, perhaps by increasing RV afterload due to increased PASP. The RV Tei Index appears to be particularly sensitive to even short-term changes in the hypoxic environment supporting previous work [33, 34].

The interaction between time and experimental conditions on the left and right ventricular S' velocity appeared to relate to a more sustained increase in long axis function with GHA in relation to NN and NH where the changes were less marked (Figs 1 and 3). We also observed marked differences in the right ventricular systolic (S' and ICV), diastolic (E and A') and global function (Tei) depending on the hypoxic environment.

This study has a number of limitations that need acknowledgment. The sample size in the HH was smaller than among the other three groups, which could lead to selection bias and reduced power to detect a difference that was not appreciated. However, the demographics on the smaller HH group were similar to the other three groups and this sample size is at least as large as several previous comparative studies. Despite allowing for a reasonable 'washout' period between studies, it is uncertain whether the order of the studies could have influenced symptom scores and cardiac performance over time due to changes in physical fitness and experience of HA exposure. Furthermore, the effects of hyobarbia was not independently assessed in this study through a hypobaric normoxia condition, as this was not feasible, and as such, the findings of this study should be reviewed in the context in which they have been presented.

In conclusion, HH, NH and HH produce similar cardiac adaptations at rest. However, notable differences emerge following exercise in the degree of hypoxemia, RVSP, RV systolic, diastolic and global function. This was most marked with HH and GH versus NH where the post exercise RV Tei and S' velocity respectively were greater. The type of hypoxic environment and exercise performed in this environment significantly influence the cardiac response. Observed changes in cardiac function with NH are not necessarily predictive of similar changes with genuine HA or HH and vice versa.

Supporting Information

S1 Table. SPSS file with information on the three hypoxia groups and their respective echo data.

(SAV)

S2 Table. Excel file with full data for all four experimental study groups.

(XLSX)

Acknowledgments

The authors would like to thank the Drummond Foundation, Leeds Beckett University, the Centre of Aviation Medicine, RAF Henlow, the Defence Medical Services and the Surgeon General's Department for their support. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the Defence Medical Services.

Author Contributions

Conceived and designed the experiments: CJB JPO AM PDH CT NR LG NDCG DRW. Performed the experiments: CJB JPO AM PDH CT NR LG NDCG DRW. Analyzed the data: CJB. Contributed reagents/materials/analysis tools: CJB JPO AM PDH CT NR LG NDCG DRW. Wrote the paper: CJB JPO AM PDH NDCG DRW.

References

1. Bärtsch P, Swenson ER (2013). Clinical practice: Acute high-altitude illnesses. *N Engl J Med* 368: 2294–302. doi: [10.1056/NEJMcp1214870](https://doi.org/10.1056/NEJMcp1214870) PMID: [23758234](https://pubmed.ncbi.nlm.nih.gov/23758234/)
2. Coppel J, Hennis P, Gilbert-Kawai E, Grocott MP (2015). The physiological effects of hypobaric hypoxia versus normobaric hypoxia: a systematic review of crossover trials. *Extrem Physiol Med* 26: 4:1–20.
3. Conkin J (2011). PH₂O and simulated hypobaric hypoxia. *Aviat Space Environ Med* 82: 1157–8. PMID: [22195399](https://pubmed.ncbi.nlm.nih.gov/22195399/)
4. Feriche B, García-Ramos A, Calderón-Soto C, Drobnic F, Bonitch-Góngora JG, Galilea PA, et al (2014). Effect of acute exposure to moderate altitude on muscle power: hypobaric hypoxia vs. normobaric hypoxia. *PLoS One* 9: e114072. doi: [10.1371/journal.pone.0114072](https://doi.org/10.1371/journal.pone.0114072) PMID: [25474104](https://pubmed.ncbi.nlm.nih.gov/25474104/)
5. Millet GP, Faiss R, Pialoux V (2012) Point: Hypobaric hypoxia induces different physiological responses from normobaric hypoxia. *J Appl Physiol* 112: 1783–4. doi: [10.1152/jappphysiol.00067.2012](https://doi.org/10.1152/jappphysiol.00067.2012) PMID: [22267386](https://pubmed.ncbi.nlm.nih.gov/22267386/)
6. Mounier R, Brugniaux JV (2012). Counterpoint: Hypobaric hypoxia does not induce different physiological responses from normobaric hypoxia. *J Appl Physiol* 112: 1784–6. doi: [10.1152/jappphysiol.00067.2012a](https://doi.org/10.1152/jappphysiol.00067.2012a) PMID: [22589489](https://pubmed.ncbi.nlm.nih.gov/22589489/)
7. Savourey G, Launay JC, Besnard Y, Guinet A, Travers S (2003). Normo- and hypobaric hypoxia: are there any physiological differences? *Eur J Appl Physiol* 89:122–6. PMID: [12665974](https://pubmed.ncbi.nlm.nih.gov/12665974/)
8. Saugy JJ, Schmitt L, Cejuela R, Faiss R, Hauser A, Wehrin JP, et al (2014). Comparison of "Live High-Train Low" in normobaric versus hypobaric hypoxia. *PLoS One* 9:e114418 doi: [10.1371/journal.pone.0114418](https://doi.org/10.1371/journal.pone.0114418) PMID: [25517507](https://pubmed.ncbi.nlm.nih.gov/25517507/)
9. Dill DB, Evans DS (1970) Report barometric pressure. *J Appl Physiol* 29: 914–6. PMID: [5485369](https://pubmed.ncbi.nlm.nih.gov/5485369/)
10. Boos CJ, Hodkinson P, Mellor A, Green NP, Woods DR (2012). The effects of acute hypobaric hypoxia on arterial stiffness and endothelial function and its relationship to changes in pulmonary artery pressure and left ventricular diastolic function. *High Alt Med Biol* 13: 105–11. doi: [10.1089/ham.2012.1009](https://doi.org/10.1089/ham.2012.1009) PMID: [22724613](https://pubmed.ncbi.nlm.nih.gov/22724613/)
11. Boos CJ, Holdsworth DA, Woods DR, Green K, Naylor J, Mellor A (2013). Cardiac biomarkers and high altitude pulmonary edema. *Int J Cardiol* 167: e65–6. doi: [10.1016/j.ijcard.2013.03.119](https://doi.org/10.1016/j.ijcard.2013.03.119) PMID: [23632120](https://pubmed.ncbi.nlm.nih.gov/23632120/)
12. Boos CJ, Mellor A, Begley J, Stacey M, Smith C, Hawkins A, et al (2014). The effects of exercise at high altitude on high-sensitivity cardiac troponin release and associated biventricular cardiac function. *Clin Res Cardiol* 103:291–9. doi: [10.1007/s00392-013-0654-2](https://doi.org/10.1007/s00392-013-0654-2) PMID: [24363107](https://pubmed.ncbi.nlm.nih.gov/24363107/)
13. Rao M, Li J, Qin J, Zhang J, Gao X, Yu S, et al (2015). Left ventricular function during acute high-altitude exposure in a large group of healthy young Chinese men. *PLoS One* 10: e0116936. doi: [10.1371/journal.pone.0116936](https://doi.org/10.1371/journal.pone.0116936) PMID: [25629435](https://pubmed.ncbi.nlm.nih.gov/25629435/)
14. Mellor A, Boos C, Holdsworth D, Begley J, Hall D, Lumley A, et al (2014). Cardiac biomarkers at high altitude. *High Alt Med Biol* 15: 452–8. doi: [10.1089/ham.2014.1035](https://doi.org/10.1089/ham.2014.1035) PMID: [25330333](https://pubmed.ncbi.nlm.nih.gov/25330333/)
15. Woods DR, Begley J, Stacey M, Hooper T, Smith C, Begley J, et al (2012). Severe acute mountain sickness, brain natriuretic peptide and NT-proBNP in humans *Acta Physiol* 205: 349–55.
16. Holloway C, Cochlin L, Codreanu I, Bloch E, Fatemian M, Szmigielski C, et al (2011). Normobaric hypoxia impairs human cardiac energetics. *FASEB J* 25: 3130–5. doi: [10.1096/fj.11-183426](https://doi.org/10.1096/fj.11-183426) PMID: [21646398](https://pubmed.ncbi.nlm.nih.gov/21646398/)
17. Beidleman BA, Fulco CS, Staab JE, Andrew SP, Muza SR (2014). Cycling performance decrement is greater in hypobaric versus normobaric hypoxia. *Extrem Physiol Med* 3: 8. doi: [10.1186/2046-7648-3-8](https://doi.org/10.1186/2046-7648-3-8) PMID: [24778792](https://pubmed.ncbi.nlm.nih.gov/24778792/)

18. Miyagawa K, Kamijo Y, Ikegawa S, Goto M, Nose H (2011). Reduced hyperthermia-induced cutaneous vasodilation and enhanced exercise-induced plasma water loss at simulated high altitude (3,200 m) in humans. *J Appl Physiol* (1985) 110:157–65.
19. Kuipers H, Keizer HA, Brouns F, Saris WH (1987). Carbohydrate feeding and glycogen synthesis during exercise in man. *Pflugers Arch* 410:652–6. PMID: [3449801](#)
20. Hackett PH & Oelz O (1992) The Lake Louise consensus on the quantification of altitude illness. In: Sutton JR, Houston CS & Coates G (eds) *Hypoxia and Mountain Medicine*: Queen City Printers, Burlington VT: 327–330.
21. Naeije R, Huez S, Lamotte M, Retailleau K, Neupane S, Abramowicz, et al (2010). Pulmonary artery pressure limits exercise capacity at high altitude. *Eur Respir J* 36:1049–55. doi: [10.1183/09031936.00024410](#) PMID: [20378601](#)
22. Paulus WJ, Tschöpe C, Sanderson JE, Rusconi C, Flachskampf FA, Rademakers FE (2007). How to diagnose diastolic heart failure: A consensus statement on the diagnosis of heart failure with normal left ventricular ejection fraction by the Heart Failure and Echocardiography Association of the European Society of Cardiology. *Eur Heart J* 28:2539–2550. PMID: [17428822](#)
23. Tei C, Dujardin KS, Hodge DO, Bailey KR, McGoon MD, Tajik AJ, et al (1996). Doppler echocardiographic index for assessment of global right ventricular function. *J Am Soc Echocardiogr* 9: 838–47. PMID: [8943444](#)
24. Kaul S, Tei C, Hopkins JM, Shah PM (1984). Assessment of right ventricular function using two-dimensional echocardiography. *Am Heart J* 107: 526–31. PMID: [6695697](#)
25. Girard O, Koehle MS, MacInnis MJ, Guenette JA, Koehle MS, Verges S, et al (2012). Comments on Point: Counterpoint: Hypobaric hypoxia induces/does not induce different responses from normobaric hypoxia. *J Appl Physiol* (1985) 112:1788–94.
26. Baulig W, Bernhard EO, Bettex D, Schmidlin D, Schmid ER (2005). Cardiac output measurement by pulse dye densitometry in cardiac surgery. *Anaesthesia* 60: 968–73. PMID: [16179040](#)
27. Compton F, Schäfer JH (2009). Noninvasive cardiac output determination: broadening the applicability of hemodynamic monitoring. *Semin Cardiothorac Vasc Anesth* 13: 44–55. doi: [10.1177/1089253208330711](#) PMID: [19147529](#)
28. Faiss R, Pialoux V, Sartori C, Faes C, Dériaz O, Millet GP (2013). Ventilation, oxidative stress, and nitric oxide in hypobaric versus normobaric hypoxia. *Med Sci Sports Exerc* 45: 253–60. doi: [10.1249/MSS.0b013e31826d5aa2](#) PMID: [22895381](#)
29. Sessler DI, McGuire J, Hynson J, Moayeri A, Heier T (1992). Thermoregulatory vasoconstriction during isoflurane anesthesia minimally decreases cutaneous heat loss. *Anesthesiology* 76:670–5. PMID: [1575332](#)
30. Schramm WM, Bartunek A, Gilly H (1997). Effect of local limb temperature on pulse oximetry and the plethysmographic pulse wave. *Int J Clin Monit Comput* 4:17–22.
31. Dempsey JA, Wagner PD (1999). Exercise-induced arterial hypoxemia. *J Appl Physiol* (1985) 87:1997–2006.
32. Naeije R (2010). Physiological adaptation of the cardiovascular system to high altitude. *Prog Cardiovasc Dis* 52: 456–66. doi: [10.1016/j.pcad.2010.03.004](#) PMID: [20417339](#)
33. Ogihara Y, Yamada N, Dohi K, Matsuda A, Tsuji A, Ota S, et al (2014). Utility of right ventricular Tei-index for assessing disease severity and determining response to treatment in patients with pulmonary arterial hypertension. *J Cardiol* 63:149–53. doi: [10.1016/j.jicc.2013.07.002](#) PMID: [24012334](#)
34. Zhou Q, Yang S, Luo Y, Qi Y, Yan Z, Shi Z, Fan Y (2012). A randomly-controlled study on the cardiac function at the early stage of return to the plains after short-term exposure to high altitude. *PLoS One* 7:e31097. doi: [10.1371/journal.pone.0031097](#) PMID: [22363556](#)
35. Stembridge M, Ainslie PN, Hughes MG, Stöhr EJ, Cotter JD, Nio AQ, Shave R (2014). Ventricular structure, function, and mechanics at high altitude: chronic remodeling in Sherpa vs. short-term lowlander adaptation. *J Appl Physiol* (1985) 117:334–43.
36. Huez S, Faoro V, Guénard H, Martinot JB, Naeije R (2009). Echocardiographic and tissue Doppler imaging of cardiac adaptation to high altitude in native highlanders versus acclimatized lowlanders.
37. Pagé M, Sauvé C, Serri K, Pagé P, Yin Y, Schampaert E (2013). Echocardiographic assessment of cardiac performance in response to high altitude and development of subclinical pulmonary edema in healthy climbers. *Am J Cardiol* 103:1605–9.
38. Eldridge MW, Braun RK, Yoneda KY, Walby WF (2006). Effects of altitude and exercise on pulmonary capillary integrity: evidence for subclinical high-altitude pulmonary edema. *J Appl Physiol* (1985) 100:972–80.