



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Campbell, MD and Walker, M and King, D and Gonzalez, JT and Allerton, DM and Stevenson, EJ and Shaw, JA and West, DJ (2016) Carbohydrate Counting at Meal Time Followed by a Small Secondary Postprandial Bolus Injection at 3 Hours Prevents Late Hyperglycemia, Without Hypoglycemia, After a High-Carbohydrate, High-Fat Meal in Type 1 Diabetes. *Diabetes care*, 39 (9). e141-e142. ISSN 0149-5992 DOI: <https://doi.org/10.2337/dc16-0709>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/2801/>

Document Version:

Article (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

Carbohydrate counting at meal time followed by a small secondary postprandial bolus injection at 3 hours prevents late hyperglycemia, without hypoglycemia, after a high-carbohydrate high-fat meal in Type 1 diabetes

Matthew D Campbell<sup>3,4</sup> PhD, Mark Walker<sup>1</sup> MD, David King<sup>2</sup> MRes, Javier T Gonzalez, PhD<sup>5</sup>, Dean Allerton<sup>2</sup> MSc, Emma J Stevenson<sup>1</sup> PhD, James A Shaw<sup>1</sup> FRCP, Daniel J West<sup>1</sup> PhD.

### **Affiliation**

<sup>1</sup>Institute of Cellular Medicine, Newcastle University, Newcastle-upon-Tyne, UK; <sup>2</sup>Faculty of Health and Life Sciences, Northumbria University, Newcastle-upon-Tyne, UK; <sup>3</sup>Carnegie School of Sport, Leeds Beckett University, UK; <sup>4</sup>Carnegie Research Institute, Leeds Beckett University, UK <sup>5</sup>Department for Health, University of Bath, UK

Short running title: insulin and high-fat meals in Type 1 diabetes

### **Corresponding author**

Daniel J. West

Institute of Cellular Medicine

Newcastle University, Newcastle-upon-Tyne,

NE2 4HH, UK.

Telephone: (+44) 191 2087076

Email: [Daniel.west@newcastle.ac.uk](mailto:Daniel.west@newcastle.ac.uk)

Manuscript word count: 500

Figures: 1

Clinical Trials Registration: NCT02595658

Type 1 diabetes patients are provided guidance and structured education on adjusting their meal time bolus-insulin dose based on meal carbohydrate content. However, recent research in patients using continuous subcutaneous insulin infusion (CSII) has highlighted the role of dietary fat in increasing prandial insulin requirements, particularly late into the postprandial period (1, 2).

Many patients are treated with basal-bolus insulin injections, which is a less flexible method of insulin therapy than CSII; for example, patients are unable to dual wave/extended bolus at meal time. It is important to consider that patients are encouraged to carbohydrate count and administer rapid-acting insulin units at meal-time, and that the time-action profiles of lispro/aspart vary in a dose dependent manner; while the peak insulin concentration after a pre-meal bolus usually occurs within the first 60 minutes (3), the peak action is usually observed between 90-120 minutes, and this duration can vary between 4-6 hours. However, peak postprandial lipemia occurs after 3-4 hours (4), which can promote acute peripheral insulin resistance and increase hepatic glucose output (5), and thus, hyperglycemia.



We investigated the influence of rapid-acting insulin dose and timing after high-carbohydrate high-fat meals in type 1 diabetes patients using insulin analogue injections.

10 male patients (mean±SD; age 26±4 years, BMI 25.4±1.6, diabetes duration 17±5 years, age at diagnosis 9±4 years; HbA<sub>1c</sub> 52.5±5.9 mmol/mol[7.0±0.5%]) using insulin aspart, and either basal insulin glargine (n=8) or detemir (n=2) attended the Newcastle NIHR Clinical Research Facility at 07:30h on four occasions.

Experimental trials were randomized and involved consuming either a 1) low-fat meal with bolus insulin dictated by carbohydrate counting (**Low-Fat100%**), 2) high-fat meal with bolus insulin dictated by carbohydrate counting (**High-Fat100%**), 3) high-fat meal with a bolus insulin dose increased by 30% (**High-Fat130%**), or 4) high-fat meal with bolus insulin dictated by carbohydrate counting, with an additional 30% administered at 3-h post-meal (**High-FatSplit**). Meals were matched for carbohydrate and protein content, but differed in fat (low fat: 68 g carbohydrate, 26 g protein, 5 g fat; High-fat: 68 g carbohydrate, 26 g protein, 55 g fat). Interval blood samples were collected over a 6 hour post-prandial period and were processed for glucose (Biosen C-Line, EKF-Diagnostic GmbH, London) and exogenous insulin (Invitron Insulin-Assay, Invitron, Monmouth, UK).

Glucose and insulin are presented in Figure 1-A and B, respectively. **High-Fat100%** was associated with late-hyperglycemia, while during **High-Fat130%**, 60% of patients experienced hypoglycemia (glucose<3.9 mmol/L), with no incidences under the other conditions. Post-prandial glycaemic excursions (time-course changes and area under the curve) were similar between **Low-Fat100%** and **High-FatSplit**, despite the additional 50 g of fat consumed in the latter trial.

When a meal has a high-carbohydrate and high-fat content, using the carbohydrate counting method for insulin dose adjustments at meal time, and administering additional bolus-insulin units 3 hours later, provides similar postprandial glucose control to a meal containing negligible fat, without causing hypoglycemia. Patients should be advised that increasing meal-time insulin dose alone is not an effective strategy, and is an approach that may increase the risk of early post-prandial hypoglycemia.



**Figure 1 A-B.** **A** Time course changes in blood glucose; **B** in Serum Insulin. Blue trace = **Low-Fat100%**; Black trace = **High-Fat100%**; Red trace = **High-FatSplit**; Green trace = **High-Fat130%**. Data presented as mean±SD (n=10). \* indicates that all conditions are significantly different from **High-Fat130%**, \*\* indicates that all conditions are significantly different from **High-Fat100%** and **High-Fat130%**.

## **ACKNOWLEDGEMENTS**

This study was funded by Newcastle and Northumbria Universities. The authors thank the study participants for their time, effort, and commitment, as well as the research team at the National Institute for Health Research Clinical Research Facility, Newcastle-upon-Tyne, for their assistance.

## **DUALITY OF INTEREST**

There are no potential conflicts of interest relevant to this article.

## **AUTHOR CONTRIBUTIONS**

**MDC** and **DJW** contributed to the study concept and design, researched data, and wrote the manuscript. **MW** and **JAS** contributed to participant recruitment and reviewed and edited the manuscript. **DK, JTG and DA** aided in data collection and data analysis. **EJS** contributed to the study concept and design, provided materials and reviewed and edited the manuscript. **DJW** is the guarantor of this work and, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and accuracy of data analysis.

## **REFERENCES**

1. Bell KJ, Smart CE, Steil GM, Brand-Miller JC, King B, Wolpert HA. Impact of fat, protein, and glycemic index on postprandial glucose control in type 1 diabetes: implications

for intensive diabetes management in the continuous glucose monitoring era. *Diabetes care*. 2015;38(6):1008-15.

2. Wolpert HA, Atakov-Castillo A, Smith SA, Steil GM. Dietary fat acutely increases glucose concentrations and insulin requirements in patients with type 1 diabetes: implications for carbohydrate-based bolus dose calculation and intensive diabetes management. *Diabetes care*. 2013;36(4):810-6.

3. Homko C, Deluzio A, Jimenez C, Kolaczynski JW, Boden G. Comparison of insulin aspart and lispro: pharmacokinetic and metabolic effects. *Diabetes care*. 2003;26(7):2027-31.

4. Roberts R, Bickerton AS, Fielding BA, Blaak EE, Wagenmakers AJ, Chong MF, et al. Reduced oxidation of dietary fat after a short term high-carbohydrate diet. *Am J Clin Nutr*. 2008;87(4):824-31.

5. Ferrannini E, Barrett EJ, Bevilacqua S, DeFronzo RA. Effect of fatty acids on glucose production and utilization in man. *The Journal of clinical investigation*. 1983;72(5):1737-47.