



LEEDS
BECKETT
UNIVERSITY

Citation:

Ispoglou, T and Deighton, K and King, R and White, H and Lees, M (2016) THE IMPACT OF ESSENTIAL AMINO ACID SUPPLEMENTS ENRICHED WITH L-LEUCINE ON APPETITE AND ENERGY INTAKE IN ELDERLY WOMEN. In: 38th ESPEN Congress, 17 September 2016 - 20 September 2016, Copenhagen, Denmark.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3086/>

Document Version:

Conference or Workshop Item (Other)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

The Impact of Essential Amino Acid Supplements Enriched With L-leucine on Appetite and Energy Intake in Elderly Women

MON-P108



Theocharis Ispoglou¹, Kevin Deighton¹, Roderick King¹, Helen White², Matt Lees¹

¹Leeds Beckett University, Institute for Sport Physical Activity & Leisure

²Leeds Beckett University, School of Clinical & Applied Sciences

Introduction

- Inadequate protein intake (PI), the main source of essential amino acids (EAAs), and reduced appetite are contributing factors to age-related sarcopenia.
- Protein supplementation frequently confers beneficial adaptations in performance and body composition.
- The satiating effects of dietary protein may negatively affect energy intake (EI), thus there is a need to explore alternative strategies to facilitate PI without compromising appetite and subsequent EI.

Results

- In **Experiment 1**, energy intake (EI) at ad libitum breakfast (ALB) was not different between conditions (282 ± 135 , 299 ± 122 , 288 ± 131 kcal for Control (C), Bar (B) and Gel (G) respectively). However, **total EI (Fig. 1a)** was significantly higher in B and G compared to C after accounting for the energy content of the supplements ($P < 0.0005$).
- Analysis revealed significantly higher appetite ratings (Fig. 2a) Area under the Curve (AUC) ($P < 0.007$), a tendency for higher acylated ghrelin AUC ($P = 0.087$), and significantly lower pancreatic polypeptide AUC ($P = 0.02$) in C compared with B and G (Fig. 2b).
- In **Experiment 2**, EI at ALB was significantly higher ($P = 0.028$) in C (306 ± 122 kcal) compared to B (245 ± 135 kcal) and G (254 ± 118 kcal). However, **total EI (Fig. 1b)** was significantly higher in B and G after accounting for the energy content of the supplements ($P < 0.007$).
- In both experiments, there were no differences between supplements for ratings of taste, aftertaste or overall palatability. Similarly no differences were observed in visual appeal, smell, taste, aftertaste and palatability of the breakfast.

Conclusion

- Supplementation with either the bar or gel increased total energy intake whether consumed 1h before or during breakfast.
- This may represent an effective nutritional means for addressing protein and total energy deficiencies in elderly women.

Total Energy Intake (EI)

Fig. 1a: Experiment 1









Fig. 1b: Experiment 2



Outline of Experimental Trials

- Key
- Conditions:
- Nothing (control)
 - Standard dose bar
 - Standard dose gel.
- ↑ Blood sample
↑ Assessment of appetite and/or appetite hormones



Methods

- Elderly women completed two experiments where they consumed either a Bar (B, 135 kcal) or a Gel (G, 114 kcal), rich in EAAs (7.5 g, 40% L-Leucine), or nothing [control (C)].
- In **Experiment 1**, subjects ($n=10$, 68 ± 5 years, mean \pm SD) consumed B, G or C with appetite sensations and appetite-related hormonal responses monitored for 1h, followed by consumption of an ad libitum breakfast (ALB).
- In **Experiment 2**, subjects ($n=11$, 69 ± 5 years) ingested B, G or C alongside an ALB.

Fig 2: Composite Appetite (CA) and Hormone Concentration

Fig. 2a: CA ratings in the control (▼), bar (●) and gel (○) trials (experiment one). Values are mean (SEM).





Fig. 2b: Plasma peptide YY (a), acylated ghrelin (b) and pancreatic polypeptide (c) concentrations in the control (▼), bar (●) and gel (○) trials (experiment one). Values are mean (SEM).



References & Acknowledgments

- Dillon, E. L., Sheffield-moore, M., Paddon-jones, D., Gilkison, C., Sanford, A. P., Casperson, S. L., Jiang, J., Chinkes, D. L. & Urban, R. J. (2009) Amino acid supplementation increases lean body mass, basal muscle protein synthesis, and insulin-like growth factor-I expression in older women. *J Clin Endocrinol Metab*, 94, 1630-7.
- Giezenaar, C., Chapman, I., Luscombe-marsh, N., Feinle-bisset, C., Horowitz, M. & Soenen, S. (2016) Ageing Is Associated with Decreases in Appetite and Energy Intake—A Meta-Analysis in Healthy Adults. *Nutrients*, 8.
- Gregorio, L., Brindisi, J., Kleppinger, A., Sullivan, R., Mangano, K. M., Bihuniak, J. D., Kenny, A. M., Kerstetter, J. E. & Insogna, K. L. (2014) Adequate dietary protein is associated with better physical performance among post-menopausal women 60-90 years. *J Nutr Health Aging*, 18, 155-60.
- Ispoglou, T., White, H., Preston, T., Mcelhone, S., Mckenna, J. & Hind, K. (2016) Double-blind, placebo-controlled pilot trial of L-Leucine-enriched amino-acid mixtures on body composition and physical performance in men and women aged 65-75 years. *European Journal of Clinical Nutrition*, 70, 182-188.
- Leidy, H. J., Clifton, P. M., Astrup, A., Wycherley, T. P., Westerterp-plantenga, M. S., Luscombe-marsh, N. D., Woods, S. C. & Mattes, R. D. (2015) The role of protein in weight loss and maintenance. *Am J Clin Nutr*.
- Morley, J. E. (1997) Anorexia of aging: physiologic and pathologic. *Am J Clin Nutr*, 66, 760-73.
- Tieland, M., Van De Rest, O., Dirks, M. L., Van Der Zwaluw, N., Mensink, M., Van Loon, L. J. & De Groot, L. C. (2012) Protein supplementation improves physical performance in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc*, 13, 720-6.
- Zhu, K., Meng, X., Kerr, D. A., Devine, A., Solah, V., Binns, C. W. & Prince, R. L. (2011) The effects of a two-year randomized, controlled trial of whey protein supplementation on bone structure, IGF-1, and urinary calcium excretion in older postmenopausal women. *J Bone Miner Res*, 26, 2298-306.

Sources of Funding: Higher Education Innovation Funding (HEIF), and Leeds Beckett University

Presented at the 38th ESPEN Congress on Clinical nutrition and Metabolism, Copenhagen, Denmark, 17-20 September 2016.