



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Cooper, L and Ryan, C and Ells, LJ and Hamilton, S and Atkinson, G and Cooper, K and Johnson, MI and Kirwan, JP and Martin, D (2016) Weight-loss interventions for overweight/obese adults with chronic musculoskeletal pain: a mixed methods systematic review protocol. JBI Database System Rev Implement Rep, 14 (5). pp. 57-67. ISSN 2202-4433 DOI: <https://doi.org/10.11124/JBISRIR-2016-002725>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3100/>

Document Version:

Article (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

## Review title

Weight loss interventions for overweight/obese adults with chronic musculoskeletal pain: a mixed methods systematic review protocol

## Reviewers

Lesley Cooper RGN, CertEd, MSc<sup>1,2</sup>

Cormac Ryan BSc, MSc, PhD<sup>1,2</sup>

Louisa Jane Ells BSc (Hons) PhD<sup>1,2</sup>

Sharon Hamilton PhD, MA, BA (Hons),RN<sup>1,2</sup>

Greg Atkinson BSc (Hons), PhD<sup>1,2</sup>

Kay Cooper BSc, MSc, PhD<sup>3</sup>

Mark I. Johnson BSc, PGCertHE, PhD<sup>4</sup>

John P. Kirwan BA(Hons), MSC, PhD<sup>5</sup>

Denis Martin BSc. MSc. DPhil<sup>1,2</sup>

1 Health and Social Care Institute, Teesside University, United Kingdom

2 Teesside Centre for Evidence Based Practice: an affiliate centre of the Joanna Briggs Institute

3 The Scottish Centre for Evidenced-Based, Multi-professional Practice: A Collaborating Centre of the Joanna Briggs Institute, Robert Gordon University, UK

4 Leeds Beckett University, UK

5 Lerner Research Institute, Department of Pathobiology, Cleveland Clinic, Ohio, USA

### Corresponding author:

Lesley Cooper, [l.cooper@tees.ac.uk](mailto:l.cooper@tees.ac.uk)

## Review question/objective

The objective of this mixed methods review is to develop an aggregated synthesis of qualitative and quantitative data on weight loss interventions for overweight/obese adults with chronic musculoskeletal pain in an attempt to derive conclusions and recommendations useful for clinical practice and policy decision making

The objective of the quantitative component of this review is to quantify the effectiveness of weight loss interventions on weight, pain, and physical and/or psychosocial function in overweight/obese adults

with chronic musculoskeletal pain.

The objectives of the qualitative component of this review are to explore the perceptions and experiences of overweight/obese adults with chronic musculoskeletal pain of the link between their weight and pain and the effectiveness and appropriateness of weight loss interventions and sustainability of weight loss efforts

## Background

Independently, overweight and chronic pain are prevalent conditions that have widespread implications for the individual, health care resources and the economy.<sup>1,2</sup> These two conditions frequently occur simultaneously and the association appears to be bi-directional, this adds to the complexity of managing either condition independently. Studies to assess the effectiveness of weight loss interventions on individuals with co-existing chronic musculoskeletal pain have shown that weight reduction can be achieved and is associated with lower pain scores<sup>3,4</sup>, however no systematic review has been conducted to determine the most effective intervention or participants' perceptions of the appropriateness and sustainability of interventions to inform clinical practice.

The World Health Organisation (WHO) defines obesity as “abnormal or excessive fat accumulation that may impair health”.<sup>5</sup> Obesity is categorised using Body Mass Index (BMI). A BMI  $\geq 25$  indicates that a person is overweight and a BMI  $\geq 30$  is deemed as obese.<sup>6</sup> This growing global public health problem has been associated with 3.4 million adult deaths annually and is linked to the development of type 2 diabetes, coronary heart disease (CHD) and some cancers.<sup>5</sup>

Worldwide prevalence of overweight/obesity has risen markedly in recent decades with rates more than doubling in some developed areas; globally 1.9 billion adults are overweight, 600 million of these are obese.<sup>5</sup> The United Kingdom (UK) has one of the highest levels of overweight/obesity in Europe; 66.6 % of men and 57.2% of women in England were classified as overweight/obese in 2012 (24% of men and 25% of women were obese).<sup>8,9</sup> Cohort studies carried out by the Centre for Longitudinal Studies found that at age 42 those in the 1970 birth cohort were markedly more likely to be overweight or obese compared to individuals in the 1958 birth cohort at the same age, highlighting the increase in adult obesity in the UK.<sup>10</sup> Obesity has been shown to reduce quality of life and limit ability to perform activities of daily living (ADLs),<sup>11</sup> it is also linked to higher levels of mental illness.<sup>10</sup> National Health Service (NHS) costs associated with obesity are currently estimated to be in excess of £5 billion annually.<sup>10,12</sup>

The causes of overweight/obesity in the general population are increased calorie intake and lack of physical exercise. However the Foresight Report describes this as a simplistic view and reports that overweight/obesity is the result of complex interaction between biological factors, psychological factors and a changing social environment wherein work patterns have become more sedentary and physical activity has decreased combined with an increase in the availability of food.<sup>12</sup> In 2013 the American Medical Association classified obesity as a disease.<sup>13</sup>

The International Association for the Study of Pain define pain as “An unpleasant sensory and emotional experience associated with actual or potential tissue damage”.<sup>14</sup> Chronic pain is persistent pain, which lasts over 12 weeks or after the time that healing would have been thought to occur after trauma or surgery.<sup>15</sup> Approximately 5 million adults in the UK annually develop chronic pain that impacts on their ability to work, thereby having an effect on business and the economy<sup>2</sup>. Although the total cost

of chronic pain is unknown the cost to the exchequer annually is in excess of £12.3 billion for back pain alone and 2.7 million people with chronic musculoskeletal pain are in receipt of incapacity benefit.<sup>15,16</sup> While back pain and osteoarthritis (OA) (particularly of the knee) account for 50% of chronic pain, other conditions such as rheumatoid arthritis, migraine, chronic daily headache and neck pain are also common. Five per cent of the UK population suffer from widespread chronic pain often referred to as fibromyalgia.<sup>16</sup> The 2011 Health Survey England (HSE)<sup>17</sup> and the National Pain Audit<sup>18</sup> found that chronic pain is more common in women, increases with age and is more likely to be reported by those in the lowest economic quintile. People reporting chronic pain also reported higher levels of anxiety and depression with a positive correlation between increasing pain and anxiety and depression scores.<sup>17,19</sup> Chronic pain sufferers are prone to have difficulty sleeping, socialising and are also more likely to become unemployed.<sup>16</sup>

Although no cause-effect relationship has yet been established there is growing acknowledgement of a link between obesity and chronic pain conditions.<sup>20-27</sup> For these individuals difficulties arise in everyday activities such as walking, climbing stairs and driving which result in a decline in independence, leading to reduction in mental health, with depression and social isolation being known to affect these patient groups.<sup>28</sup> As chronic pain interferes with daily functioning of obese individuals, it can have a negative effect on weight loss.<sup>11</sup>

The mechanical-structural nature of the relationship between obesity and chronic pain is generally accepted,<sup>22</sup> however it is not the definitive explanation for this comorbidity. In addition to mechanical explanations metabolic factors may be involved in altered pain sensitivity among obese individuals.<sup>29</sup> Furthermore evidence suggests that poor general health behaviours may have a role to play in the development and persistence of the comorbidity. Obesity may contribute to the development/maintenance of pain while the presence of pain may be a risk factor for the development/maintenance of obesity. Treatment needs to address both aspects of the comorbidity.<sup>30,31</sup>

National Institute for Clinical Excellence (NICE) recommend that overweight and obese adults are offered Lifestyle Weight management programs<sup>6</sup> while weight reduction is advocated as an integral component of chronic pain management.<sup>20,30</sup> Weight loss interventions may have an additional positive impact on pain by reducing mechanical-structural factors, metabolic factors, systemic inflammation and increasing health related quality of life (hrQoL).<sup>32</sup> Interventions designed to target weight may have added positive effects, e.g. reduction of mental distress, such as anxiety or depression, both of which are strongly correlated with obesity and chronic pain.<sup>10,175</sup> Improvement in mental health may enable individuals who are overweight or obese with co-existing chronic pain to sustain positive changes and maintain new patterns of behaviour.

Physical activity is a primary intervention for overweight/obesity, however, many overweight/obese individuals report chronic pain acts as a barrier to physical activity and thus weight loss.<sup>11</sup> Individuals with higher BMI have been shown to have increased prevalence of chronic pain.<sup>33</sup> Physical activity is also used as a treatment technique for those with chronic pain; however, obesity can be a barrier to physical activity for individuals who are obese with chronic pain.<sup>34,35</sup> The cause and consequence relationship between obesity and chronic pain can become a vicious cycle that impacts on the function, health and well-being of this population. The rising prevalence of both obesity and chronic pain is a worldwide concern, as alongside patient health and well-being concerns there is an increased financial burden on healthcare and economic systems.<sup>33,36,37</sup>

The complex relationship between chronic pain and overweight/obesity appears to be bi-directional in nature; for example increased weight may cause an individual to experience pain that reduces activity or pain may restrict activity which results in increased weight that produces more pain, therefore a break in this cycle could have positive outcomes for individuals. A comprehensive search has failed to identify any systematic reviews that have been completed or any protocols for systematic reviews currently in progress on the effectiveness, appropriateness and sustainability of weight loss interventions in an overweight/obese population with co-existing chronic musculoskeletal pain. This review will bring together quantitative and qualitative evidence to increase knowledge of the effectiveness of interventions in this population and understanding of individual perceptions of the links between overweight/obesity and chronic pain, the effectiveness, and appropriateness of weight loss interventions and sustainability of weight loss efforts. This review should inform future interventions for the promotion of healthier lifestyles, thus increasing weight loss and decreasing chronic pain in a more sustainable manner. In addition to individual positive health outcomes, there is potential for wider societal and economic benefits<sup>33,37</sup> e.g. considerable savings could be made on healthcare spending on care of patients with OA knee if cases attributed to overweight/obesity (currently 26.4%) were reduced.

## **Inclusion criteria**

### **Types of participants**

The quantitative and qualitative components of this review will consider studies that include male or female adult participants (aged 18 or over) of any ethnic origin who are overweight/obese and have co-existing chronic pain. We will consider studies that include participants diagnosed with chronic musculoskeletal pain conditions such as, chronic lower back pain (CLBP), OA or rheumatoid arthritis (RA), as well as those who suffer non-specific or widespread musculoskeletal pain conditions. The diagnosis of overweight/obesity should be consistent with the WHO definition (BMI  $\geq 25$  overweight; BMI  $\geq 30$  obese)<sup>5</sup> The diagnosis of chronic pain should be consistent with the BPS definition (persistent pain, which lasts over 12 weeks or after the time that healing would have been thought to occur after trauma or surgery).<sup>15</sup>

Studies that include participants with; chronic pain related to cancer; pregnant or breast-feeding women; or syndromic obesity will be excluded from the review.

### **Types of intervention(s)/phenomena of interest**

The quantitative component of the review will consider studies that evaluate weight management treatment programmes. Interventions can include bariatric drugs (e.g. orlistat), surgery (e.g. gastric banding), and lifestyle modifications such as diet, physical activity or psychological interventions delivered as part of a multi or single component study. Comparator: intervention will be compared with no treatment (true control) or usual care in any experimental study design.

The qualitative component of this review will include studies that explore the perceptions and experiences of co-existing overweight/obesity and chronic musculoskeletal pain, engaging with weight loss interventions and sustaining weight loss efforts long-term.

### **Types of outcomes**

The quantitative component of this review will consider studies that include the following outcome

measures - primary outcome: weight; any objective validated measure of adiposity (BMI, waist circumference). Secondary outcomes; pain - any validated measure of pain - numeric rating scale (NRS)/ visual analogue scale (VAS); adverse effects; psychological health - measured using standard scales (e.g. Hospital Anxiety and Depression Scale (HADS)); physical and/or psychological function, quality of life (e.g. Health Related Quality of Life (HRQoL)); use of analgesic medication (e.g. Naproxen, Codiene, opioids); contacts with health care professionals and hospital admission.

### **Types of studies**

The quantitative component of the review will be restricted to experimental study designs including randomised controlled trials and quasi-experimental trials for inclusion.

The qualitative component of the review will consider qualitative studies of experience and perceptions including (but not limited to) designs such as interview studies, focus group studies, ethnography, phenomenology, grounded theory and action research.

## **Search strategy**

The search strategy aims to find both published and unpublished studies. A three-step search strategy will be utilised in this review. An initial limited search of MEDLINE and CINAHL will be undertaken followed by analysis of the text words contained in the title and abstract, and of the index terms used to describe the article. A second search using all identified keywords and index terms will then be undertaken across all included databases. Thirdly, the reference list of all key identified reports and articles will be searched for additional studies. Only studies published in English will be considered for inclusion in this review. Studies published 1990- present will be considered for inclusion in this review. This time frame was selected as overweight/obesity rates have been increasing steadily during this period.<sup>3</sup>

The databases to be searched include:

The Cochrane Library, MEDLINE, CINAHL, PubMed, Scopus, PsycINFO, EMBASE, Education Resources Information Centre (ERIC), Web of Science

The search for unpublished studies will include:

Clinical trials registries (e.g. clinicaltrials.gov), dissertations indexed with ProQuest Dissertations & Theses

Initial keywords to be used will be:

Overweight, obesity, obese, chronic pain, chronic lower back pain, chronic joint pain, osteoarthritis, rheumatoid arthritis, chronic widespread pain, fibromyalgia, persistent pain, chronic headache, weight loss, weight management, weight control, diet control, exercise therapy, exercise, structured exercise, cognitive behavioural therapy, CBT, pain coping skills training, bariatric surgery, gastric surgery, gastric bypass, gastric banding, orlistat, experiences, perceptions

## **Assessment of methodological quality**

Quantitative papers selected for retrieval will be assessed by two independent reviewers for

methodological validity prior to inclusion in the review using standardised critical appraisal instruments from the Joanna Briggs Institute Meta Analysis of Statistics Assessment and Review Instrument (JBI-MAStARI) (Appendix I. Any disagreements that arise between the reviewers will be resolved through discussion, or with a third reviewer.

Qualitative papers selected for retrieval will be assessed by two independent reviewers for methodological validity prior to inclusion in the review using standardised critical appraisal instruments from the Joanna Briggs Institute Qualitative Assessment and Review Instrument (JBI-QARI) (Appendix I. Any disagreements that arise between the reviewers will be resolved through discussion, or with a third reviewer.

## Data extraction

### Stage 1 data extraction

Quantitative data will be extracted from papers included in the review using the standardised data extraction tool from JBI-MAStARI (Appendix II. The data extracted will include specific details about the interventions, populations, study methods and outcomes of significance to the review question and specific objectives.

Qualitative data will be extracted from papers included in the review using the standardised data extraction tool from JBI-QARI (Appendix II. The data extracted will include specific details about the phenomena of interest, populations, study methods and outcomes of significance to the review question and specific objectives.

### Stage 2 data extraction

The results of each single method synthesis included in the mixed method review will be extracted in numerical, tabular or textual format. For example, for syntheses of quantitative data, this will consist of appropriate elements of the meta-analysis Forest plot or, where applicable an evidence table; for qualitative reviews, it will consist of appropriate elements of the QARI-view table.

## Data synthesis

### Stage 1 data synthesis for each single-method synthesis

Treatment effect sizes will, where possible be pooled in a meta-analysis using Comprehensive Meta-analysis (CMA) and Stata. All results will be subject to double data entry. We anticipate most of the data from each study will be effect sizes expressed either as odds ratios (for categorical data) and/or the mean difference between baseline and follow-up (for continuous data). Pooled effects sizes (and associated 95% confidence intervals) will be quantified in a weighted fashion using the inverse variance approach. Heterogeneity will be quantified using I-squared and Tau-squared statistics. If appropriate, heterogeneity sources will be explored with subgroup analyses and/or meta-regression approaches. Where statistical pooling is not possible the findings will be presented in narrative form including tables and figures to aid in data presentation where appropriate.

Qualitative research findings will, where possible be pooled using JBI-QARI. This will involve the

aggregation or synthesis of findings to generate a set of statements that represent that aggregation, through assembling the findings (Level 1 findings) rated according to their quality, and categorising these findings on the basis of similarity in meaning (Level 2 findings). These categories are then subjected to a meta-synthesis in order to produce a single comprehensive set of synthesised findings (Level 3 findings) that can be used as a basis for evidence-based practice. Where textual pooling is not possible the findings will be presented in narrative form.

Stage 2 data synthesis for mixed method synthesis

The findings of each single-method synthesis included in this review will be aggregated according to the JBI Reviewers' Manual Methodology for JBI Mixed Methods Systematic Reviews<sup>38</sup>. This will involve the configuration of the findings to generate a set of statements that represent that aggregation through coding any quantitative findings to attribute a thematic description to all quantitative data; assembling all of the resulting themes from quantitative and qualitative syntheses; and the configuration of these themes to produce a set of synthesised findings in the form of a set of recommendations or conclusions

## Conflicts of interest

None

## Acknowledgements

## References

1. Morgan, E., Dent, M. The economic burden of obesity. Oxford: National Obesity Observatory. 2010
2. Phillips, C.J., The cost and burden of chronic pain. *Reviews in Pain* 2009 3(1) 2-5
3. Somers, T.J., Blumenthal, J.A., Guilak, F., Kraus, V.B., Schmitt, D.O., Babyak, M.A., Craighead, L.W., Caldwell, D.S., Rice, J.R., McKee, D.C., Shelby, R.A., Campbell, L.C., Pells, J.J., Sims, E.L., Queen, R., Carson, J.W., Connelly, M., Dixon, K.E., LaCaille, L.J., Huebner, J.L., Rejeske, W.J., Keefe, F.J. Pain Coping Skills Training and Lifestyle Behavioural Weight Management in Patients with Knee Osteoarthritis; A Randomized Controlled Study. *Pain*.2012; 153(6): 1109-1129.
4. Foy, C.G., Lewis, C.E., Hairston, K.G., Miller, G.D., Lange, W., Jakicic, J.M., Rejeski, W.J., Ribisi, P.M. Intensive Lifestyle Intervention Improves Physical Function Among Obese Adults With Knee Pain: Findings From the Look AHEAD Trial. *Obesity (Silver Spring)*.2011; 19(1): 83-93.
5. World Health Organisation. Obesity and Overweight. Factsheet No 311. 2014.
6. National Institute for Clinical Excellence. Managing Overweight and obesity in adults - lifestyle weight management services. 2014; ph55.

7. National Institute for Clinical Excellence. Osteoarthritis Care and Management. CG177. 2014.
8. Department of Health. Health Survey for England 2012. 2013.
9. Health and Social Care Information Centre. Statistics on Obesity, Physical Activity and Diet - England 2014. 2014.
10. Centre for Longitudinal Studies. Overweight and obesity in midlife: Evidence from the 1970 British Cohort Study at age 42. . Institute of Education. University of London. 2014.
11. Wachholtz, A.; Binks, M.; Eisenson, H.; Kolotkin, R.; Suzuki, A. Does Pain Predict interference with daily functioning and weight loss in an obese residential treatment-seeking population?. International Society of Behavioural Medicine.2010; 17: 118-124.
12. Butland, B.;Jebb, S.; Kopleman, P. Tackling obesities: future choices - project report (2nd Edition). London. the Stationery Office. 2007.
13. American Medical Association House of Delegates. Resolution 420 (A-13). <http://www.ama-assn.org/assets/meeting/2013a/a13-addendum-refcomm-d.pdf> 2013.
14. International Association for the Study of Pain. Global year against pain. Acute Pain Factsheets 3:Mechanisms of Acute Pain. <http://www.iasp-pain.org/Taxonomy?navItemNumber=576> 2015
15. British Pain Society. FAQ's. <https://www.britishpainsociety.org/media-resources/#faqs> 2008
16. Donaldson, Sir L. Pain: Breaking the Barrier. Chief Medical Office Annual Report. 2008
17. Department of Health. Health Survey for England 2011. 2012.
18. Price, C.; Hoggart, B.; Olukoga, O.; Williams , A.; Bottle, A. National Pain Audit 2010-2012. 2012.
19. McCarthy,L.H.; Bigal,M.E.; Katz,M.; Derby,C.; Lipton,R.B. Chronic pain and obesity in elderly people: results from the Einstein Aging Study. Journal of the American Geriatrics Society.2009; 57(1): 115-119.
20. Silverwood, V.; Blagojevic-Bucknall, M.; Jinks, C.; Jordan, J.L.; Protheroe, J.; Jordan,K.P. Current evidence on risk factors for knee osteoarthritis in older adults: a systematic review and meta-analysis. Osteoarthritis and Cartilage.2015; 23(4): 505-515.
21. Blagojevic, M.; Jinks, C.; Jeffry, A.; Jordan, K.P. Risk Factors for the onset of knee osteoarthritis in older adults: a systematic review and meta-analysis. Osteoarthritis & Cartilage.2010; 18: 24-33.
22. Shiri, R.;Karppinen, J.; Leino-Argas, P.; Solovieva S.; Viikari-Juntura, E. The association between obesity and low back pain: a meta-analysis. American Journal of Epidemiology.2010; 17(2): 135-154.
23. Somers, T.J.; Wren, A.A.; Keefe, F.J. Understanding chronic pain in older adults: Abdominal fat is where it is at. PAIN.2011; 152: 8-9.
24. Heuch,I.; Heuch,I.; Hagen,K.; Zwart,J.A. Body Mass Index as a Risk Factor for Developing Chronic Low Back Pain: A Follow-up in the Nord-Trøndelag Health Study. Spine.2013; 38(2): 133-139.
25. Okifuji,A.; Donaldson,G.W.; Barck,L.; Fine,P.G. Relationship between fibromyalgia and obesity in pain, function, mood, and sleep. Journal of Pain.2010; 11(12): 1329-1337.
26. Ray,L.; Lipton,R.B.; Zimmerman,M.E.; Katz,M.J.; Derby,C.A. Mechanisms of association between

obesity and chronic pain in the elderly. *PAIN*.2011; 152(1): 53-59.

27. Thomazeau, J.; Perin, J.; Nizard, R.; Bouhassira, D.; Collin, E.; Nguyen, E.; Perrot, S.; Bergmann, J.F.; Lloret-Linares, C. Pain management and pain characteristics in obese and normal weight patients before joint replacement. *Journal of evaluation in clinical practice*.2014; 20(5): 611-616.

28. Gatineau, M.; Hancock, C.; Dent, M.. Adult obesity and disability.. National Obesity Observatory Oxford.2013

29. Janke, A.E.; Collins, A.; Kozak, A.T.. Overview of the relationship between pain and obesity: What do we know? Where do we go next?. *Journal of rehabilitation research & development*.2007; 44(2): 245-262.

30. Bonakdar, R.A. Obesity Related pain; time for a new approach that targets systemic inflammation. Supplement To The *Journal of Family Practice*.2013; 62(9): 22-28.

31. McVinnie, D.S.. Obesity and Pain. *British Journal of Pain*.2013; 7(4): 163-170.

32. Morden, A.; Jinks, C.; Nio Ong, B.. "...I've found once the weight had gone off, I've had a few twinges, but nothing like before". Exploring weight and self-management of knee pain. *Musculoskeletal Care*.2014; 12: 63-73.

33. Wedin, S.; Byrne, K.; Morgan, K.; LePage, M.; Goldman, R.; Crowley, N.; Galloway, S.; Borckardt, J.J. Presurgical weight is associated with pain, functional impairment and anxiety among gastric bypass surgery patients. *Pain Research and Treatment*.2012: 1-5.

34. Janke, A.E.; Kozak A.T.. "The more pain I have the more I want to eat": Obesity in the context of chronic pain. *Obesity*.2012; 20: 2027-2034.

35. de Rooij, M.; Steultjens, M.P.M.; Avezaat, E.; Häkkinen, A.; Klaver, R.; van, der Leeden; M, T.; Roorda, L., D.; van, der Velde; L.W.F.; Dekker, J. Restrictions and contraindications for exercise therapy in patients with hip and knee osteoarthritis and comorbidity. *Physical Therapy Reviews*.2013; 18(2): 101-111.

36. Wright, L.J.; Schur, C.; Noonan, C.; Ahumada, S.; Buchwald, D.; Afari, N. Chronic Pain, Overweight and Obesity: Findings from a Community-based Twin Registry. *The Journal of Pain*.2010; 11(7): 628-635.

37. Wang, Y.C.; McPherson, K.; Marsh, K.; Gortmaker, S.L.; Brown, M.. Health and economic burden of the projected obesity trends in the USA and the UK. *The Lancet*.2011; 378: 815-825.

38. The Joanna Briggs Institute. Joanna Briggs Institute Reviewers' Manual 2014 edition/Supplement.

## Appendix I: Appraisal instruments

### QARI Appraisal instrument

#### JBI QARI Critical Appraisal Checklist for Interpretive & Critical Research

Reviewer ..... Date .....

Author ..... Year ..... Record Number .....

|  | Yes                      | No                       | Unclear                  | Not Applicable           |
|--|--------------------------|--------------------------|--------------------------|--------------------------|
| 1. Is there congruity between the stated philosophical perspective and the research methodology?   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Is there congruity between the research methodology and the research question or objectives?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Is there congruity between the research methodology and the methods used to collect data?   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Is there congruity between the research methodology and the representation and analysis of data?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Is there congruity between the research methodology and the interpretation of results?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Is there a statement locating the researcher culturally or theoretically?   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. Is the influence of the researcher on the research, and vice-versa, addressed?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 8. Are participants, and their voices, adequately represented?   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 9. Is the research ethical according to current criteria or, for recent studies, and is there evidence of ethical approval by an appropriate body? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 10. Do the conclusions drawn in the research report flow from the analysis, or interpretation, of the data?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Overall appraisal:  Include  Exclude  Seek further info.

Comments (Including reason for exclusion)

---



---



---

**MAStARI Appraisal instrument**

**JBI Critical Appraisal Checklist for Randomised Control / Pseudo-randomised Trial**

Reviewer ..... Date .....

Author ..... Year ..... Record Number .....

|   | Yes                      | No                       | Unclear                  | Not Applicable           |
|---|--------------------------|--------------------------|--------------------------|--------------------------|
| 1. Was the assignment to treatment groups truly random?                             | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Were participants blinded to treatment allocation?                               | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Was allocation to treatment groups concealed from the allocator?                 | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Were the outcomes of people who withdrew described and included in the analysis? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Were those assessing outcomes blind to the treatment allocation?                 | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Were the control and treatment groups comparable at entry?                       | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. Were groups treated identically other than for the named interventions?          | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 8. Were outcomes measured in the same way for all groups?                           | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 9. Were outcomes measured in a reliable way?  | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 10. Was appropriate statistical analysis used?                                      | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
|   | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

Overall appraisal:    Include                     Exclude                     Seek further info.

Comments (Including reason for exclusion)

---



---



---

## Appendix II: Data extraction instruments

### QARI data extraction instrument

#### **JBI QARI Data Extraction Form for Interpretive & Critical Research**

Reviewer ..... Date .....

Author ..... Year .....

Journal ..... Record Number .....

#### **Study Description**

Methodology

---

Method

---

Phenomena of interest

---

Setting

---

Geographical

---

Cultural

---

Participants

---

Data analysis

---

Authors Conclusions

---

Comments

---

Complete

Yes

No

| Findings | Illustration from Publication (page number) | Evidence    |          |             |
|----------|---|-------------|----------|-------------|
|          |   | Unequivocal | Credible | Unsupported |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |
|          |   |             |          |             |

Extraction of findings complete

Yes

No

**MAStARI data extraction instrument**

**JBI Data Extraction Form for  
Experimental / Observational Studies**

Reviewer ..... Date .....

Author ..... Year .....

Journal ..... Record Number .....

**Study Method**

RCT                       Quasi-RCT                       Longitudinal   
Retrospective                       Observational                       Other

**Participants**

Setting \_\_\_\_\_

Population \_\_\_\_\_

**Sample size**

Group A \_\_\_\_\_ Group B \_\_\_\_\_

**Interventions**

Intervention A \_\_\_\_\_

Intervention B \_\_\_\_\_

Authors Conclusions:  
\_\_\_\_\_  
\_\_\_\_\_

Reviewers Conclusions:  
\_\_\_\_\_  
\_\_\_\_\_