



Citation:

Worrillow, N and Briggs, M and South, J (2016) An Analysis of the Identification and Management of Frequent Users of Urgent and Emergency Healthcare. *Emergency Medicine Journal*, 33 (9). ISSN 1472-0205 DOI: <https://doi.org/10.1136/emermed-2016-206139.5>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3368/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Background

There is a dearth of literature available that examines the characteristics of frequent users of urgent and emergency healthcare. Most research and literature explores this population by focusing within one clinical service rather than throughout the whole urgent and emergency healthcare system in England.

The aim was to examine how frequent users of: Out of Hours General Practitioners; NHS 111; Ambulance Services; and Emergency Departments are defined, what thresholds are used to identify them and what approaches are used to manage this population.

Method

A systemic mapping exercise was undertaken to capture a wide range of data to gain a comprehensive understanding of how these services are currently working with regard to frequent users.

The search provided 5094 articles which were screened against the inclusion criteria, leaving 156 articles to be analysed. Data was extracted and thematic synthesis created four main themes: typology with regard to definitions assigned; threshold variance; separation of the main population into subgroups; and case management approaches.

Results

The systematic mapping exercise revealed that diverse definitions and thresholds were being employed by all services; the language used to describe these individuals was often negative and subjective. A single definition of High Intensity User (HIUs) has been selected and advocated for standardisation within all services. This transposable terminology and identification process will aid in supporting and managing patients within all these services.

A taxonomy of HIUs has been developed from this research to sub-divide the main heterogeneous group into 5 main classification groups. This meets a gap in current knowledge.

Conclusions

The standardised identification process and taxonomy will facilitate interoperability and clarity between clinicians, thus benefitting the patient and their outcome. The support and management of HIUs will become more comparable between all urgent and emergency services and within the research community.