



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Serna, H and Balaskas, K and Mahmoodi, N and Aslam, T (2017) Experience of Anti-VEGF Treatment and Clinical Levels of Depression and Anxiety in Patients With Wet Age-Related Macular Degeneration. American Journal of Ophthalmology. ISSN 0002-9394 DOI: <https://doi.org/10.1016/j.ajo.2017.03.005>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3649/>

Document Version:

Article (Accepted Version)

---

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

Journal of Ophthalmology

Elsevier Editorial System(tm) for American

Manuscript Draft

Manuscript Number: AJO-16-1566R3

Title: Experience of Anti-VEGF Treatment and Clinical Levels of Depression and Anxiety in Patients with Wet Age-Related Macular Degeneration

Article Type: Original Article

Keywords: Wet Age-Related Macular Degeneration; Anti-VEGF; Depression; Anxiety; Experience.

Corresponding Author: Dr. Hugo Senra, PhD

Corresponding Author's Institution: University of Manchester

First Author: Hugo Senra, PhD

Order of Authors: Hugo Senra, PhD; Konstantinos Balaskas, MD; Neda Mahmoodi, PhD; Tariq Aslam, MD; PhD

## Abstract

**Purpose:** To investigate detailed patient experiences specific to receiving vascular endothelial growth factor inhibitors (anti-VEGF) for wet Age-Related Macular Degeneration (wAMD), and to acquire a snapshot of the frequency of clinically significant levels of depression, anxiety and post-traumatic stress among patients and levels of burden in patients' carers.

**Design:** Observational cross-sectional mixed-methods study

**Methods:** 300 patients with wAMD receiving anti-VEGF treatment and 100 patient carers were recruited. Qualitative data on patients' experience of treatment were collected using a structured survey. Standardised validated questionnaires were used to quantify clinically significant levels of anxiety, depression, post-traumatic stress, as well as cognitive function and carers' burden.

**Results:** Qualitative data showed that 56% of patients (n =132) reported anxiety related to anti-VEGF treatment. The main sources of anxiety were fear of going blind due to intravitreal injections, and concerns about treatment effectiveness rather than around pain. From validated questionnaires, 17% of patients (n=52) showed clinical levels of anxiety, and 12% (n=36) showed clinical levels of depression. Depression levels, but not anxiety, were significantly higher in patients who received up to 3 injections compared to patients who received from 4 to 12 injections (ANOVA  $P=.027$ ) and compared to patients who received more than 12 injections (ANOVA  $P=.001$ ).

**Conclusions:** Anti-VEGF treatment is often experienced with some anxiety related to treatment regardless of the number of injections received. Clinical levels of depression seem to be more frequent in patients at early stages of anti-VEGF treatment. Strategies to improve patient experience of treatment and minimise morbidity are suggested.

**Key Words:** Wet Age-Related Macular Degeneration; Anti-VEGF; Depression; Anxiety; Experience.

**Title:** Experience of Anti-VEGF Treatment and Clinical Levels of Depression and Anxiety in Patients with Wet Age-Related Macular Degeneration

Authors:

Hugo Senra<sup>1</sup>, PhD; Konstantinos Balaskas<sup>2</sup>, MD; Neda Mahmoodi<sup>3</sup>, PhD; Tariq Aslam<sup>1,2</sup>, MD, PhD

<sup>1</sup>Division of Pharmacy and Optometry, School of Health Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, UK

<sup>2</sup>Manchester Royal Eye Hospital, Central Manchester Foundation Trust, Manchester, UK.

<sup>3</sup> Faculty of Health and Social Sciences, School of Psychology, Leeds Beckett University, U.K

**Correspondence:**

Tariq Aslam, MD, PhD

Manchester Royal Eye Hospital

Oxford Road

Manchester M13 9PL

UK

Email: Tariq.Aslam@cmft.nhs.uk

**Short Title:** Patient Experience of Anti-VEGF treatment for Wet AMD

## Introduction

Age-related macular degeneration (AMD) is one of the leading causes of vision loss and blindness in people aged 50 years and older, particularly in the developed world (1,2). Currently, wet AMD (wAMD) is the only form of AMD that is treatable, usually involving the use of vascular endothelial growth factor inhibitors (anti-VEGF) such as bevacizumab, ranibizumab, or aflibercept (3,4). Anti-VEGF treatments are regarded as a generational breakthrough in the treatment of macular diseases such as wAMD (5). Recent studies have shown that anti-VEGF treatment can significantly reduce patients' risk of severe visual impairment as it has great potential for halting disease progression (3-5). However, this treatment is administered by invasive intravitreal injections, often at the conclusion of lengthy, frequent and repeated visits after transport to a suitably equipped hospital. The specific act of these intraocular anti-VEGF injections can be experienced by patients as a stressful event, with anticipatory anxiety and expectations of pain and discomfort (6-8). Evidence on this topic is still limited due the lack of studies exploring the complexity and diversity of patient experiences of anti-VEGF treatment for wAMD (9). A recent review of the literature (9) only found 3 studies exploring in-depth patients' experiences of receiving anti-VEGF treatments using qualitative designs (8,10,11). The remaining studies addressing the experience of receiving anti-VEGF treatment were mainly focused on medical aspects such as the anaesthetic's effectiveness to prevent pain when receiving an intravitreal injection (12-17). The studies suggest that anticipatory anxiety associated with anti-VEGF treatment tends to be circumscribed to the first injections, i.e. at early stages of treatment (17,18). The aspects of anti-VEGF treatment in which patients have reported more discomfort and anxiety are (6-8,10,18): needle entry; application of drops; insertion of speculum; waiting for injection; experiences of pain, fear of losing sight, fear for the unknown and side effects. One qualitative study (11) also stressed the potential importance of other variables for the way patients experience anti-VEGF treatment and cope with anxiety and the fear for the "unknown". These variables are: communication between healthcare professionals and patients, especially about treatment procedures; and the way service is delivered, with a special attention for anticipatory anxiety induced by waiting times before the injection. In another qualitative study on patient experience of anti-VEGF treatment (8), cautious optimism regarding treatment success and disease stabilization was found as a frequent and valid response to treatment from wet AMD patients. There is, overall, a need for more research exploring the complexity of patient experience of anti-VEGF treatment to clarify reasons underlying patients' anxieties when receiving regular intravitreal injections (9). The relationship between AMD and depression and anxiety has been well documented in literature (19-24). Research shows a wide prevalence range from 15.7% to 44% for depressive symptoms and 9.6% to 30.1% for anxiety symptoms among AMD patients (19). Previous studies have suggested that vision-related disability (caused by visual impairment) (25-28), visual acuity (29,30), and social support (31,32) can play an important role in the link between visual impairment and depression. Other studies have highlighted the link between ageing and depression (33) which might also play a role in the relationship between AMD and depression. However, most studies on depression associated with AMD were conducted prior to the implementation of anti-VEGF treatments or were conducted on general AMD patients (dry and wet) (22,34). This might be explained by the fact of anti-VEGF treatment is relatively recent and previous research on AMD and depression was conducted before the dissemination of anti-VEGF as the first line treatment for wAMD (34). Considering the positive effects of anti-VEGF to prevent further vision loss in wAMD patients, the question has been raised as to whether patients receiving anti-VEGF treatment might present a different prevalence of depression and anxiety (34,35). Existing studies in this field are still limited in generating evidence and often with low numbers (9,34). A recent

1 review on this topic (9) found four studies where the prevalence of depression was examined  
2 in patients receiving anti-VEGF treatments (22-24,36). Findings were preliminary but  
3 suggested that the wAMD patient group receiving anti-VEGF treatment still present higher  
4 rates of depression and anxiety in comparison with general population. Finally, previous  
5 research has also shown that despite depression and anxiety being common among adults  
6 with vision loss, they tend to not be tackled by health services and therefore remain untreated  
7 (36-39).

8 The primary purpose of this study is to investigate the patient experience of receiving anti-  
9 VEGF treatment for wAMD, with a particular focus on patient sources of anxiety related to  
10 anti-VEGF treatment. The secondary aim of this study is to achieve a snapshot of the  
11 frequency of clinical levels of anxiety, depression, and post-traumatic stress (PTS), among  
12 patients attending for treatment of wet AMD, and to explore levels of burden among patient  
13 carers. To the best of our knowledge there is no literature supporting the existence of any link  
14 between AMD or anti-VEGF and PTS, but we included the assessment of PTS as exploratory  
15 of any serious unmet mental health needs that could influence patients' experience of  
16 treatment. Additionally, we want to explore clinical levels of depression and anxiety between  
17 patients with different number of injections received and between patients with different  
18 visual acuity. Mental health problems in our patient group could be related to their AMD, the  
19 injection treatments, ageing, personal history, burden of travel or other internal and external  
20 factors and we do not attempt to prove any specific link between anti-VEGF and  
21 psychological symptoms. Instead we aimed to explore any existent mental health needs in our  
22 patient group, exploring clinical levels of depression and anxiety across different patient sub-  
23 groups to derive a general picture of patients' mental health and whether these needs are  
24 being met by health services. With regard to patient carers, we also wanted to achieve a  
25 snapshot of the burden experienced by carers of a patient who regularly receives anti-VEGF  
26 treatment. Our study aims were built upon two premises: the fact our patient group is  
27 receiving regular invasive treatment that has been described as potential stressful; and the fact  
28 of AMD patients are generally considered at risk of having comorbid mental health problems  
29 that tend to not be addressed by health services. Our ultimate hope is that information from  
30 this study can be used to help managing the psychological as well as ophthalmic needs of our  
31 patients in AMD clinics.

32 Literature on depression and anxiety among wAMD patients receiving anti-VEGF treatment  
33 is still very limited and scarce, which necessitates a more exploratory scope for our study.  
34 However, we hypothesise that patient anxiety related to anti-VEGF is more frequent among  
35 patients at early stages of treatment, i.e., in patients who have received up to 3 injections than  
36 in patients who are receiving treatment for longer, in light of what was suggested in previous  
37 studies (17,18). Additionally, we hypothesise that clinical levels of depression and anxiety  
38 are significantly more frequent in patients who are at early stages of treatment (who have  
39 received up to 3 injections) than in patients receiving treatment for longer, i.e., the higher the  
40 number of anti-VEGF injections received, the less frequent anxiety and depression are seen.  
41 Finally we also hypothesise that the frequency of clinical levels of depression and anxiety are  
42 negatively correlated with visual acuity, as previous studies suggested that depression is  
43 associated with poorer visual acuity in patients with vision disorders (29,30).

## 44 **Methods**

### 45 **Ethics**

46 North West Research Ethics Committee (NHS, UK) approved study procedures (reference  
47 15/NW/0288). The study adheres to the tenets of the declaration of Helsinki. We offered all  
48 participants a participant information sheet, and obtained informed consent prior to  
49 recruitment.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1 **Study Design**

2 We carried out an observational cross-sectional mixed-methods study. The qualitative strand  
3 used a structured survey to explore patient experience of receiving anti-VEGF treatment and  
4 sources of anxiety associated with this treatment. The quantitative strand of this study used  
5 standardised psychological instruments to investigate depression and anxiety, and post-  
6 traumatic stress in our sample of patients and levels of burden among patients' carers. This  
7 mixed-methods design allows us to combine the measurement of psychological health in  
8 patients undergoing treatment for wAMD (depression, anxiety and post-traumatic stress) with  
9 in-depth data (qualitative strand) that explores experience of treatment and sources of anxiety  
10 associated with treatment.  
11  
12

## 13 **Participants**

14 We invited consecutive patients with formal diagnoses of wAMD attending the macular  
15 treatment centre at Manchester Royal Eye Hospital to take part in the study. Patients and  
16 carers were recruited from May 2015 to November 2015. Inclusion criteria for patients were:  
17 to have a confirmed diagnosis of wAMD, to have received at least one anti-VEGF injection  
18 prior to the interview for this study, and to be followed-up in the macular treatment centre of  
19 the corresponding hospital. We included in our study patients who were receiving and who  
20 were not receiving psychological and psychiatric treatment. We excluded patients whose first  
21 anti-VEGF injection was on the day of interview and patients who were not able to  
22 communicate effectively. For carers the inclusion criteria was only to be carer of a patient  
23 currently receiving anti-VEGF treatment for wAMD. In our study, all researchers were  
24 blinded to carer-patient relationship, i.e., we interviewed carers without knowing who was the  
25 corresponding patient of each carer, in order to allow carers to be as open and honest as  
26 possible.  
27  
28  
29  
30  
31

## 32 **Procedures**

33 Participants were approached by an experienced clinical psychologist (HS) on the same day  
34 they were scheduled to have the medical appointment and receive treatment. All patients  
35 were invited to participate in an individual interview before receiving the injection, in a  
36 private room where all study instruments were administered. Participants were issued with a  
37 patient information sheet, and given the opportunity to ask questions about their participation.  
38 They provided written informed consent if they agreed to participate. Questionnaires and  
39 forms were read to patients who were unable to read as a result of their vision loss or having  
40 received topical eye medication. Carers of patients were approached by another researcher  
41 (NM) on the same day, following the same procedures.  
42  
43  
44  
45

## 46 **Measures for Qualitative Strand**

47 For the qualitative strand of this study, we were keen to acquire a broad range of information  
48 specific to AMD. A structured survey was developed from input provided by a patient and  
49 public involvement (PPI) group, previous literature on the topic, and considering our prior  
50 experience of dealing with AMD patients. The PPI event consisted of an informative meeting  
51 with wAMD patients, researchers and healthcare professionals who worked with these  
52 patients on a daily basis. The PPI group helped to identify key issues of patients' perceptions  
53 and concerns about anti-VEGF treatment and the way it has been delivered, which shaped the  
54 development of the survey. The PPI meeting lasted 2 hours and was audio-recorded and  
55 transcribed to identify the main and key concerns raised by participants about the treatment  
56 for wet AMD and the way the treatment is being delivered by our service. Although our  
57 survey was built upon the main topics raised by PPI participants, we have also included  
58  
59  
60  
61  
62  
63  
64  
65

1 questions addressing topics that have been highlighted in previous literature on patient  
2 experience of anti-VEGF treatment (9), such as sources of anxiety related to anti-VEGF  
3 treatment, and experiences of pain when receiving an injection. The final version of our  
4 survey comprised of 17 closed-ended questions addressing patients' demographics, clinical  
5 data and experience and perception of anti-VEGF treatment, and one open-ended question  
6 addressing patient concerns and anxieties on the anti-VEGF treatment (for more details on  
7 the survey please see appendix A). Topics covered by the survey included anxieties and  
8 concerns about anti-VEGF treatment, treatment preferences and options, doctor/patient  
9 relationship, pain experienced when receiving treatment and perception of carer's burden.

### 11 **Measures for Quantitative Strand**

12 We defined patients' visual acuity (VA) as the best-corrected VA of the eye with better  
13 vision. Best-corrected VA was converted into the logarithm of the minimum angle of  
14 resolution during analysis. We divided patients into 3 groups, based on the World Health  
15 Organization (WHO) classification of visual impairment (40): no visual impairment or mild  
16 visual impairment ( $VA \geq 6/18$ ), moderate impairment ( $6/60 < VA < 6/18$ ) and severe  
17 impairment ( $VA < 6/60$ ).

18 Depression and anxiety were assessed by the Hospital Anxiety and Depression Scale (HADS)  
19 (41). HADS is a 14-item self-report scale, comprising two subscales evaluating levels of  
20 anxiety (HADS-Anxiety) and levels of depression (HADS-Depression). The HADS has been  
21 used extensively in the hospital setting as a standardised psychological screening tool for  
22 emotional disorders (41). Each subscale includes 7 items, generating possible scores of 0 to  
23 21 for anxiety (Cronbach's  $\alpha = 0.89$ ) and depression (Cronbach's  $\alpha = 0.91$ ). A score of 8 and  
24 above is widely used to indicate the presence of clinical levels of anxiety or depression that  
25 may warrant further psychological investigation (41,42).

26 Post-traumatic stress (PTS) symptoms related to receiving an intravitreal injection was  
27 assessed by the Impact of Events Scale-Revised (IES-R) (43). IES-R is a validated measure  
28 of self-reported posttraumatic stress symptoms experienced in the past seven days in relation  
29 to a specific stressor (40). IES-R consists of 22 items, each ranging from 0 ("Not at all") to 4  
30 ("Extremely"), with the total score ranging from 0 to 88. Three symptom clusters associated  
31 with post-traumatic stress are assessed with the IES-R, including intrusion, avoidance and  
32 hyperarousal. Intrusion relates to intrusive thoughts, nightmares and imagery associated with  
33 the traumatic event (8 items; Cronbach's  $\alpha = 0.90$ ), avoidance relates to avoidance of any  
34 feelings or situations (8 items; Cronbach's  $\alpha = 0.95$ ) and hyper-arousal is an inability to  
35 concentrate, anger, irritability and hypervigilance (6 items; Cronbach's  $\alpha = 0.96$ ). Higher  
36 scores indicate more symptoms of posttraumatic stress, with scores above 24 indicating the  
37 potential need for clinical assessment and care (43).

38 The Mini-Mental State Examination (MMSE) (44) was used to assess patients' cognitive  
39 status. MMSE consists of two sections: the first section requires oral responses regarding  
40 temporal and spatial orientation, memory and attention; the second section requires the  
41 subject to name simple objects, follow verbal written commands, write a sentence  
42 spontaneously, and copy two intersecting pentagons, in order to assess language and visuo-  
43 spatial functions. MMSE final score ranges from 0 to 30 (the maximum score). MMSE  
44 scores from 25 to 30 suggest a questionably significant degree of cognitive impairment.  
45 Scores from 21 to 24 indicate mild cognitive impairment. Scores from 10 to 20 suggest  
46 moderate cognitive impairment, and scores below 10 suggest severe cognitive impairment.  
47 We adjusted some vision-related tasks to make them suitable for patients with vision  
48 impairment: for the task "writing a sentence" we helped the patient to find the paper and  
49 writing space; for the task "reading a sentence" we provided larger letters printed in bold; and  
50 for "identifying a pen and a watch" we adjusted the distance of those objects to the patients'



1 eyes. We only excluded the last MMSE task which consists of asking the patient to copy a  
2 pair of intersecting pentagons, as this task involves more complex visuospatial planning and  
3 executive skills. We anticipated that some patients would perform inaccurately in this task  
4 due to reduced visual acuity which would compromise the reliability of results. Final scores  
5 were adjusted from 0 (minimum) to 29 (maximum) because one point refers to the last test  
6 task involving the copy of a pair of pentagons. Cut-off scores were also adjusted, i.e., we  
7 used the original test cut-off scores minus 1 point. The internal consistency of the MMSE  
8 obtained Cronbach's alphas of 0.82 in elderly patients admitted to a medical service and 0.84  
9 in elderly nursing home residents (44,45).

10 The Burden Assessment Scale (BAS) (46) was used to assess burden in patients' carers. BAS  
11 is a 19-item measure that evaluates burden in an objective way, that is, the amendment of  
12 caring for someone with limitations imposed on the level of activities and resources of the  
13 caregiver. BAS employs a 4-point scale (1–4), with higher scores indicating greater levels of  
14 caregiving burden. Cronbach's alphas for the subscales and total scale were as follows: a =  
15 0.82 for limitation in activity, a = 0.64 for feelings of worry and guilt, a = 0.71 for social  
16 burden, and a = 0.81 for total scale.

## 20 **Data Analysis**

21 Results are presented mainly with descriptive statistics using SPSS software. We had no prior  
22 sample to perform power calculations. As part of pilot analyses, Chi-square test was used to  
23 investigate whether or not anxiety related to treatment is more frequent in patients who have  
24 received up to 3 injections than in patients who have received more injections, i.e., patients  
25 receiving treatment for longer. The basic assumptions for this statistical test were satisfied in  
26 that there were distinct categories and all expected frequencies were greater than 5. We  
27 created three sub-groups of patients who received up to 3 injections, i.e. patients who are at  
28 early stages of treatment (Group 1, n=21); patients who received 4 to 12 injections, i.e.,  
29 patients who are not at early stages of treatment but are more experienced with treatment and  
30 are not yet regarded as chronically treated (Group 2, n= 119); and patients who have received  
31 more than 12 injections and therefore are more likely to be chronically treated with anti-  
32 VEGF (Group 3, n=160). The choice of sub-groups was based upon patient and public  
33 involvement meetings and upon timeframes cited in previous literature indicating patient  
34 anxiety associated with earlier stages of treatment (17,18). Univariate ANOVA was used to  
35 investigate differences in HADS scores for depression and anxiety between all sub-groups of  
36 patients (G1, G2 and G3). Chi-square test was used to investigate differences in the frequency  
37 of patients with HADS scores indicating clinical levels of depression between all sub-groups  
38 of patients (G1, G2 and G3). Chi-square post-hoc analysis was performed using adjusted  
39 residuals (47) to investigate which sub-groups of patients present significant differences in  
40 number of cases of clinical levels of depression. The samples were from a number of  
41 independent groups and the variables were normally distributed with similar variance in each  
42 group. Pearson correlation was used to investigate associations between HADS scores and  
43 visual acuity (LogMAR), and between HADS scores and number of injections received.  
44 Data from the survey was analysed in terms of frequency of responses for each closed-ended  
45 question. The final open-ended question of the survey was analysed for the frequency of each  
46 source/type of anxiety related to anti-VEGF treatment that patients reported.

## 54 **Results**

### 55 *Patient Characteristics*

56 We invited 345 patients and 100 carers to participate in our study. Of 345 patients 41 refused  
57 to participate and 4 were not eligible because they were not able to communicate effectively  
58 with the interviewer. A total of 300 wAMD patients and 100 carers gave their consent and  
59  
60  
61  
62  
63  
64  
65

1 agreed to participate in this study. Patients' demographic and clinical data are presented in  
2 Table 1. Our sample of patients had a higher percentage of women (59.7%), and patients who  
3 were receiving treatment for more than 1 year (72%). The great majority of patients (85%)  
4 had good visual acuity or mild visual impairment. More than 95% of our patients were not  
5 receiving any psychological or psychiatric support at the time of the interview.  
6

7 *Please insert Table 1 here*  
8  
9

### 10 ***Qualitative Strand – Broad Experience of Receiving Anti-VEGF Treatment from Survey***

11 Patients' experience of receiving anti-VEGF treatment is presented in table 2 and table 3.  
12 Generally patients reported a positive feedback on doctor/patient relationship and the quality  
13 of care received. Most patients were satisfied with the support received from the hospital and  
14 also satisfied with the information received about the condition and treatment. Most patients  
15 also preferred to rely on the doctors to make treatment decisions. With regard to patients'  
16 treatment concerns and anxiety, we found that 132 patients (56%) reported some type of  
17 anxiety related to anti-VEGF treatment. Further analysis showed that the patients' main  
18 concerns and sources of anxiety related to anti-VEGF treatment were: fear of going blind due  
19 to intravitreal injections / fear of the needle causing damage in the eye (n=118; 39.3%); hope  
20 the treatment works properly / fear of getting worse because the treatment didn't work  
21 (n=111; 37%); waiting in the patient waiting room / anticipatory anxiety (n=104; 34.6%); and  
22 fear of the unknown in relation to treatment outcomes and disease progression (n=91; 30%).  
23 In contrast, anxiety around injection pain (n=10; 3.3%) were relatively infrequently raised.  
24 127 patients (42.3%) did report experiences of pain when receiving an intravitreal injection,  
25 but the level of pain was relatively minor with mean score of 2.4 ( $\pm 3.17$ ) on a scale from 0 to  
26 10 with 67% of patients experiencing a level of pain equal or below to 3 (scale 0 to 10).  
27 Among those patients who reported pain when receiving an injection, 42 patients (33%)  
28 experienced pain only during the initial treatment injection(s), 55 patients (43.3%)  
29 experienced pain half way through their treatment, and 30 patients (23.6%) experienced pain  
30 only during the last few injections received. More than half of patients (53.7%) reported to  
31 have expected more pain than they really experienced when receiving an intravitreal  
32 injection.  
33  
34  
35  
36  
37  
38

39 *Please insert Table 2 and Table 3 here*  
40

### 41 ***Anxiety related to Anti-VEGF treatment and Number of Injections Received***

42 Frequency of anxiety related to treatment in each patient sub-group for the number of  
43 injections received is presented in Table 4. Chi-Square showed no significant differences  
44 ( $P > .05$ ) in the frequency of anxiety related to anti-VEGF treatment between patients who  
45 have received different number of injections, i.e., between all patient sub-groups (G1, G2 and  
46 G3).  
47  
48

49 *Please insert Table 4 here*  
50  
51

### 52 ***Quantitative Strand – Clinical Levels of Depression, Anxiety, Post-traumatic stress and Carers Burden***

53 Results from the standardized measures on depression, anxiety, post-traumatic stress,  
54 cognitive function and burden are presented in Table 1. HADS-Anxiety scores indicated that  
55 17% (n=52) of all wAMD patients had clinical levels of anxiety, 10.6% (n=5) of which were  
56 receiving psychological or psychiatric treatment at the time of interview. HADS-Depression  
57 scores indicated that 12% (n=36) of all wAMD patients had clinical levels of depression, 9%  
58  
59  
60  
61  
62  
63  
64  
65

(n=3) of which were receiving psychological or psychiatric treatment for depression. Therefore 89% (n=47) of patients who showed anxiety, and 91% (n=33) of patients who showed depression were not receiving appropriate psychological and psychiatric treatment for their condition. 1 % (n=3) of patients showed symptoms of post-traumatic stress disorder according to the IES-R. Most patients (90.3%; n=271) did not exhibit impaired cognition according to MMSE. All patients were able to perform the administrated MMSE vision-related tasks successfully. Finally, according to BAS scores, 73% (n=73) of carers reported little or no burden, 19% (n=19) reported mild to moderate burden, and 8% (n=8) reported moderate to severe burden. No carers reported severe burden.

*Please insert Table 5 here*

### ***Depression and Anxiety Investigated for Number of Anti-VEGF Injections Received and Visual Acuity***

Table 5 displays HADS scores in different sub-groups of patients regarding the number of anti-VEGF injections received. HADS-Depression scores were significantly higher in patients who had received up to 3 injections (Group 1) in comparison with patients who had received from 4 to 12 injections (Group 2) (ANOVA  $P=.027$ ), and in comparison with patients who had received more than 12 injections (Group 3) (ANOVA  $P=.001$ ). No significant differences were found in HADS-Anxiety scores between the three sub-groups of patients for the number of injections received (ANOVA  $P=.22$ ).

Table 6 displays the frequency of patients with a HADS-Depression score indicating clinical levels of depression in the three sub-groups of patients (HADS-Depression $\geq$ 8). Chi-square test showed significant differences between groups ( $P=.003$ ). Chi-square post-hoc analysis indicated that the number of patients with clinical levels of depression were significantly higher in Group 1 (patients who had received up to 3 injections) ( $P=.002$ ), considering the adjusted Bonferroni corrected  $P$ -value of .008.

HADS-Depression scores were significantly but weakly correlated with number of anti-VEGF injections received ( $R=-.126$ ;  $P=.03$ ). HADS-Anxiety scores were not significantly correlated with number of injections ( $P=.325$ ). Both HADS-Anxiety ( $P=.536$ ) and HADS-Depression scores ( $P=.635$ ) were not significantly correlated with VA (LogMAR).

*Please insert Table 6 here*

### **Discussion**

The qualitative findings of the study demonstrated that patients generally reported to have a good understanding of anti-VEGF treatment, and a good relationship with healthcare professionals. Additionally, they acknowledged the need for receiving treatment and its potential benefits for their eye health. However, despite presenting positive feedback on anti-VEGF treatment, more than half of the patients reported anxiety associated with receiving anti-VEGF treatment. In addition, our findings suggest that the frequency of anxiety related to treatment is similar between patients who are at early stages of treatment, i.e., patients who had received up to 3 injections, and patients who are receiving treatment for longer. This finding rejects our hypothesis that patients anxiety related to treatment is higher in patients who have received up to 3 injections and suggests that the anxiety related to anti-VEGF might persist throughout the treatment, which is not consistent with some previous literature (17,18). A possible explanation for this finding is the fact of patients' three most frequent sources of anxiety related to treatment in our study were related to fear of going blind due to injections, concerns about vision getting worse from treatment failure and waiting in the waiting room. These sources of anxiety are known to be frequent in this patient group

1 (8,10,11) and are triggered by treatment preconceptions, and anticipatory anxiety induced by  
2 regular waiting times in the waiting area before receiving treatment (11). Further cohort  
3 studies will better clarify patient anxiety related to anti-VEGF treatment, in terms of variance  
4 throughout the treatment and identifying potential anxiety triggers.

5 Previous literature has suggested that the main sources of patient discomfort and anxiety are  
6 related to the injection procedure and the fear of a needle causing damage in the eye  
7 (6,8,10,11). Our survey acknowledges this but suggests greater patient fears of going blind  
8 either as the final route of wAMD and / or as a consequence of an unexpected adverse event  
9 occurred during the injection, which is very unlikely to occur. These fears occurred despite  
10 patients having received routine information from the medical staff about the potential  
11 benefits of anti-VEGF treatment and the safety of treatment procedures. The findings from  
12 the survey suggest patients may benefit from additional and very specific information before  
13 they start the treatment with regard to the success rates of anti-VEGF treatment in halting  
14 disease progression, the disease process and how treatment can reduce the risk of becoming  
15 blind in the future. Additionally, it would be useful for patients to be informed that the needle  
16 in the eye will not be applied in the central area of the ocular globe, i.e., the patient will not  
17 see the needle coming into the eye. Finally patients need to be better informed about low  
18 likelihood of serious problems following the treatment, and that most adverse effects can be  
19 easily manageable after the treatment. Further patient resources and/or communication skills  
20 training for health professionals would help to better inform patients.

21 The quantitative finding of the study illustrated a considerable frequency of patients with  
22 clinical levels of depression and anxiety, and low percentage of patients with symptoms of  
23 post-traumatic stress. According to epidemiologic studies on depression and anxiety (48,49),  
24 the frequency of clinical levels of anxiety and depression we found in our patient group is  
25 higher than general population, and within the range of what has been found in studies on  
26 general AMD patients (dry and wet AMD) (19). However, it is also noticeable that the  
27 frequency of patients with clinical levels of anxiety that we found in our validated  
28 questionnaires (17%) is clearly lower than the frequency of patients reporting anxiety related  
29 to treatment in our survey (56%). This difference is explained by the concept that patient  
30 anxiety related to treatment does not necessarily have clinical value or entail clinical levels of  
31 anxiety, i.e., a patient can have concerns and anxiety about the treatment but not being  
32 clinically anxious.

33 Another important issue was that most of our patients were not receiving any psychiatric or  
34 psychological treatment at the time of interview. These findings draw attention to the need of  
35 tackling mental health needs among adults with vision loss, especially in health services  
36 dedicated to provide care for ophthalmologic diseases. Previous research has highlighted the  
37 tendency for these needs to remain unaddressed (36-39), which might have a negative impact  
38 on patient healthy life-style, and eventually compromise medical treatment outcomes.

39 The study results partially reject our previous hypothesis that the frequency of clinical levels  
40 of depression and anxiety is higher at early stages of anti-VEGF treatment. We only  
41 confirmed the hypothesis that the frequency of clinical levels of depression is higher in  
42 patients who were at early stages of treatment (had received up to 3 intravitreal injections),  
43 compared to patients who had received more than 3 injections, i.e., patients who were at later  
44 stages of treatment. This hypothesis is also supported by the correlation we found between  
45 the number of injections received and HADS-Depression scores. However, this finding  
46 should be confirmed in further studies conducted on a larger sample size, as our group 1  
47 sample was small and the correlation coefficient we found was significant but weak. Previous  
48 literature on this topic is very limited and inconclusive about the frequency of depression in  
49 patients at different anti-VEGF treatment stages (9,22,28,34). One possible hypothesis to  
50 explain why depression is higher in patients at early stages of treatment is the fact of AMD is

1 often seen as a threatening condition that entails further blindness (50). Patients recently  
2 diagnosed with wAMD can therefore have that expectation of going blind. Previous  
3 qualitative and mixed-methods studies on adults with serious vision disorders suggested that  
4 the fear of further vision loss and blindness can be an important source of emotional distress  
5 (32,51,52), which can ultimately lead to depression (32). In light of this, patients at early  
6 stages of anti-VEGF treatment might be at higher risk of having depression due to their fear  
7 of going blind because of wAMD. However, as anti-VEGF treatment proceeds, patients can  
8 become more optimistic about treatment success and disease stabilization (8).

9 Patients in this study were on anti-VEGF therapy and the great majority of them had good  
10 visual acuity and did not have visual impairment according to the WHO's criteria (40).  
11 We still found that a percentage of wAMD patients are clinically depressed and anxious  
12 within the same prevalence range found in previous studies conducted with adults with visual  
13 impairment (53-55). Additionally, we rejected our hypothesis that depression and anxiety are  
14 negatively correlated with visual acuity, which is not consistent with some previous literature  
15 suggesting that patients with lower visual acuity are more likely to suffer from depression  
16 than patients with better visual acuity (29,30). However this is possibly explained by the fact  
17 of the great majority of our patients did not have visual impairment and therefore they were  
18 not likely to suffer from vision-related disability which may trigger depression (25-28).

19 Social support has been described as playing an important role in patient experience of illness  
20 and depression (31,32). Most of patient carers did not report experience of burden while  
21 being carers of someone with wAMD. This finding is consistent with the fact that most of our  
22 patients didn't perceive themselves as a source of burden for carers which might suggest that  
23 the majority of our patients had no issues with social support.

24 Our study highlights common sources of anxiety related to the process of treatment for wet  
25 macular degeneration and an awareness of these would be useful for any clinician to improve  
26 the patient experience with wAMD. It is suggested that some patient anxiety such as the fear  
27 of going blind due to injections, fear for the unknown and fear of getting worse, might be  
28 attenuated with effective communication from healthcare professionals and some counselling  
29 if needed. To ask patients about their preconceptions and expectations of treatment and  
30 prepare them for initiating anti-VEGF treatment, explaining how injections are administered  
31 and how unlikely it is to go blind because of injections, might help patients to cope with  
32 treatment. Physicians should also be aware of the prevalence of undiagnosed anxiety and  
33 depression in their AMD patients and be alert to the opportunity to manage this. Inclusion of  
34 standardized tools to screen wAMD patients for symptoms of anxiety and depression in the  
35 clinical routine of a macular treatment unit could help to better identify such patients. This  
36 should be applied irrespective of how long patients have been having injections.

37 In the current study, some limitations need to be acknowledged. This is a cross-sectional  
38 study and therefore it is more limited in providing insights into dynamic changes in clinical  
39 levels of depression and anxiety over time which would be valuable to understand these  
40 symptoms in our sample. In our study there was potential for selective bias in data collection  
41 because we were not able to identify patients who discontinued treatment after initial  
42 injection and patients with very good response (no recurrence after the first loading phase of  
43 3 injections) or very poor response to treatment (vision drop below 6/60 despite treatment),  
44 because these patients' follow-up is usually carried out outside the hospital. One of the  
45 researchers who collected data for this study (HS) also contributed to the study design and  
46 data analysis which may have entailed some bias to the results of study. Furthermore some of  
47 the researchers involved in the study have had prior clinical experience with wAMD patients,  
48 and therefore may hold preconceptions about patients' coping behaviours with treatment,  
49 which may also contribute as a source of bias. In our study the researcher were blinded to the  
50 patient-carer relationship, and did not link carers to the corresponding patients to ensure they  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 would provide a more authentic input of their experiences and perceptions of being a carer.  
2 However, a limitation of this was that we were not able to know about the potential  
3 importance of patients' social support for their experience of treatment and for the occurrence  
4 of clinical levels of depression. Finally, the authors were not able to have data on the clinical  
5 characteristics of the 41 patients who refused to participate and therefore didn't provide  
6 informed consent, and the 4 patients who were excluded because they were unable to  
7 communicate effectively with the interviewer.

8 In conclusion, this study represents one of the largest and most detailed examinations of  
9 patients undergoing anti-VEGF therapy to date. We found that although anti-VEGF  
10 treatments themselves are generally well-accepted, particular causes of anxiety exist which  
11 could potentially be reduced by improved education and minor alterations in practice. Our  
12 additional study using validated objective questionnaires demonstrated that in our examined  
13 population, a substantial percentage of patients with wAMD receiving anti-VEGF treatment  
14 had undiagnosed clinical levels of anxiety and depression, despite the good visual acuity and  
15 the promising outcomes offered by anti-VEGF treatment in preventing further visual  
16 impairment. Doctors should be vigilant to this finding that might impair patients' quality of  
17 life and consider measures to detect and address this. Further longitudinal studies and  
18 controlled trials are needed to provide a more in-depth understanding about depression and  
19 anxiety in wAMD patients and to aid development of new intervention tools, at both the  
20 patient and clinical level, aiming to reduce these symptoms and to improve the well-being of  
21 patients receiving anti-VEGF treatment.  
22  
23  
24  
25  
26

#### 27 **Abbreviations and Acronyms:**

28 **wAMD:** Wet Age-Related Macular Degeneration; **Anti-VEGF:** Vascular endothelial growth  
29 factor inhibitors; **HADS:** Hospital Anxiety and Depression Scale; **IES-R:** Impact of Events  
30 Scale Revised; **MME:** Mini-Mental State Examination; **VA:** Visual Acuity; **BAS:** Burden  
31 Assessment Scale  
32  
33  
34

#### 35 **Financial Disclosure(s):**

36 This work was supported by a grant received from Bayer – reference: R117779.

37 Hugo Senra received an educational grant from Bayer.

38 Konstantinos Balaskas is receiving Travel/educational grants from Novartis, Bayer, and  
39 Alimera.

40 Neda Mahmoodi has no conflicts of interest.

41 Tariq Aslam is receiving travel and educational grants from Bayer, Novartis, Bausch and  
42 Lomb, Alimera, Oraya, and Thea.  
43  
44  
45

#### 46 **Author Contributions:**

47 Study conception and design: Tariq Aslam, Konstantinos Balaskas, Hugo Senra

48 Analysis and interpretation: Hugo Senra, Tariq Aslam, Neda Mahmoodi

49 Data collection: Hugo Senra, Neda Mahmoodi

50 Obtained funding: Tariq Aslam, Konstantinos Balaskas

51 Overall responsibility: Tariq Aslam  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## REFERENCES:

1. Coleman HR, Chan CC, Ferris FL 3rd, Chew EY. Age-related macular degeneration. *Lancet* 2008;372(9652):1835-1845.
2. Wong WL, Su X, Li X, et al. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *Lancet Glob Health* 2014;2(2):e106-116.
3. Solomon SD, Lindsley KB, Krzystolik MG, Vedula SS, Hawkins BS. Intravitreal Bevacizumab Versus Ranibizumab for Treatment of Neovascular Age-Related Macular Degeneration: Findings from a Cochrane Systematic Review. *Ophthalmology* 2016;123(1):70-77.e1
4. Ba J, Peng RS, Xu D, et al. Intravitreal anti-VEGF injections for treating wet age-related macular degeneration: a systematic review and meta-analysis. *Drug Des Devel Ther* 2015;28(9):5397-5405.
5. Shao J, Choudhary MM, Schachat AP. Neovascular Age-Related Macular degeneration. *Dev Ophthalmol* 2016;55:125-136.
6. Boyle J, Vukicevic M, Koklanis K, Itsiopoulos C. Experiences of patients undergoing anti-VEGF treatment for neovascular age-related macular degeneration: a systematic review. *Psychol Health Med* 2015;20(3):296-310.
7. Segal O, Segal-Trivitz Y, Nemet AY, Cohen P, Geffen N, Mimousi M. Anxiety levels and perceived pain intensity during intravitreal injections. *Acta Ophthalmol* 2016; 94(2):203-204.
8. McCloud C, Khadka J, Gilhotra JS, Pesudoivs K. Divergence in the lived experience of people with macular degeneration. *Optom Vis Sci* 2014;91(8):966-974.
9. Senra H, Ali Z, Balaskas K, Aslam T. Psychological impact of anti-VEGF treatments for wet macular degeneration-a review. *Graefes Arch Clin Exp Ophthalmol*. 2016;254(10):1873-1880.
10. Henriksen K, Adhami S. Patient experience of intravitreal injections in AMD. *IJOP* 2010;1(2):68–72.
11. Thetford C, Hodge S, Harding S, Taylor S, Knox P. Living with age-related macular degeneration treatment: Patient experiences of being treated with ranibizumab (Lucentis)® intravitreal injections. *Br J Vis Impair* 2013;31(2):89–101.
12. Blaha G, Tilton E, Barouch F, Marx JL. Randomized trial of anesthetic methods for intravitreal injections. *Retina* 2011;31(3):535–539.
13. LaHood B, Sherwood D, Suter A. Comparative assessment of the effectiveness of anaesthesia for intravitreal bevacizumab injection. *Clin Exp Ophthalmol* 2011;39(2):184–185.
14. Moisseiev E, Regenbogen M, Bartfeld Y, Barak A. Evaluation of pain in intravitreal bevacizumab injections. *Curr Eye Res* 2012;37(9):813–817.
15. Rifkin L, Schaal S. Factors affecting patients' pain intensity during in office intravitreal injection procedure. *Retina* 2012;32(4):696–700.
16. Roth D, Scott I, Gulati N, Prenner J. Patient perceptions of discomfort and changes in vision and functional status associated with intravitreal triamcinolone injection. *Am J Ophthalmol* 2006;142(3):492–494.
17. Tailor R, Beasley R, Yang Y, Narendran N. Evaluation of patients' experiences at different stages of the intravitreal injection procedure – What can be improved? *Clin Ophthalmol* 2011;5:1499-1502.

18. Chua P, Mitrut I, Armbrrecht A, Vani A, Aslam T, Dhillon B. Evaluating patient discomfort, anxiety, and fear before and after ranibizumab intravitreal injection for wet age related macular degeneration. *Arch Ophthalmol* 2009;127(7):939–940.
19. Dawson SR, Mallen CD, Gouldstone MB, Yarham R, Mansell G. The prevalence of anxiety and depression in people with age-related macular degeneration: a systematic review of observational study data. *BMC Ophthalmol* 2014;14:78.
20. Eramudugolla R, Wood J, Anstey KJ. Co-morbidity of depression and anxiety in common age-related eye diseases: a population-based study of 662 adults. *Front Aging Neurosci* 2013;5:56.
21. Mathew RS, Delbaere K, Lord SR, Beaumont P, Vaegan, Madigan MC. Depressive symptoms and quality of life in people with age-related macular degeneration. *Ophthalmic Physiol Opt* 2011;31(4):375-380.
22. Casten R, Rovner BW, Leiby BE, Tasman W. Depression despite anti-vascular endothelial growth factor treatment of age-related macular degeneration. *Arch Ophthalmol* 2010;128(4):506-508.
23. Lee WJ, Cho HY, Kim DH, et al. Depression of Late Age-Related Macular Degeneration Patients in Korea. *Asia Pac J Ophthalmol* 2013;2(1):23-27.
24. Sloan FA, Hanrahan BW. The effects of technological advances on outcomes for elderly persons with exudative age-related macular degeneration. *JAMA Ophthalmol* 2014;132(4):456-463.
25. Gopinath B, Liew G, Burlutsky G, Mitchell P. Age-related macular degeneration and 5-year incidence of impaired activities of daily living. *Maturitas* 2014;77(3):263-266.
26. Walker JG, Anstey KJ, Lord SR. Psychological distress and visual functioning in relation to vision-related disability in older individuals with cataracts. *Br J Health Psychol* 2006;11(pt2):303-317.
27. Banerjee A, Kumar S, Kulhara P, Gupta A. Prevalence of depression and its effect on disability in patients with age-related macular degeneration. *Indian J Ophthalmol* 2008;56(6):469-474.
28. Casten R, Rovner B. Depression in Age-Related Macular Degeneration. *J Vis Impair Blind* 2008;102(10):591-599.
29. Wulsin LR, Jacobson AM, Rand LI. Psychosocial adjustment to advanced proliferative diabetic retinopathy. *Diabetes Care* 1993;16:1061–1066.
30. Dreer L, Elliott T, Berry J, et al. Cognitive appraisals, distress and disability among persons in low vision rehabilitation. *Br J Health Psychol* 2008;13:449–461.
31. Papadopoulos K, Papakonstantinou D, Montgomery A, Solomou A. Social support and depression of adults with visual impairments. *Res Dev Disabil* 2014;35:1734-1741.
32. Senra H, Vieira C, Nicholls E, Leal I. Depression and Experience of Vision Loss in a Group of Adults at Rehabilitation Setting: a mixed-methods pilot study. *J Rehab Res Dev* 2013;50(9):1301-1314.
33. Vink D, Aartsen MJ, Schoevers RA. Risk factors for anxiety and depression in the elderly: a review. *J Affect Disord.* 2008;106(1-2):29-44.
34. Casten RJ, Rovner BW. Update on depression and age-related macular degeneration. *Curr Opin Ophthalmol* 2013;24(3):239-243.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
35. Varano M, Eter N, Winyard S, Wittrup-Jensen KU, Navarro R, Heraghty J. The emotional and physical impact of wet age-related macular degeneration: findings from the wAMD Patient and Caregiver Survey. *Clin Ophthalmol* 2016;3(10):257-267.
  36. Rees G, Tee HW, Marella M, Fenwick E, Dirani M, Lamoureux EL. Vision-specific distress and depressive symptoms in people with vision impairment. *Invest Ophthalmol Visual Sci* 2010;51(6):2891–2896.
  37. Nyman SR, Gosney MA, Victor CR. Emotional well-being in people with sight loss. *Br J Vis Impair* 2010;28(3):175–203.
  38. Rees G, Ponczek E, Hassell JB, Keeffe J, Lamoureux E. Psychological outcomes following interventions for people with low vision: a systematic review. *Expert Rev Ophthalmol* 2010;5(3):385–403.
  39. Holloway EE, Sturrock BA, Lamoureux EL, Keeffe JE, Rees G. Help seeking among vision-impaired adults referred to their GP for depressive symptoms: patient characteristics and outcomes associated with referral uptake. *Aust J Prim Health* 2015;21(2):169-175.
  40. World Health Organization Website. Available at <http://www.who.int/blindness/Change%20the%20Definition%20of%20Blindness.pdf> Accessed 28 January 2016.
  41. Zigmond A, Snaith R. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983;67(6):361-370.
  42. Poole N, Morgan J. Validity and reliability of the Hospital Anxiety and Depression Scale in a hypertrophic cardiomyopathy clinic: the HADS in a cardiomyopathy population. *Gen Hosp Psychiatry* 2006;28(1):55-58.
  43. Weiss D, Marmar C. *The Impact of Event Scale-Revised*. New York: The Guilford Press; 1997.
  44. Folstein M, Folstein SE, McHugh PR. “Mini-Mental State” a Practical Method for Grading the Cognitive State of Patients for the Clinician. *J Psych Res* 1975;12(3):189-198.
  45. Crum RM, Anthony JC, Bassett SS, et al: Population-based norms for the Mini-Mental State Examination by age and educational level. *JAMA* 1993;269:2386–2391.
  46. Reinhard SC, Gubman GD, Horwitz AV, Minsky S. Burden assessment scale for families of the seriously mentally ill. *Eval Program Plann* 1994;17(3):261-269.
  47. Garcia-pErez MA, Nunez-Anton V. Cellwise residual analysis in two-way contingency tables. *Educ Psychol Meas* 2003;63(5):825-839.
  48. NICE (2011). Common mental health disorders | Guidance and guidelines | NICE. [online] Available at <http://www.nice.org.uk/guidance/cg123>. Accessed 25 August 2015.
  49. McManus S, Meltzer H, Brugha T, Bebbington P, Jenkins R (eds), 2009. *Adult Psychiatric Morbidity in England 2007: results of a household survey*. NHS Information Centre for Health and Social Care. [online] Available at <http://www.hscic.gov.uk/pubs/psychiatricmorbidity07>. Accessed 22 April 2015.
  50. Bennion AE, Shaw RL, Gibson JM. What do we know about the experience of age related macular degeneration? A systematic review and meta-synthesis of qualitative research. *Soc Sci Med* 2012;75:976-985.

- 1 51. Nyman SR, Dibb B, Victor CR, Gosney MA. Emotional well-being and adjustment to  
2 vision loss in later life: a metasynthesis of qualitative studies. *Disabil Rehabil*  
3 2012;34(12):971–981.
- 4 52. De Leo D, Hickey PA, Meneghel G, Cantor CH. Blindness, fear of sight loss, and  
5 suicide. *Psychosomatics* 1999;40(4):339-344.
- 6  
7 53. Evans JR, Fletcher AE, Wormald RP. Depression and anxiety in visually impaired  
8 older people. *Ophthalmology* 2007;114(2):283–288.
- 9  
10 54. Zhang X, Bullard KM, Cotch MF, et al. Association between depression and  
11 functional vision loss in persons 20 years of age or older in the United States,  
12 NHANES 2005-2008. *JAMA Ophthalmol* 2013;131(5):573-581.
- 13  
14 55. Ribeiro MV, Hasten-Reiter Júnior HN, Ribeiro EA, Jucá MJ, Barbosa FT, Sousa-  
15 Rodrigues CF. Association between visual impairment and depression in the elderly: a  
16 systematic review. *Arq Bras Oftalmol* 2015;78(3):197-201.
- 17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Table 1 – Demographical and Clinical Data

Variable		No. (%)	Range	Mean ± SD
Age			56-95	80±7.4
Gender	Female	179 (59.7%)		
	Male	121 (40.3%)		
Years of education	<10	193 (64.3%)	4-21	10.8±2.5
	≥10	107 (35.7%)		
Relationship status	Single	22 (7.3%)		
	Married	122 (40.7%)		
	Widowed	123 (41%)		
	Cohabited	7 (2.3%)		
	Divorced	36 (8.7%)		
Visual acuity (LogMAR)	Good VA / Mild visual impairment ( $\geq 6/18$ )	255 (85%)	0.02-1.0	0.6±0.26
	Moderate VA ( $6.60 \leq VA < 6/18$ )	42 (14%)		
	Poor VA ( $< 6/60$ )	3 (1%)		
Type of anti-VEGF treatment	Ranibizumab	120 (40%)		
	Aflibercept	175 (58.3%)		
	Bevacizumab	5 (1.6%)		
Elapsed time since the first anti-VEGF injection (Treatment Duration)	≤ 1 Year	84 (28%)	0-96	30.9±24
	> 1 Year	216 (72%)		
Number of injections received to date			1-66	16.6±11.6
Receiving any type of psychiatric or psychological treatment	Yes	13 (4,3%)		
	No	287 (95,7%)		
Symptoms of Clinical Depression (HADS-D)	No symptoms of depression	264 (88%)		
	Symptoms of depression	36 (12%)		
Symptoms of Clinical Anxiety (HADS-A)	No symptoms of anxiety	248 (82.7%)		
	Symptoms of anxiety	52 (17.3%)		
Symptoms of PTS (IES-R)	No symptoms of PTS	297 (99%)		
	Symptoms of PTS	3 (1%)		
Cognitive Impairment	No cognitive impairment	271 (90.3%)		

(MMES)	Mild cognitive impairment	25 (8.3%)		
	Moderate to severe cognitive impairment	4 (1.3%)		
Carers' Perception of Burden (BAS)	Little or no burden (0-20)	73 (73%)	0-55	14.4±13.07
	Mild to moderate burden (21-40)	19 (19%)		
	Moderate to severe burden (41-60)	8 (8%)		
	Severe burden (61-88)	0		

Table 2 Results from Survey about Patient Experiences of Anti-VEGF treatment

Survey Topic	Sub-Topics	No.(%)	
Anxiety related to anti-VEGF Treatment	Anxiety / concerns related to treatment not reported	132 (44%)	
	Anxiety / concerns related to treatment reported	168 (56%)	
Preferences about the treatment	Hospital Visits	Prefer having fewer hospital visits	163 (54.3%)
		Don't mind about the number of hospital visits	137 (45.7%)
	Frequency of Intravitreal Injections	Prefer receiving less frequent injections	174 (58%)
		Don't mind about the frequency of injections	126 (42%)
Expectation of Pain before start the Treatment	No pain expected	128 (42.7%)	
	Pain expected	161 (53.7%)	
Experience of Pain When Receiving an Intravitreal Injection	No painful experience	162 (54%)	
	One or more painful experiences	127 (42.3%)	
Doctor/patient relationship	Understanding of Treatment Options	Good understanding of the different treatment options	192 (64%)
		Poor understanding of the different treatment options	21 (7%)
		Have only been told about one treatment option	87 (29%)
	Making Decisions on the Eye Treatment	Prefer just rely on the doctors for making decisions on the AMD treatment	197 (65.7%)
		Prefer to be more in charge for making decisions on the AMD treatment	8 (2.7%)
		Both (to rely on the doctors and to be in charge for making decisions)	38 (31.7%)
	Contradictory	Have received contradictory	22 (7.3%)

	Information about the Condition and Treatment	information about the diagnosis and treatment	
		Did not have received contradictory information about the diagnosis and treatment	277 (92.3%)
	Satisfaction with the support received from the hospital	Some disappointment	38 (12.7%)
		Not disappointed	262 (87.3%)
	Explanation about treatments and exams	Insufficient	35 (11.7%)
		Good	265 (88.3%)
Perception of carer's burden	The carer has been burdened with hospital visits		44 (14.6%)
	The carer has not been burdened with hospital visits		130 (43.3%)
	Not applicable / no carer		126 (42%)

Table 3 Results from Survey about Patient Concerns and Sources of Anxiety Related to Anti-VEGF Treatment

<b>Concerns / Anxieties Associated with anti-VEGF Treatment</b>	<b>No.(% of total sample)</b>
Fear of going blind due to injections / fear of the needle in the eye	118 (39.3%)
Hope the treatment works properly / Fear of getting worse because the treatment didn't work (treatment effectiveness)	111 (37%)
Waiting in the waiting room / Anticipatory anxiety	104 (34.7%)
Fear of the unknown in relation to treatment outcomes and disease progression	91 (30%)
Anxiety caused by being in the eye hospital for a medical appointment, exam or eye-treatment	52 (17.3%)
Fear of pain when receiving an injection	10 (3.3%)
Fear of side effects	8 (2.6%)

Table 4 – Frequency of Anxiety related to Anti-VEGF Treatment between Sub-Groups of Patients

	G1: Up to 3 injections received (N=21)	G2: From 4 to 12 injections received (N=119)	G3: More than 12 injections received (N=160)	P-value*
Patients who reported anxiety related to treatment (N=168)	14 (66,7%)	64 (53,8%)	90 (56,3%)	.546
Patients who did not report anxiety related to treatment (N=132)	7 (33,3%)	55 (46,2%)	70 (43,8%)	

G: Sub-Group of Patients; \*Chi-Square test



Table 5 – HADS scores in Sub-Groups of Patients

	G1: Up to 3 injections received (N=21)	G2: From 4 to 12 injections received (N=119)	G3: More than 12 injections received (N=160)	<i>P</i> -value* between G1-G2	<i>P</i> -value* between G1-G3	<i>P</i> -value* between G2-G3
HADS-Anxiety (±SD)	5.1±3.9	3.8±4.1	3.56±3.51	.34	.19	.83
HADS-Depression (±SD)	5.8±4.5	3.7±3.8	2.9±2.9	.027**	.001**	.11

G: Sub-Group of Patients; VA: Visual Acuity; VI: Visual Impairment

\* Univariate ANOVA; \*\* $P < 0.05$

Table 6 – Frequency of Clinical Levels of Depression between Sub-Groups of Patients

	G1: Up to 3 injections received (N=21)	G2: From 4 to 12 injections received (N=119)	G3: More than 12 injections received (N=160)	<i>P</i> Value*
Patients with Clinical Levels of Depression (HADS-D $\geq$ 8)	7 (33,3%) <i>P</i> =.001**	16 (13,4%) <i>P</i> =.54**	13 (8,1%) <i>P</i> =.03**	.003

\* Chi-Square Test before Post-Hoc analysis; \*\* Chi-Square Post-Hoc Analysis (Adjusted Bonferroni Corrected *P*-value=.008)

**Manuscript Title:** Experience of Anti-VEGF Treatment and Clinical Levels of Depression and Anxiety in Patients with Wet Age-Related Macular Degeneration

**Table of Contents**

This mixed-methods study highlights that although anti-VEGF treatments were generally well-accepted by patients with wet AMD, most of them reported particular types of anxiety related to treatment, especially the fear of going blind due to injections. A substantial percentage of wet AMD patients receiving treatment had undiagnosed and untreated clinical anxiety and depression, despite the good visual acuity and the promising outcomes of anti-VEGF treatment. Strategies to improve practice and address patient needs were suggested.

**UNIVERSITY OF MANCHESTER  
MANCHESTER ROYAL EYE HOSPITAL  
SURVEY FOR POPPIES STUDY**

**1. DEMOGRAPHIC AND CLINICAL DATA**

Date: \_\_\_\_ / \_\_\_\_ / 2015

Patient's Reference: \_\_\_\_\_

Gender: F / M

Birth Date: \_\_\_\_ / \_\_\_\_ / \_\_\_\_\_

Marital Status: \_\_\_\_\_

Years of Education: \_\_\_\_\_

Date of AMD Diagnosis: \_\_\_\_ / \_\_\_\_ / \_\_\_\_\_

Type of Medical Treatment for AMD: \_\_\_\_\_

AMD Treatment Regimen: Ranibizumab monthly PRN/Ranibizumab treated and extended,  
/ First year Aflibercept fixed regimen/ Second year Aflibercept monthly PRN/ Second year Aflibercept  
bimonthly PRN / Aflibercept first or second year after Ranibizumab

Date of First Injection for AMD: \_\_\_\_ / \_\_\_\_ / \_\_\_\_\_

Number of Injections Received to Date \_\_\_\_\_

Visual Acuity at the Date of this Interview: \_\_\_\_\_

Pre-Morbid Medical Conditions: \_\_\_\_\_  
\_\_\_\_\_

Co-morbid Medical Conditions (Present): \_\_\_\_\_  
\_\_\_\_\_

Premorbid Depression? Yes \_\_\_\_ ; No \_\_\_\_ ; Type of treatment \_\_\_\_\_

Premorbid Other Psychiatric Conditions? Yes \_\_\_\_ ; No \_\_\_\_ ; If so, specify \_\_\_\_\_

Comorbid Other Psychiatric Conditions? Yes \_\_\_\_ ; No \_\_\_\_ ; If so, specify \_\_\_\_\_

Are you receiving psychological / psychotherapeutic treatment? Yes \_\_\_\_ ; No \_\_\_\_

**2. QUESTIONS ABOUT THE TREATMENT**

1. Do you have any particular anxieties as you wait to be seen in the waiting area?

Not at all \_\_\_\_ ; Yes sometimes \_\_\_\_ ; Yes Frequently \_\_\_\_

2. Do you find the uncertainty of whether you need an injection or not difficult to cope with?

Not at all \_\_\_\_ ; Yes sometimes \_\_\_\_ ; Yes Frequently \_\_\_\_

3. Would you prefer to have slightly fewer hospital visits?

No \_\_\_\_ ; Yes \_\_\_\_ ; No Preference \_\_\_\_

4. If possible, would you be happy with having fewer injections than you currently have?

No \_\_\_\_ ; Yes \_\_\_\_ ; No Preference \_\_\_\_

5. Do you feel you have adequate understanding of the different treatment options available for your eye problem to decide which one would be more appropriate for you?  
Yes \_\_\_\_ ; No \_\_\_\_; I was only told about one treatment option \_\_\_\_\_
6. Do you get anxious when you are asked to make decisions on the treatment for your eyes?  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_; I was never asked to make decisions on the treatment for my condition \_\_\_\_\_
7. Do you prefer relying more on your doctors for making the decisions for your care or do you prefer being fully in charge of your own care?  
I prefer relying more on my doctors \_\_\_\_ ; I prefer being in charge of my own care \_\_\_\_; Both \_\_\_\_
8. Have you experienced receiving contradictory information from the healthcare professionals about the exams or treatments you had to do / or have done?  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_\_
9. Are you disappointed with the support you have received from the hospital to get information after the treatment, booking new appointments or exams?  
Not disappointed \_\_\_\_\_ ; Disappointed \_\_\_\_\_ ; A Little Disappointed \_\_\_\_\_
10. Have you found insufficient explanation about your condition and treatments  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_\_
11. Have you ever experienced losing control when managing your attendance to medical appointments, treatment sessions, exams, etc.?  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_\_
12. Have you ever feared for your vision and becoming blind when receiving the treatment?  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_\_
13. To what extent do you think your carer(s) or accompanying person has been burdened with your hospital visits?  
Never \_\_\_\_\_ ; Sometimes \_\_\_\_\_ ; Frequently \_\_\_\_\_; Not Applicable \_\_\_\_
14. Have you experienced some financial problems related to travel to the hospital for a treatment session or a medical appointment?  
Never \_\_\_\_ ; Sometimes \_\_\_\_ ; Frequently \_\_\_\_\_
15. Have you ever experienced pain when receiving injections?  
Yes \_\_\_\_ ; No \_\_\_\_; If so please rate - from no pain (0) to the maximum level of pain (10) \_\_\_\_\_
16. When did you experience more pain?  
In the first injection \_\_\_\_ ; In the first injections \_\_\_\_ ; Middle of treatment \_\_\_\_ ; Last one (s) \_\_\_\_
17. Before you start the treatment did you expect more pain when receiving injections than you really experienced?  
Yes \_\_\_\_; No \_\_\_\_
18. What is/are your greatest concern(s) / source(s) of anxiety with regard to your treatment?  
Please identify \_\_\_\_\_