



LEEDS
BECKETT
UNIVERSITY

Citation:

Read, D and Till, K and Beasley, G and Clarkson, M and Heyworth, R and Lee, J and Jones, B (2017) Peak running intensities in rugby union match-play: comparisons to whole match data. In: 40th Annual Conference of National Strength and Conditioning Association, 12 July 2017 - 15 July 2017, Las Vegas, NV, USA.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3703/>

Document Version:

Conference or Workshop Item (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

PEAK RUNNING INTENSITIES IN RUGBY UNION MATCH-PLAY: COMPARISONS TO WHOLE MATCH DATA

Peak running intensities in comparison to whole match data

Dale Read^{1,2}, Kevin Till^{1,2}, Grant Beasley³, Michael Clarkson⁴, Rob Heyworth⁴, Josh Lee⁴, Ben Jones^{1,2,5}

¹Leeds Beckett University, UK

²Yorkshire Carnegie Rugby Union Football Club, UK

³Rugby Football Union, UK

⁴Catapult Sports, Australia

⁵Rugby Football League, UK

Purpose: Data from global positioning systems (GPS) such as relative distance ($\text{m}\cdot\text{min}^{-1}$) is typically presented as an average from the whole match. The purposes of the study were to establish the peak running intensities of rugby union match-play and secondly, to make comparisons between the peak intensities and a whole match average.

Methods: With ethical approval, 125 rugby union players were recruited from 6 English rugby union academies (age: 17.6 ± 0.6 years; stature: 182.3 ± 12.2 cm; body mass: 92.6 ± 12.9 kg). Players were divided into forwards ($n = 70$) and backs ($n = 55$). One game from each academy during the 2016 / 2017 season was assessed with each player wearing a 10 Hz GPS unit (S5 Optimeye, Catapult Sports) during the match. The file was trimmed so that it only contained data from the match and analysed using a customised spreadsheet (Microsoft Excel). Peak values for relative distance were calculated for different moving averages (1, 5, 10 min) alongside the average for the whole match. For example, a 1 min moving average analysed 600 consecutive data points and the highest value for the duration was recorded for each player. Data were analysed using Cohen's d effect sizes (ES) and classified as *trivial* (0.00-0.19); *small* (0.20-0.59); *moderate* (0.60-1.19); *large* (1.20-1.99); *very large* (2.00-3.99) and *extremely large* (>4.00).

Results: The difference between average relative distance and peak 1 min was *extremely large* for forwards (73.6 ± 8.4 vs. 157.3 ± 18.1 $\text{m}\cdot\text{min}^{-1}$, ES = 10.00) and backs (79.8 ± 10.5 vs. 174.9 ± 23.9 $\text{m}\cdot\text{min}^{-1}$, ES = 9.02). The difference between average relative distance and peak 5 min was *very large* for forwards (98.5 ± 15.1 $\text{m}\cdot\text{min}^{-1}$, ES = 2.97) and backs (109.2 ± 14.5 $\text{m}\cdot\text{min}^{-1}$, ES = 2.79). The difference between average relative distance and peak 10 min was *large* for forwards (84.8 ± 11.6 $\text{m}\cdot\text{min}^{-1}$, ES = 1.33) and backs (94.8 ± 11.8 $\text{m}\cdot\text{min}^{-1}$, ES = 1.43). The difference between 1 and 5 min peak was *large* for forwards (ES = -1.74) and backs (ES = -1.71). The difference between 5 and 10 min peak was *moderate* for forwards (ES = -0.94) and backs (ES = -0.96).

Conclusions: This study highlights that peak running intensities are substantially higher than whole match averages. Additionally, the intensity decreases with the duration of time. It provides sport and position specific intensities that players should be exposed to during training for a range of durations, which prepares athletes for the most demanding phases of play.

Practical Applications: The quantification of peak running intensities allows the conditioning coach to prescribe high intensity running drills during training. The study provides evidence that coaches should not use whole match data for conditioning purposes. In team sports where the play is intermittent in nature, the use of whole match data is likely to underestimate the demands of match-play.

Figure 1. Peak average relative distance ($\text{m}\cdot\text{min}^{-1}$) of rugby union players for whole match, 1, 5 and 10 min durations