



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Till, KA and Morley, D and O'Hara, J and Jones, B and Chapman, C and Beggs, CB and Cooke, CB and Cobley, S (2017) A retrospective longitudinal analysis of anthropometric and physical qualities that associate with adult career attainment in junior rugby league players. *Journal of Science and Medicine in Sport*, 20 (11). pp. 1029-1033. ISSN 1440-2440 DOI: <https://doi.org/10.1016/j.jsams.2017.03.018>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/3706/>

Document Version:

Article (Accepted Version)

---

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

1 **A retrospective longitudinal analysis of anthropometric and physical qualities that associate**  
2 **with adult career attainment in junior rugby league players**

1 **Abstract**

2 *Objectives:* To retrospectively compare the longitudinal physical development of junior rugby league  
3 players between the Under 13 and 15 age categories in relation to their adult career attainment  
4 outcome.

5 *Design:* Retrospective longitudinal design

6 *Methods:* Fifty-one former junior rugby league players were retrospectively grouped according to  
7 their career attainment outcome as adults (i.e., amateur, academy or professional). As juniors, players  
8 undertook a physical testing battery on three consecutive annual occasions (Under 13s, 14s, 15s)  
9 including height, body mass, sum of four skinfolds, maturation, vertical jump, medicine ball chest  
10 throw, 10-60 m sprint, agility 505 and estimated  $\dot{V}O_{2max}$ .

11 *Results:* Future professional players were younger than academy players with a greater estimated  
12  $\dot{V}O_{2max}$  compared to amateur players. Between Under 13s and 15s, professional players ( $5.8 \pm 2.5$  cm)  
13 increased sitting height more than amateur ( $4.4 \pm 2.1$  cm) and academy ( $4.1 \pm 1.4$  cm) players. Logistic  
14 regression analyses demonstrated improvements in sitting height, 60m sprint, agility 505 and  
15 estimated  $\dot{V}O_{2max}$  between amateur and professional players with a high degree of accuracy  
16 (sensitivity = 86.7%, specificity = 91.7%).

17 *Discussion:* Findings demonstrate that the development of anthropometric, maturational and physical  
18 qualities in junior rugby league players aged between 13 and 15 years contributed to adulthood career  
19 attainment outcomes. Results suggest that age, maturity and size advantages, commonly observed in  
20 adolescent focused talent identification research and practice, may not be sensitive to changes in later  
21 stages of development in order to correctly identify career attainment. Practitioners should identify,  
22 monitor and develop physical qualities of adolescent rugby league players with long-term athlete  
23 development in mind.

24

25 **Key Words:** Talent identification, talent development, maturation, adolescence

## Introduction

Sport national governing bodies and professional clubs invest considerable resources (e.g., time, finances, staff) to identify and develop young talented athletes in the hope that they may become the professionals of tomorrow<sup>1</sup>. Many organisations implement talent identification practices designed to recognise current participants who have the potential to excel in particular sport contexts<sup>2</sup>. This has led to an increased research interest in the process of talent identification over the last decade, with research in this field having been undertaken in a wide variety of sports (e.g., rugby union<sup>3</sup>, rugby league<sup>4</sup>, Australian Rules Football [AFL]<sup>5</sup>). Although talent identification research and practice is now common, it is often limited by the cross-sectional nature of the methodologies used<sup>2</sup>. While these methodologies have merit, they tend to assume that current adolescent performance can be used to predict outcomes in adulthood, an approach which fails to consider the dynamic nature of athlete development, which is impacted upon by factors such as growth and maturation<sup>2,6,7</sup>.

Longitudinal and retrospective research methodologies are two approaches that can address the limitations of cross-sectional methods and detect athlete change over time<sup>8</sup>. Longitudinal designs involve data collection on the same individuals for two or more time periods<sup>9</sup> to track and evaluate performance change. Longitudinal approaches have been utilized in rugby league<sup>10</sup> and rugby union<sup>11</sup> to account for the role developmental factors (e.g., growth and maturation) play in physical performance during adolescence. For instance, Till et al.<sup>10</sup>, demonstrated large inter-individual differences and changes in physical performance characteristics of 13-15 year old athletes; with later maturing players demonstrating greater performance improvement than earlier maturing athletes (e.g., 60m sprint, Early = -0.46s; Later = -0.85s<sup>10</sup>). These findings highlight the value of longitudinal tracking in understanding physical development variability and dynamics in youth athletes.

Retrospective designs use an athlete's future career attainment and retrospectively analyse individuals cross-sectional data from an earlier timepoint<sup>12</sup>. This methodology allows researchers and practitioners to identify characteristics, potentially within adolescent athletes, that may be important for future sporting success<sup>12</sup>. Previously, in talent development research, such designs have helped determine the physical characteristics of adolescent athletes associated with adult career attainment (i.e., amateur or professional) in soccer<sup>13,14</sup> and rugby league<sup>12,15</sup>. For example, Till et al.<sup>12</sup> reported

1 that future professional rugby league players had lower sum of skinfolds and advanced fitness  
2 characteristics at 13-15 years of age compared to those who attained amateur status with no  
3 differences in characteristics observed between future professional and academy level players.

4 In the emerging research field of talent identification and athlete development, any study,  
5 which combines longitudinal and retrospective methodological approaches, should in theory yield  
6 new insights that are important. Although prior studies have implemented longitudinal (e.g., <sup>7,10</sup>) and  
7 retrospective (e.g., <sup>12,15</sup>) research methodologies, no study to the authors knowledge has combined  
8 these two methods to evaluate the developmental changes in adolescent athletes that are associated  
9 with future career attainment outcomes. Therefore, the aim of this study was to retrospectively  
10 compare the longitudinal physical development of adolescent athletes (i.e., junior rugby league  
11 players) between the Under 13 and 15 age groups in relation to their long-term career attainment  
12 outcome.

13

14

#### Methods

15

16

17

18

19

20

21

22

23

24

25

26

27

Fifty-one male junior rugby league players (mean age 13.59±0.25 years at Under 13s) who were selected to the Rugby Football League's former talent identification and development programme, the Player Performance Pathway (PPP; see <sup>16</sup> for a detailed description), participated in the study. Players were selected to the PPP by rugby coaches using subjective decisions based on current performance and future potential <sup>16</sup>. Players in the present study were selected to the PPP on three consecutive annual occasions in 2005 (Under 13s), 2006 (Under 14s) and 2007 (Under 15s). All players were then subsequently tracked in relation to their career attainment in July 2008 and August 2015. By 2008, participants were classified as either: (a) selected to join a professional rugby league club's academy; (b) continued to play amateur rugby league; or (c) no longer participating in rugby league. By 2015, players were potentially able to progress into playing senior professional rugby league within the European Super League. Consistent with previous investigations <sup>12,15</sup>, for the purposes of this study, players were classified into three career attainment outcomes, (1) not selected to an academy squad and classed as 'amateur' (n=12); (2) selected to a professional 'academy' but did

1 not play Super League (n=24), and (3) played 'professional' rugby league by the 2015 Super League  
2 season (n=15).

3 All participants were assessed on their physical qualities in July 2005, 2006 and 2007. All  
4 protocols received institutional ethics approval with consent provided by both players and  
5 parents/guardians. The physical assessment included standard anthropometry (height, sitting height,  
6 body mass, sum of 4 skinfolds), maturation (age at peak height velocity; PHV) and fitness (lower and  
7 upper body power, speed, change of direction speed, estimated  $\dot{V}O_{2max}$ ) characteristics. Intraclass  
8 correlation coefficients and typical error measurements for each measure have been presented  
9 previously<sup>4,17</sup> and all measurement reliability and objectivity conformed to published expectations<sup>18</sup>.

10 Height, sitting height, body mass and sum of four skinfolds were collected in the morning in a  
11 fasted state with participants wearing only shorts. Height and sitting height were measured to the  
12 nearest 0.1cm using a Seca Alpha stadiometer. Body mass was measured to the nearest 0.1kg using  
13 calibrated Seca alpha (model 770) scales. The sum of four skinfold thickness was determined using  
14 calibrated Harpenden skinfold callipers (British Indicators, UK) with procedures in accordance with  
15 Hawes & Martin<sup>19</sup>. An age at PHV prediction equation was used to measure maturity status<sup>20</sup>. Years  
16 from PHV (YPHV) were calculated by subtracting age at PHV from chronological age.

17 A standardised warm-up was conducted prior to fitness testing with tests performed in the  
18 following order. Running speed was assessed over 10m, 20m, 30m and 60m using timing gates  
19 (Brower Timing Systems, IR Emit, USA) recorded to the nearest 0.01s from three trials, separated by  
20 3 minutes rest. Change of direction speed was assessed using the agility 505 test<sup>21</sup>. Three attempts  
21 were performed on each foot with times recorded to the nearest 0.01s. A vertical jump test was used to  
22 assess lower body power using a Takei vertical jump metre (Takei Scientific Instruments Co. Ltd,  
23 Japan). A counter-movement jump was performed with hands positioned on hips, with jump height  
24 measured to the nearest cm from three trials separated by 30 seconds rest<sup>22</sup>. Upper body power was  
25 assessed using the 2kg medicine ball (Max Grip, China) chest throw<sup>23</sup>. Participants were seated with  
26 their backs against a wall and were instructed to throw the ball horizontally as far as possible.  
27 Distance was measured to the nearest 0.1cm from the wall to where the ball landed with the furthest

1 of three trials used as the score. Maximal oxygen uptake ( $\dot{V}O_{2max}$ ) was estimated using the multistage  
2 fitness test<sup>24</sup>. Players were required to run 20m shuttles keeping in time with a series of beeps.  
3 Player's running speed increased progressively until they reached volitional exhaustion. Regression  
4 equations were used to estimate  $\dot{V}O_{2max}$  from the level reached during the multistage fitness test.

5 Mean and standard deviation (SD) scores were calculated for all dependent variables  
6 according to age category and career attainment outcome. A repeated measures multivariate analysis  
7 of variance test (MANOVA) was initially conducted to identify significant main effects for time  
8 between age category, for group according to career attainment outcome, and whether an age category  
9 x career attainment outcome interaction existed. Bonferroni pairwise comparisons were then  
10 conducted to examine univariate effects between each dependent variable. Partial eta squared ( $\eta^2$ )  
11 effect sizes were also calculated and interpreted as 0.01 = small, 0.06 = medium and 0.14 = large<sup>25</sup>.  
12 All analyses were conducted with SPSS version 21.0 with significance levels set at  $p < 0.05$ .

13 To assess the physical qualities that best predicted career attainment outcome, binomial  
14 logistic regression analysis was performed using the changes in physical qualities between the age  
15 categories, with career attainment outcome coded as a binary variable (1 = Professional, 0 =  
16 Amateur). These groups were selected because binomial logistic regression can only cope with two  
17 groups, and these groups represented opposite ends of the player spectrum. Analysis was performed  
18 using in-house algorithms written in 'R' (open source statistical software), with all study variables  
19 included in the initial model. Variable selection was undertaken using a step-wise approach, with  
20 variables excluded if non-significant. The general applicability of the predictive logistic regression  
21 models was tested using 10-fold cross-validation. To maximize sensitivity (true positive rate) and  
22 specificity (true negative rate), receiver operating characteristic (ROC) analysis was used to calculate  
23 optimum cut-off values.

24

25

## Results

26

27

Table 1 shows the physical qualities at each age category according to career attainment  
outcome. Table 2 shows the repeated measures MANOVA univariate analyses and pairwise

1 comparison results according to age category, career attainment outcome and the age category x  
2 career attainment outcome interactions.

3 \*\*\*Insert Table 1 near here\*\*\*

4 \*\*\*Insert Table 2 near here\*\*\*

5 Repeated measures MANOVA analyses identified significant main effects for age category  
6 ( $F=6120.7$ ,  $p<0.001$ ,  $\eta^2=1.00$ ) and all dependent variables. Pairwise comparisons showed all variables  
7 significantly improved across the three annual-age categories except sum of four skinfolds and agility  
8 505 left and right.

9 For career attainment outcome, analyses identified significant main effects ( $F=2.12$ ,  $p=0.005$ ,  
10  $\eta^2=0.48$ ) with significant differences found for chronological age and estimated  $\dot{V}O_{2max}$ . Pairwise  
11 comparisons found professional players were younger than academy players, and professional and  
12 academy players had a greater estimated  $\dot{V}O_{2max}$  than amateur players across the three age categories.

13 For age category x career attainment outcome interactions, analyses identified significant  
14 main effects ( $F=1.66$ ,  $p=0.049$ ,  $\eta^2=0.72$ ) with significant differences found for sitting height, 10m and  
15 20m sprint. Greater improvements in sitting height were found for professional ( $5.8\pm 2.5$  cm)  
16 compared to amateur ( $4.4\pm 2.1$  cm) and academy ( $4.1\pm 1.4$  cm) between Under 13s and 15s. For 10m  
17 and 20m sprint, professional ( $-0.09\pm 0.07$ ;  $-0.16\pm 0.10$  s) and amateur ( $-0.08\pm 0.06$ ;  $-0.19\pm 0.13$  s)  
18 players demonstrated greater improvements than academy ( $-0.05\pm 0.06$ ;  $-0.09\pm 0.11$  s) players.

19 Logistic regression analysis revealed that physical changes between Under 13s and 14s in  
20 YPHV ( $\beta=-34.320$ ,  $p=0.029$ ), sitting height ( $\beta=4.564$ ,  $p=0.025$ ) and body mass ( $\beta=1.309$ ,  $p=0.031$ )  
21 contributed to a predictive model (LR model 1) with a cross-validation accuracy of 88.9%. Between  
22 Under 14s and 15s, 10m sprint ( $\beta=22.225$ ,  $p=0.025$ ) contributed to a model (LR model 2) with 66.7%  
23 cross-validation predictive accuracy. Between Under 13s and 15s, sitting height ( $\beta=-0.896$ ,  $p=0.024$ ),  
24 60m sprint ( $\beta=-6.199$ ,  $p=0.032$ ), agility 505 left ( $\beta=-8.060$ ,  $p=0.045$ ) and estimated  $\dot{V}O_{2max}$  ( $\beta=-$   
25  $0.431$ ,  $p=0.025$ ) contributed to a model (LR model 3) with 81.5% cross-validation predictive  
26 accuracy. Table 3 shows the results of the ROC analysis for the respective logistic regression models.  
27 The models were able to distinguish with a high degree of accuracy at the Under 13s to 14s



1 (sensitivity = 93.3%, specificity = 91.7%;  $p < 0.001$ ) and Under 13 to 15s (sensitivity = 86.7%,  
2 specificity = 91.7%;  $p < 0.001$ ) between the future professional and amateur players.

3 \*\*\*Insert Table 3 near here\*\*\*

## 4 5 **Discussion**

6 Originality in the current study is highlighted by the longitudinal, retrospective research  
7 design that allowed for changes in physical qualities between Under 13s-15s to be evaluated against  
8 adult career attainment outcome. The longitudinal development of physical qualities in a sample of  
9 junior rugby league players selected to a talent development program on three consecutive occasions  
10 (i.e., Under 13s, 14s, 15s) was related to adult career attainment outcome (i.e., amateur, academy or  
11 professional). Findings demonstrated that future professional players were chronologically younger  
12 than academy players, and had a greater estimated  $\dot{V}O_{2\max}$  than amateur players. Future professional  
13 players increased sitting height more than academy and amateur players. Further, amateur and  
14 professional players improved 10m and 20m sprint performance more than academy players. Logistic  
15 regression analysis demonstrated that anthropometric and maturational characteristics differentiated  
16 between career attainment outcome between Under 13s and 14s age categories and 10m sprint  
17 between Under 14s and 15s. The development of sitting height, speed, change of direction speed and  
18 estimated  $\dot{V}O_{2\max}$  differentiated between career attainment outcome between Under 13s and 15s.

19 When physical qualities were compared between career attainment level across the Under  
20 13s-15s age categories, significant differences were found for chronological age and estimated  
21  $\dot{V}O_{2\max}$ . Future professional players were found to be younger, both chronologically and relatively,  
22 than the future academy players, supporting the findings of previous research in rugby league <sup>15</sup>,  
23 rugby union <sup>26</sup> and ice hockey <sup>27</sup>. Furthermore, this suggests that relatively younger athletes selected  
24 to a talent identification and development programme during adolescence tend to achieve greater  
25 success in future career attainment. As such, this suggests that perceived advantages (e.g., selection  
26 opportunities associated with increased age within chronological annual-age groups) may not be  
27 advantageous for longer-term future career attainment in a sport context.

1 Estimated  $\dot{V}O_{2\max}$  was found to be greater in academy and professional players compared to  
2 amateur players across the Under13-15s age categories; a finding consistent with previous cross-  
3 sectional <sup>4</sup> and retrospective <sup>12,15</sup> research in rugby league. Such findings suggest that enhanced  
4 aerobic power during adolescence may contribute to an increased career attainment in rugby league.  
5 Interestingly, unlike previous cross-sectional <sup>4</sup>, longitudinal <sup>10</sup> and retrospective <sup>12,15</sup> studies in rugby  
6 league, no other physical qualities (e.g., vertical jump, speed, agility 505) demonstrated a significant  
7 difference according to career attainment level. The reduced sample size, compared to previous  
8 investigations <sup>12,15</sup> and large inter-player variability may have led to no significant differences in these  
9 physical qualities.

10 Findings also demonstrated that sitting height improved to a greater extent in professional,  
11 compared to amateur and academy players. This suggests that future professional players were more  
12 likely to mature later due to the relationship between maturational status and development of sitting  
13 height <sup>20,28</sup>. This is further supported by the moderate effects (although not significant) found for  
14 changes in height and body mass, with greater gains in professional than amateur players (i.e., height,  
15 professional = 9.0, amateur = 6.7 cm; body mass, professional = 15.7, amateur = 12.0 kg). These  
16 findings are also consistent with prior rugby league <sup>15</sup> and soccer <sup>14</sup> studies, highlighting how earlier  
17 maturation in adolescence does not necessarily translate into advanced career attainment outcomes,  
18 irrespective of selection to a talent identification and development programme. Later maturers may in  
19 fact have a greater likelihood of career attainment success, possibly due to greater potential for  
20 improvement <sup>7,10</sup> and/or due to the required development of other technical, tactical or psychological  
21 factors in more challenging environments <sup>29</sup>. The assessment of anthropometric characteristics and  
22 maturation status within adolescent players could be considered to allow potential dispensation  
23 criteria to supplement age grade grouping to provide greater participation and selection opportunities  
24 for later maturing players as proposed in rugby union <sup>30</sup>.

25 Recent research in AFL <sup>5</sup> has used logistic regression analysis to determine the characteristics  
26 important for talent identification. Findings from our application of logistic regression analysis  
27 suggest that the development of anthropometric and maturational characteristics is highly influential

1 in future career attainment in rugby league. Using a combination of logistic regression and ROC  
2 analysis, we were able to use changes in YPHV, sitting height and body mass between Under 13s and  
3 14s to correctly identify 93.3% of the future professionals and 91.1% of amateur players. Thus, the  
4 development of body size in players selected into the talent identification programme seems important  
5 for career attainment, particularly between 13 and 14 years of age which is the timing of maturation  
6 <sup>28</sup>. 10 m speed improvement was identified as the only variable discriminating professional and  
7 amateur players between Under 14 and 15 age categories with a sensitivity and specificity score  
8 suggest that it may be an important variable. However, between Under 13s and 15s the development  
9 of sitting height, 60m sprint, agility 505 and estimated  $\dot{V}O_{2max}$  all contributed to logistic regression  
10 model and accurately distinguished between future professional and amateur players. This analysis  
11 provides evidence that the improvement of a range of physical qualities contributes to successful  
12 career attainment in rugby league as previously suggested <sup>4,12,15</sup> but may be limited by the collinearity  
13 of some measures. On this basis, practitioners should aim to monitor maturation, alongside the  
14 monitoring and development of anthropometric and physical qualities, within adolescent rugby league  
15 players to support talent identification and development practices.

16

## 17 **Conclusions**

18 In summary, findings showed that physical qualities and the rate of development in  
19 anthropometric, maturational and physical qualities of junior rugby league players aged between 13  
20 and 15 years contributed to future career attainment (i.e., professional levels). Younger and later  
21 maturing individuals selected to the talent development programme between Under 13s and 15s age  
22 groups appeared to have greater likelihood of attaining professional levels. Likewise, players with  
23 advanced estimated  $\dot{V}O_{2max}$ , may have a greater likelihood of higher career attainment outcome. The  
24 development of sitting height, speed, change of direction speed and estimated  $\dot{V}O_{2max}$  during  
25 adolescence appear to be important factors for future career attainment outcomes and practitioners  
26 should identify, monitor and develop such physical qualities of adolescent rugby league players with  
27 long-term athlete development in mind.

1 **Practical Implications**

- 2 • Practitioners should understand that advanced age and earlier maturation within chronological  
3 annual-age groups might not be an accurate indicator of longer-term future career attainment.
- 4 • Advanced physical qualities, particularly estimated  $\dot{V}O_{2\max}$ , of adolescent rugby league  
5 players may contribute to long-term career attainment, and could be more carefully  
6 considered in talent identification and development practices.
- 7 • The systematic training and development of physical qualities including speed, change of  
8 direction speed and estimated  $\dot{V}O_{2\max}$ , should be emphasised in adolescent player  
9 development to increase the likelihood of higher level career attainment.

10  
11 **Acknowledgements**

12 This research was supported by the Rugby Football League (RFL) and the authors would like  
13 to thank the RFL for providing the data. There was no financial assistance associated with this  
14 research.

## References

1. Cobley S, Baker J, Schorer J. Identification and development of sport talent: A brief introduction to a growing field of research and practice, Chapter 1 in *Talent identification & development in sport: International perspectives*, Baker J, Cobley S, Schorer J editors, London, Routledge, 2012.
2. Vaeyens R, Lenoir M, Williams AM, et al. Talent identification and development programmes in sport: Current models and future directions. *Sports Med* 2008; 38(9): 703-714.
3. Pienaar AE, Spamer MJ, Steyn HS. A scientific approach towards the identifying of rugby talent among ten and eleven year-old boys. *Kinesiology* 1998; 28(1): 48-53.
4. Till K, Cobley S, O'Hara J, et al. Using Anthropometric and Performance Characteristics to Predict Selection in Junior UK Rugby League Players. *J Sci Med Sport* 2011; 14(3): 264-269.
5. Woods C, Raynor AJ, Bruce I, et al. Predicting playing status in junior Australian Football using physical and anthropometric parameters. *J Sci Med Sport* 2015; 18(2): 225-229.
6. Subotnik RF, Olszewski-Kubilius P, Worrell FC. Rethinking Giftedness and Gifted Education: A Proposed Direction Forward Based on Psychological Science. *Psychological Science* 2011; 12(1): 3-54
7. Cobley S., Till, K., O'Hara, J. et al. Variable and changing trajectories in youth athlete development: Further verification in advocating a long-term inclusive tracking approach. *J Strength Cond Res* 2014; 28(7): 1959-1970.
8. Cobley S, Till K. Longitudinal tracking of athlete development: Its importance, methods & future considerations. . In *The Handbook of Talent Identification and Development in Sport*, Baker J, Cobley S, Schorer J, Wattie, N. editors, London, Routledge, in press.
9. Falk B, Lidor R, Lander Y, et al. Talent identification and early development of elite water polo players: a 2 year follow up. *J Sports Sci* 2004; 22(4): 347-355.
10. Till K, Cobley S, O'Hara J, et al. Considering maturation and relative age in the longitudinal evaluation of junior rugby league players. *Scand J Sci Med in Sport* 2014; 24(3): 569-576.
11. Spamer EJ, Hare E. A longitudinal study of talented youth rugby players with specific reference to skill, growth and development. *J Hum Move Stud* 2001; 41(106): 39-57

- 1 12. Till K, Cobley S, Morley D, et al. Retrospective analysis of anthropometric and fitness  
2 characteristics associated with long-term career progression in rugby league. *J Sci Med Sport*  
3 2015; 18(3): 310-314.
- 4 13. Emmonds S, Till K, Jones B, et al. Anthropometric, speed and endurance characteristics of  
5 English academy soccer players: Do they influence obtaining a professional contract at 18 years  
6 of age? *Int J Sports Sci Coaching* 2016; 11(2): 212-218.
- 7 14. Le Gall F, Carling C, Williams M, et al. Anthropometric and fitness characteristics of  
8 international, professional and amateur male graduate soccer players from an elite youth  
9 academy. *J Sci Med Sport* 2010; 13(1): 90-95.
- 10 15. Till K, Cobley S, Morley D, et al. The influence of age, playing position, anthropometry and  
11 fitness on career attainment outcomes in rugby league. *J Sports Sci* 2016; 34(13): 1240-1245.
- 12 16. Till K, Cobley S, Wattie N, et al. The prevalence, influential factors and mechanisms of relative  
13 age effects in UK rugby league. *Scand J Sci Med Sport* 2010; 20: 320-329.
- 14 17. Till K, Cobley S, O'Hara J, et al. Anthropometric, physiological and selection characteristics in  
15 high performance UK junior Rugby League players. *Talent Development Excellence* 2010; 2(2):  
16 193-207.
- 17 18. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med* 2000; 30: 1-15.
- 18 19. Hawes MR, Martin AD. Human Body Composition, chapter 1 in *Kinanthropometry and exercise*  
19 *physiology laboratory manual: Tests, procedures and data second edition. Volume 1:*  
20 *Anthropometry*. London, Routledge, 2001
- 21 20. Mirwald RL, Baxter-Jones ADG, Bailey DA, et al. An assessment of maturity from  
22 anthropometric measurements. *Med Sci Sport Exerc* 2002; 34: 689-694.
- 23 21. Gabbett TJ, Herzig PJ. Physiological characteristics of junior elite and sub-elite rugby league  
24 players. *Strength Cond Coach* 2004; 12: 19-24.
- 25 22. Hunter JP, Marshall, RN. Effects of power and flexibility training on vertical jump technique.  
26 *Med Sci Sports Exerc* 2002; 34(3): 478-486.

- 1 23. Stockbrugger BA, Haennel RG. Contributing factors to performance of a medicine ball explosive  
2 power test: a comparison between jump and non jump athletes. *J Strength Cond Res* 2003; 17:  
3 768-774.
- 4 24. Ramsbottom R, Brewer J, Williams C. A progressive shuttle run test to estimate maximal oxygen  
5 uptake. *Br J Sports Med* 1988; 22: 141–144.
- 6 25. Cohen, J. *Statistical power analysis for the behavioural sciences*, 2nd ed. Hillsdale, 9 NJ.  
7 Erlbaum. 1988
- 8 26. McCarthy N, Collins D. Initial identification and selection bias versus the eventual confirmation  
9 of talent: evidence for the benefits of a rocky road? *J Sports Sci* 2014; 32: 1604-1610.
- 10 27. Deaner R, Lowen A, Cogley S. Born at the wrong time: Selection bias in the 11 NHL Draft.  
11 *PLoS One* 2013; 8(2): 1-7. 12
- 12 28. Malina RM, Bouchard C, Bar-Or O. *Growth, Maturation, and Physical Activity* (2nd ed.).  
13 Champaign, IL: Human Kinetics, 2004.
- 14 29. Cogley S, Baker J, Wattie N, et al. Annual age-grouping and athlete development: a meta-  
15 analytical review of relative age effects in sport. *Sports Med* 2009; 39: 235–256.
- 16 30. Patton DA, McIntosh AS, Denny G. A review of the anthropometric characteristics, grading and  
17 dispensation of junior and youth rugby union players in Australia. *Sports Med* 2016; 46(8): 1067-  
18 1081.

19  
20  
21  
22  
23

**Table 1:** Anthropometric and physical qualities of players selected to the PPP at Under 13s, 14s and 15s age categories according to career attainment outcome

	Amateur (n=12)			Academy (n=24)			Professional (n=15)		
	U13s	U14s	U15s	U13s	U14s	U15s	U13s	U14s	U15s
Age (years)	13.59±0.24	14.59±0.24	15.59±0.24	13.71±0.12	14.71±0.12	15.71±0.12	13.42±0.31	14.42±0.31	15.42±0.31
Age at PHV (years)	13.34±0.59	13.53±0.66	13.60±0.58	13.51±0.48	13.57±0.40	13.71±0.44	13.38±0.62	13.44±0.50	13.55±0.37
Years PHV (years)	0.24±0.58	1.06±0.66	1.98±0.58	0.20±0.50	1.13±0.42	2.00±0.45	0.04±0.70	0.98±0.55	1.90±0.48
Height (cm)	171.4±6.7	175.0±6.1	178.1±5.0	170.5±4.6	174.7±4.7	177.3±5.0	170.6±7.9	176.6±5.8	179.6±4.2
Sitting Height (cm)	87.2±4.4	89.2±4.5	91.3±4.0	86.6±3.5	89.3±2.8	91.0±2.7	85.6±4.2	89.3±3.2	91.4±2.2
Body Mass (kg)	65.4±12.4	70.1±12.3	77.4±11.4	62.6±7.6	69.5±9.0	76.2±10.4	63.0±11.4	71.6±10.6	78.7±10.3
Skinfolds (mm)	41.4±20.3	44.5±17.4	46.2±19.0	35.8±14.8	35.4±16.2	42.3±18.2	33.4±13.7	37.4±14.3	36.8±13.3
Vertical Jump (cm)	37.5±4.5	39.3±3.3	41.3±3.9	38.6±4.7	42.2±4.2	43.5±4.9	38.7±4.3	41.3±3.9	43.9±5.4
MBT (m)	5.4±0.8	5.8±0.8	6.4±0.9	5.4±0.5	5.9±0.4	6.5±0.5	5.3±0.8	6.0±0.6	6.7±0.5
10m (s)	1.97±0.09	1.95±0.09	1.89±0.08	1.94±0.06	1.91±0.07	1.89±0.07	1.95±0.09	1.88±0.10	1.86±0.10
20m (s)	3.41±0.18	3.34±0.15	3.22±0.12	3.32±0.13	3.23±0.11	3.22±0.15	3.34±0.15	3.22±0.15	3.18±0.14
30m (s)	4.81±0.26	4.67±0.23	4.50±0.17	4.67±0.20	4.50±0.17	4.44±0.18	4.66±0.23	4.49±0.21	4.38±0.22
60m (s)	9.17±0.60	8.62±0.51	8.27±0.36	8.75±0.44	8.28±0.32	8.19±0.39	8.69±0.49	8.33±0.41	8.09±0.42
Agility 505 L (s)	2.60±0.13	2.52±0.13	2.48±0.23	2.50±0.15	2.48±0.12	2.46±0.12	2.56±0.12	2.47±0.10	2.41±0.11
Agility 505 R (s)	2.61±0.18	2.53±0.16	2.52±0.19	2.51±0.16	2.46±0.12	2.53±0.14	2.57±0.13	2.48±0.11	2.43±0.09
Estimated $\dot{V}O_{2max}$ (ml.kg <sup>-1</sup> .min <sup>-1</sup> )	45.5±7.2	45.7±5.4	47.9±4.6	47.7±5.9	51.8±4.5	52.2±5.3	48.6±3.8	50.6±3.7	53.7±2.9



**Table 2:** Repeated Measures MANOVA results examining annual-age category, career attainment outcome (and interaction) on anthropometric and physical qualities.

	Annual-Age Category			Career Attainment Outcome			Annual-Age Category x Career Attainment Outcome	
	P	$\eta^2$	Pairwise	P	$\eta^2$	Pairwise	P	$\eta^2$
Age (years)	<0.001	1.00	13s<14s<15s	0.002	0.24	Acad>Pro	0.50	0.03
Age at PHV (years)	<0.001	0.30	13s<14s<15s	0.648	0.02		0.54	0.03
Years PHV (years)	<0.001	0.97	13s<14s<15s	0.704	0.01		0.45	0.04
Height (cm)	<0.001	0.80	13s<14s<15s	0.720	0.01		0.10	0.09
Sitting Height (cm)	<0.001	0.81	13s<14s<15s	0.949	0.00		0.049	0.11
Body Mass (kg)	<0.001	0.87	13s<14s<15s	0.850	0.01		0.06	0.09
Skinfolds (mm)	<0.001	0.11	13s<15s	0.369	0.04		0.21	0.06
Vertical Jump (cm)	<0.001	0.40	13s<14s<15s	0.284	0.05		0.67	0.02
MBT (m)	<0.001	0.71	13s<14s<15s	0.805	0.01		0.32	0.05
10m (s)	<0.001	0.41	13s>14s>15s	0.368	0.04		0.023	0.11
20m (s)	<0.001	0.51	13s>14s>15s	0.226	0.06		0.026	0.12
30m (s)	<0.001	0.62	13s>14s>15s	0.098	0.09		0.26	0.05
60m (s)	<0.001	0.66	13s>14s>15s	0.070	0.11		0.075	0.09
Ag 505 L (s)	0.001	0.15	13s>14s	0.183	0.07		0.32	0.05
Ag 505 R (s)	0.014	0.10	13s>14s	0.210	0.06		0.075	0.09
Estimated $\dot{V}O_{2\max}$ (ml.kg <sup>-1</sup> .min <sup>-1</sup> )	<0.001	0.23	13s<14s<15s	0.006	0.19	Amat<Acad, Pro	0.19	0.06

Table 3. Results of the ROC analysis using the logistic regression model outcome predictions

	<b>Logistic regression model</b>	<b>Area under curve</b>	<b>Cut-off value</b>	<b>True positives</b>	<b>False negatives</b>	<b>True negatives</b>	<b>False positives</b>	<b>Sensitivity (%)</b>	<b>Specificity (%)</b>	<b>P value</b>
<b>Age range: U13 – U14</b>										
Professional vs amateur	LR model 1	0.956	0.372	14	1	11	1	93.3	91.7	<0.0001
<b>Age range: U14 – U15</b>										
Professional vs amateur	LR model 2	0.778	0.521	11	4	9	3	73.3	75.0	0.0010
<b>Age range: U13 – U15</b>										
Professional vs amateur	LR model 3	0.928	0.577	13	2	11	1	86.7	91.7	<0.0001

