

Citation:

Surr, CA and Gates, C and Irving, D and Oyebode, J and Smith, SJ and Parveen, S and Drury, M and Dennison, A (2017) Effective dementia education and training for the health and social care workforce: A systematic review of the literature. Review of Educational Research. ISSN 0034-6543 DOI: https://doi.org/10.3102/0034654317723305

Link to Leeds Beckett Repository record: https://eprints.leedsbeckett.ac.uk/id/eprint/3892/

Document Version: Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please contact us and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

RER

Review of Educational Research

Effective dementia education and training for the health and social care workforce: A systematic review of the literature

Journal:	RER
Manuscript ID	RER-16-Jun-MS-134.R3
Manuscript Type:	Manuscript



Running head: EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Effective Dementia Education and Training for the Health and Social Care Workforce:

RER

A Systematic Review of the Literature

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Abstract

Ensuring an informed and effective dementia workforce is of international concern, however there remains limited understanding of how this can be achieved. This review aimed to identify features of effective dementia educational programmes. Critical Interpretive Synthesis underpinned by Kirkpatrick's return on investment model was applied. One hundred and fifty-two papers of variable quality were included. Common features of more efficacious educational programmes included the need for educational programmes to: be relevant to participants' role and experience; involve active face-to-face participation; underpin practice-based learning with theory; be delivered by an experienced facilitator; have a total duration of at least eight hours with individual sessions of 90 minutes or more; support application of learning in practice; and provide a structured tool or guideline to guide care practice. Further robust research is required to develop the evidence base; however, the findings of this review have relevance for all working in workforce education. *Key words* Alzheimer's, staff training, workforce development, critical interpretive synthesis

RER

 EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Effective Dementia Education and Training for the Health and Social Care Workforce: A Systematic Review of the Literature

The care of people with dementia is of global concern (Alzheimer's Disease International, 2010; WHO/Alzheimer's Disease International, 2012). People with dementia account for two-thirds of UK care home residents (Knapp et al., 2007) and occupy around one quarter of acute hospital beds (Alzheimer's Society, 2009) and have more hospital and skilled nursing facility stays and home health care visits in the US than older people generally (Alzheimer's Association, 2015). However, care quality concerns have been raised, including the adequacy of workforce skills and knowledge to provide effective care (Department of Health, 2009; US Department of Health and Human Services, 2013). In the UK, policy initiatives (Department of Health, 2009, 2012) have aimed to address this skills gap, leading to increases in dementia education and training provision. However, there remains limited available evidence of education and training efficacy. Concern with the effective transfer of knowledge developed within educational settings, to the workplace and in particular the realistic connection of theory to practice, has been a growing concern within educational research (Smith, 2003). However, to date limited attention has been given to the specific professional development needs of the dementia workforce and effective approaches to ensuring this skills gap is met. Similar initiatives to increase specialist skills or clinical expertise into the broader workforce occur frequently in healthcare e.g. mental health, end-oflife care (Department of Health, 2016). Therefore, a review of the existing evidence base, to inform future dementia workforce development initiatives, is required. Similar issues with workforce skills gaps are found in many sectors including manufacturing, technology and services, food and agriculture (Creaser, Hargreaves, & Ong, 2014; Dychtwald, Erikson, & Morison, 2006) internationally and thus the results are likely to have applicability to broader, professional development and adult, life-long and workplace education.

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Background

There has been increasing emphasis placed on ongoing knowledge and skills development of workforces, with the emergence of knowledge based economies (Dall'Alba & Sandberg, 2006). The health and social care workforce is diverse in its previous access to and experience of post-secondary education. The majority of the dementia care workforce is unqualified, low paid, low status and has no clear career path. In the UK and internationally within social care particularly, there are low levels of literacy and numeracy and many staff have English as a second, or additional, language (All-Party Parliamentary Group on Dementia, 2009; Wilson, 2014), in part due to an increasing reliance on migrant workers (Bettio, Simonazzi, & Villa, 2006; Bourgeault, Atanackovic, Rashid, & Parpia, 2010; Cangiano & Shutes, 2010). Conversely, health professions are predominantly degreequalified roles, but specialisms may provide limited provision of, or access to, dementia specific education. This diversity within the workforce of existing dementia knowledge and previous educational exposure has required employers to engage with provision of workplace learning opportunities. While Further and Higher education institutions have met some of this demand, a significant amount of dementia training provision is provided in-house or by private providers.

RER

In England currently there is no requirement for continuing professional development education or training to be accredited, and a nationally agreed sector framework for dementia educational content and learning outcomes was only published in October 2015 (Health Education England Skills for Health and Skills for Care, 2015). A similar picture is apparent internationally with recent publication of competency or education and training frameworks (Health and Social Care Board, 2016; Traynor, Cumming, & Britten, 2015; Wales, 2016), but little further support or provision for regulation or quality monitoring. As a result the content

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

and quality of dementia training and education in England are variable and low levels of dementia knowledge remain commonplace (Royal College of Psychiatrists, 2011). The government targets for numbers of NHS staff trained on dementia (Department of Health, 2013, 2014) may, in some cases, have led to a volume rather than quality or efficacy driven approach. Therefore, greater understanding and consideration of what effective dementia education and training for this workforce entails, is required. This is particularly important if this imperative to educate the workforce is to lead to improved outcomes for people with dementia. The challenges experienced with dementia care workforce development mirror those found in development of the workforce across other professional sectors, including teaching, business and the social and natural sciences (Webster-Wright, 2009). While there are differences between the roles of those providing health and social care to people with dementia and other professions, for example teaching, there are also many parallels of relevance to the continuing professional development of these professional groups including, the diversity of education of the workforce (i.e. qualified teachers/health professional vs unqualified staff/assistants), working within an organisational context and culture governed by external guidelines and quality standards and assurance, team-based working and the need to translate theoretical and knowledge-based learning into real-world practice. In the practice setting both may be under pressure to adopt a task focussed approach in order to achieve targets in a time and resource pressured environment and in what is an unpredictable setting, with complex group dynamics. Opfer and Pedder (Opfer & Pedder, 2011) have argued that teacher education must be conceptualised as a complex system, with various dynamics at the individual (teacher), meso (institutional) and macro (school system) levels at play in influencing under what conditions teachers learn, why and how. Thus we argue that health and social care workforce education too, should be conceptualised as a complex system, with

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

many facets at the individual, meso and macro levels at play that must be understood in understanding learning processes.

RER

The effective transfer of theoretical education to practice settings has received much attention in educational practice and research (Allen & Wright, 2014; Kessels & Korthagen, 1996) and has been a particular concern within the initial training and ongoing professional development of the health and social care workforce (Corlett, 2000; Jackson, Bluteau, & Furlong, 2013; Landers, 2000; Scully, 2011; Upton, 1999). In particular, effective methods to bridge the gap between the desired practices taught in the classroom and the reality of working in real-life health and social care practice have challenged those delivering health education (Jackson et al., 2013). Attempts to address the perceived gap within health and wider educational research have included a focus on the design and content of educational programmes, the knowledge and skills of the educator/teacher and teaching and learning processes utilised (i.e. delivery methods and assessment strategies) (Gallagher, 2004), the use of critical reasoning skills and reflection by learners who must be supported to be proactive in building bridges between theory and practice (Hatlevik, 2011; Severinsson, 1998; Wilkinson, Smallidge, Boyd, & Giblin, 2015) and use of simulation (Roth, Mavin, & Dekker, 2014; Wall, Andrus, & Morrison, 2014). However, little is known about whether such methods also apply to dementia education and practice in the context of workplace learning.

Workplace learning has been conceptualised in a variety of ways, which recognise the need to acquire appropriate attitudes, conceptual knowledge and practical skills that lead to the development of practical competencies (Dall'Alba & Sandberg, 2006), and research suggests that specific forms of learning may be preferred by learners located in the workplace (Smith, 2003), as opposed to other types of education. In particular, Smith argues learners have a reducing requirement for a proximal guide/facilitator and a greater need for interaction and construction as their expertise grows. Dreyfus's (1982) model of five stages to skill

 RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

development (Stage 1: novice to Stage 5: expert), applied to nurse education by Benner (1982), highlights the role of both knowledge and experience in guiding appropriate practice and decision-making. It connects theoretical knowledge with practical experience in task performance, with novice practitioners being characterised by being unable to deviate from taught rules to guide action and experts being those who are able to work holistically drawing on both formal educational preparation and acquired experience to inform practice. Thus, expertise is context specific and more advanced skills can only be achieved through experience in real-world situations (Dall'Alba & Sandberg, 2006).

The diversity of the dementia care workforce means those requiring workplace education may be at varying stages of proficiency with regard to exposure to both clinical and dementia education and practice. Some learners may, for example, have a degree or postgraduate qualification in a specialist area of clinical practice (e.g. nursing, physiotherapy, radiography, medicine) and many or few years' clinical experience, but limited exposure to specialist care of people with dementia. Other learners may work daily in specialist dementia care, but have had limited previous exposure to formal clinical or practice education (e.g. social care workers, nursing assistants/auxiliaries). This has implications for educational approaches adopted, since learners' preferred teaching and learning methods and those required for effective learning may differ depending on degree of learner expertise (Smith, 2003). This can be particularly challenging for educators who may need to provide professional development to groups of learners who span a range of levels of expertise, whether this is in health and social care education or professional development and workplace learning in another profession. Therefore, an understanding of teaching and learning methods that motivate learners and which support attitude and practice change is required.

Despite the body of educational theory and research on professional and workplace education, understanding of what constitutes effective education and training for the dementia

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

workforce is poorly understood and seldom considered when developing programmes. To date a number of systematic reviews have examined the evidence base underpinning dementia education and training interventions for the health and social care workforce. However, they are limited by their focus on: only one aspect of efficacy (Elliott, Scott, Stirling, Martin, & Robinson, 2012); a single pedagogical approach (Brody & Galvin, 2013; Fossey et al., 2014); efficacy within a single workforce group or setting (Alushi, Hammond, & Wood, 2015; Beeber, Zimmerman, Fletcher, Mitchell, & Gould, 2010; Kuske et al., 2007; Perry et al., 2011; Tullo & Allan, 2011); or a single aspect of care (Eggenberger, Heimerl, & Bennett, 2013; McCabe, Davison, & George, 2007; Raymond et al., 2014; Spector, Orrell, & Goyder, 2013; Zientz et al., 2007). Overall existing reviews consistently note the generally poor quality of existing research and variability in results, with some reported benefits of education and training particularly in areas such as staff knowledge and attitudes but no consistency across studies.

Kirkpatrick's (1979, 1984) 'Return on Investment' model developed specifically for evaluation of training consists of four levels, and is widely adopted in the evaluation of training and education provision (Bates, 2004).

- Level 1: Learner's *reaction* to and satisfaction with, the programme;
- Level 2: The extent to which *learning* has occurred including knowledge, skills, confidence and attitude change;
- Level 3: The extent to which staff *behaviour* or practices have changed as a result of completing the programme;
- Level 4: The *results* or outcomes that have occurred because of training.

Criticisms of Kirkpatrick's model have been presented including: being over simplified and incomplete in its understanding of the transfer of learning to practice; implying a hierarchy of evidence with behavioural and outcome change deemed more important to

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

measure than reaction; assuming that each level of the model is associated with the previous and following level, without evidence to support this; and a lack of empirical testing (Giangreco, Carugati, & Sebastiano, 2008; Holton, 1996; Tamkin, Yarnall, & Kerrin, 2002). Whilst Kirkpatrick's model has been subject to such critique, it remains a widely applied approach to considering the levels at which it is helpful to evaluate training, in order to understand potential return on investment.

Brody and Galvin (2013) highlight that much of the published dementia educational research fails to evaluate training impact across more than one level within Kirkpatrick's framework, meaning any relationship between levels, for example between changes to staff knowledge and outcomes for people with dementia, is not understood. All of the systematic reviews on dementia education and training conclude further research is required. While some reviews identify features present in efficacious training programmes, none examine these across each of the four Kirkpatrick levels of evaluation or combine them to provide an overall, wider picture of the ingredients most likely to lead to efficacy.

Aims

The aims of this review were to identify the factors associated with effective dementia education and training for health and social care staff, across service settings in order to draw out the implications for those involved in dementia education and more broadly for all with an interest in workforce education and training.

Method

The review protocol is registered with the PROSPERO international prospective register of systematic reviews (CRD42015027475).

Search Strategy

Protocol

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Search strategies were agreed by the authorship team based on keywords developed from initial scoping searches. The following databases were searched: MEDLINE, PsycINFO, CINAHL, AMED, British Education Index, Education Abstracts, ERIC (EbscoHost), The Cochrane Library-Cochrane reviews, Economic evaluations, CENTRAL (Wiley), HMIC (Ovid), ASSIA, IBSS (Proquest), Conference Proceedings Citation Indexes (Web of Science). Searches consisted of a combination of text words and subject headings for the following themes: Dementia/Alzheimer's, training/education, staff knowledge and patient outcomes. Reference lists of key papers and e-alerts were used to include additional papers published between search completion and the end of November 2015.

Procedure

All hits were downloaded into Endnote software, where duplicate entries were removed. Papers were excluded based on an initial title screen for relevance (see Figure 1), followed by abstract review of potentially relevant papers and finally full paper review of remaining articles. Inclusion criteria were studies written in English and published between 2000 and April 2015, whose focus was research evaluating a dementia education or training programme. Exclusion criteria were conference abstracts, Masters Dissertations and training programmes for family caregivers or people with dementia. Data extraction from all relevant papers was completed using a standard template in Excel format (see Tables S1 and S2 for data extraction headings).

At the full text review/data extraction stage, to ensure included papers were relevant to answering the review aims, further inclusion criteria were added. They were: study reports on primary research, evaluates a dementia training programme or pedagogical approach to delivery of the training, is delivered to staff working in health or social care settings and reports on at least one of Kirkpatrick's (1984) four levels of training evaluation.

Analysis

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Data analysis was conducted using Critical Interpretive Synthesis (CIS) (Dixon-Woods, Cavers, Agearwal, et al., 2006), a non-traditional systematic review method that draws on systematic qualitative enquiry, incorporating interpretive approaches. CIS permits synthesis of large amounts of diverse literature. CIS is particularly useful when studies to be reviewed use different research methods, stem from a range of disciplines and where the review is intended to inform generation of theory, evidence-based practice and decisionmaking. CIS involves a process of synthesis of the evidence, involving understanding studies in relation to themselves and other studies through thematic and conceptual comparison (Kangasniemi, Kallio, & Pietila, 2014). The product of this synthesis is not aggregation of data, as in a traditional systematic review, but the generation of new knowledge in the form of overarching themes, grounded in the studies included in the review (Dixon-Woods, Cavers, Agerwal, et al., 2006). Another feature of the review process is the selection of papers that are most relevant to explaining or theory building around the concept(s) within the review, rather than on selection of papers utilising a specific methodology or methods. Therefore, exclusion of studies on the basis of quality is not undertaken, except in the case of those deemed to be 'fatally flawed' (Dixon-Woods, Cavers, Agerwal, et al., 2006). Given the expected diverse nature of the evidence base surrounding effective dementia education and training and the review aims to develop new knowledge in the form of overarching themes, regarding the most effective approaches to adopt, CIS was chosen as the most appropriate review method. The analytic process and knowledge building was underpinned by Kirkpatrick's (1984) four-level model for the evaluation of training interventions.

Quality Review

All included papers were subject to a quality review using an adapted version of criteria developed by Caldwell et al (Caldwell, Henshaw, & Taylor, 2005) and the Critical Skills Appraisal Programme (Critical Appraisal Skills Programme (CASP), 2014) with a

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

maximum possible quality score of 14 (see Table 3). In line with CIS, this was not used to exclude papers but to provide a description of quality of the evidence base, as part of the analytic process. Papers were provided with an overall quality rating of high (score 11-14), medium (score 6-10) or low (score \leq 5). Papers with an overall rating of 'high' necessarily met all or the majority of the reporting/quality criteria to at least an adequate level and thus a degree of trust can be attributed to their findings. Papers with a rating of 'low' met few of the reporting or research quality criteria and therefore, their results, while potentially interesting or relevant require treating with some caution. An initial sample of 15 papers was rated independently for quality, using the adapted criteria, by two of the authors CS and CG. The ratings were compared, disagreements discussed and an agreed rating decided for each paper. Following this a further five studies were independently rated by both authors and compared to assess for inter-rater agreement, which was achieved to a satisfactory level (1 point or less difference) on all papers. Following this CS and CG each undertook quality ratings for half of the remaining papers.

Results

In total 152 papers were included in the review (see Figure 1). A table of included studies can be found in supplementary Table S1 and further details on training approaches, quality and Kirkpatrick outcomes are reported in supplementary table S2 (online only). **Summary of Studies**

Setting and participants. Table 1 provides an overview of the 152 included papers. The majority originated from the USA (38%) and United Kingdom (20%). Training programmes were predominantly delivered to staff working in Care Homes (49%) and were aimed at Nurses (34%) and Nursing Assistants/Aides (37%).

Methodology and quality. The majority of studies adopted a quantitative methodology (63%). One-third (34%) were rated as high quality, just over half (52%)

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

medium quality and 14% were rated as low quality. Across all three study designs (qualitative, quantitative and mixed methods), the largest proportion of studies were rated as moderate quality. Despite parity of quality criteria across all study designs, quantitative studies had the largest proportion rated as high quality (38% of all quantitative studies) and the lowest rated as low quality (11%). Qualitative studies had the lowest overall proportion rated as high quality (32%).

A breakdown of the individual quality criteria scores by overall quality rating of studies (low, moderate and high) is provided in figure 2. Low quality studies were particularly poor at: providing details of ethics issues and considerations, with over 80% of studies scoring 0 on this criterion; using appropriate data analysis techniques or reporting on the analysis techniques used (70+% of studies scoring zero); and recruiting an appropriate sample size, or providing adequate detail of the sample (65+% scoring 0). However, over 40% of moderate quality studies scored zero on adequate reporting of ethical issues and less than 15% of studies scored a two on appropriate sample size and reporting of sample characteristics, and reliability and validity of data collection methods used. Moderate quality studies scoring two on this criterion. High quality studies were characterised by each criteria being awarded a score of two for 50% or more of studies. However, over 40% of high quality studies, scored one for an appropriate sample size and reporting of sample characteristics, and 8% scored a zero for adequate reporting of ethical issues.

Teaching and learning methods adopted. Small or large group face-to-face delivery was the most common approach to training/educational delivery, adopted in 120 (79%) studies (Table 2). This approach included a diverse range of teaching and learning methods including didactic lectures, discussion, video clips, exercises and activities. In 45%

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

of studies, classroom teaching alone was used and in 34% of studies, this was used alongside other teaching methods.

Staff Reactions to Training

Seventy-four of the 152 papers (49%) reported on learners' reactions to the training; 54% used a quantitative methodology, 18% qualitative and 28% mixed methods. The quality of studies varied, with a greater proportion of low quality studies (23%) and fewer high quality studies (25%) reporting reactions than in the overall sample. Of the 74 papers, 73% reported wholly positive staff reactions to the training and 1% a predominantly negative learner response. Both positive and negative aspects of the training were reported in 22% and 4% of studies compared two or more approaches to training or education. A variety of methods for assessing reaction was used across the studies from informal discussions and semi-structured interviews to questionnaires with open or fixed response questions. These methods provided either a broad and sometimes blunt approach or a narrow and somewhat restrictive focus for learner feedback. In some cases, the restrictions of the method of gathering data limited the aspects of satisfaction learners were able to report, to those in which the trainer(s)/educator(s) were interested. However, other studies provided substantial detail on areas of the training that learners had found particularly beneficial or unhelpful. Therefore, due to some of the biases in questions asked, some caution has to be used when interpreting these results.

Four themes, drawn from 34 of the studies, of which the majority were moderate or high quality (88%), emerged from our CIS:

- 1. Relevance of training to learners' role and practice
- 2. Teaching and learning approaches adopted
- 3. The quality of the training materials
- 4. Trainer/educator qualities

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Relevance and applicability to role and practice. The perceived applicability of training to a participant's role and practice was an important factor in whether learners felt completion was beneficial. Studies indicated that training content, such as scenarios and case examples, must be relevant to the setting in which learners worked and to their role. Staff were unlikely to make efforts to apply learning in practice where training was perceived to lack relevance (Alnes, Kirkevold, & Skovdahl, 2013; Pereles et al., 2003). This finding supports assumptions within established models of adult learning (Knowles, 1970; Knowles, Holton, & Swanson, 2015; Taylor & Hamdy, 2013) and evidence from research on learning motivation in adult education more broadly (De Rijdt, Stes, van der Vleuten, & Dochy, 2013; Sogunro, 2015)

Teaching and learning approaches. Studies indicated that learners across a range of professional backgrounds, and at a range of levels, shared a preference for training that included interactive group work. Learners also indicated they valued training combining classroom-based theory with in-service/practice-based or experiential learning. Where e-learning was utilised, learners stated a combination of individual study with opportunities for on-line or face-to-face discussion were preferable. While the flexibility of e-learning was cited as a benefit, the requirements for specialist technical support and the time demands for learners and facilitators of engaging regularly with on-line discussions, which they felt were particularly beneficial to learning, meant this was suggested to be a resource intensive form of study, if done well.

A positive learning experience tended to be reported in studies employing written, video/DVD or role-play based case examples or vignettes, or through involving carers in delivery of training. They were particularly useful when they demonstrated good and poor practices for learners to reflect on and discuss. Opportunities within training to demonstrate and practise skills through in-service/practice-based learning were valued. However, the use

Page 16 of 55

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

of role-play or simulated learning could be found distressing by learners, particularly where this was employed during an individual session where there had not been time to develop a trusting relationship between learners and facilitator.

Quality of training materials. Learners identified the quality of training materials as important. Higher quality materials were clear, easy to follow, informative and unambiguous, used straightforward language and were concise. This was particularly important for learners who were learning in a second or additional language.

The perceived attractiveness of learning activities to learners has been highlighted within the theory of reasoned action, as an important antecedent for learners in their decisionmaking process regarding intention to participate in workplace learning (Kyndt & Baert, 2013). Therefore, the relevance of the programme, teaching and learning approaches used, quality of the materials and feedback from peers who have already attended the training about their perceptions of these elements, may all impact on future uptake of offered training.

Trainer skills and qualities. Studies highlighted the need for a skilled trainer/facilitator. Trainer/facilitator qualities identified included: the ability to create a comfortable environment where learners were encouraged to ask questions; being knowledgeable about the topic; listening to participants; being able to tailor training to the individual needs of a group; being able to identify and address concerns and needs quickly; and responding positively and skilfully to challenge unhelpful learner views and attitudes. This finding adds to a growing body of research in workplace training and education, that not only should effective trainers/facilitators be experienced and skilled in facilitating training with adult learners (e.g. Gauld & Miller, 2004); a concept which has long-since been established. It also supports evidence that trainers/facilitators should be able to use their own vocational knowledge and interpersonal skills to be "trainee-centred" (Arghode & Wang, 2016).

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

A number of authors have recently suggested that by using their knowledge of a relevant subject area, and by being able to tailor learning to a given learner group and their roles, trainers are in turn more effective (Arghode & Wang, 2016) and produce higher trainee satisfaction (Ghosh, Satawadi, Joshi, Ranjam, & Singh, 2012). This has especially been found to be the case in healthcare education (Khamarko, Koester, Bie, Baron, & Myers, 2012). Therefore, when combined with this growing body of evidence, our findings suggest that organisations (especially, but not limited to, healthcare organisations) could benefit from considering both previous facilitation experience and knowledge in the subject area, when recruiting or selecting trainers.

Learning

Learning-related outcomes such as knowledge, attitudes, confidence, perceived competence, and self-efficacy were reported in 109 studies, 28 of which (26%) used mixed methods, 64 (59%) were quantitative studies and the remaining 17 (16%) employed a qualitative design. Of these papers, 80% reported wholly positive outcomes for learning, 15% reported mixed outcomes with positive effects found in some areas and not in others and 6% reported no change in outcomes. The quality of the studies was largely moderate to good with only 14% rated as low quality, 55% as moderate quality and 31% as high quality. Methods of data collection included validated measures, non-validated questionnaires, surveys or scales, interviews, focus groups and analysis of documents such as assignments or written assessments. Of the 92 studies employing a mixed-methods or quantitative design, 51 (55%) used at least one validated measure of learning. While use of qualitative self-report or non-validated methods of assessing learning gains presents validity issues with regard to interpretation of findings, such data does offer a perspective on whether health and social care staff perceive they have learned anything. In some cases, staff may feel there have or

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

have not been learning gains, whether or not this is not seen on validated measures. This can have positive or negative impacts on motivation to change behaviour or practice. However, some learning gains may be difficult to quantify and thus measure, for example improved relationships with people with dementia or greater knowledge of individual life history and its use in day-to-day care. Therefore, whilst qualitative studies must be treated with caution in terms of their indication of learning gains, they retain value in understanding staff perceptions of changes in their knowledge, skills and confidence to deliver complex dementia care, particularly in difficult to measure areas.

Our analysis revealed indicators associated with the efficacy of teaching methods for learning generally, which are discussed first. Three additional sub-categories of components of learning were identified and analysed separately: knowledge or understanding; attitudes and beliefs; and confidence, competence and self-efficacy, which are discussed subsequently in this section.

Teaching methods. While caution needs to be employed in interpreting the results, due to the variable study quality, predominantly positive findings across all studies and the low numbers of studies evaluating some teaching methods, the evidence suggested some teaching methods were consistently successful or unsuccessful. In-service/practice-based learning when used as the sole training approach, watching an individual or group DVD and learning through reading a written resource (hard copy or electronic) were not effective in the majority of studies utilising these approaches. The latter two approaches represent passive learning styles, and do not reflect the best practice, active learning approaches, advocated for effective learning across the spectrum of educational research (Darling-Hammond et al., 2008; Freeman et al., 2014; Jensen, Kummer, & Godoy, 2014; Mansouri & Lockyer, 2007; Pulsford, Jackson, O'Brien, Yates, & Duxbury, 2013; Zaher & Ratnapalan, 2012).

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

A small number of studies compared efficacy of different training approaches for learning, allowing comparison beyond a 'no training' baseline or control. Generally, they found that active learning was more successful than passive approaches, for example classroom and on-line multi-media methods, were more effective than learning through reading written materials and multi-media on-line learning was more effective than passively watching a video lecture of the same content. However, in one study (Vollmar et al., 2010) no significant differences were found between learning through didactic lecture compared to multi-media on-line learning. There was a completion rate of only 50% for on-line learning, therefore, suggesting that passive face-to-face learning may be more useful by virtue of higher attendance from staff. Four studies compared face-to-face combined with in-service learning, to face-to-face learning alone (Resnick, Cayo, Galik, & Pretzer-Aboff, 2009) and in-service learning alone (Corwin, Owen, & Perry, 2008; McCabe et al., 2014; Smythe et al., 2014). All found in-service learning alone was less effective than either face-to-face delivery alone, or in-service learning in combination with face-to-face delivery. This indicates that theoretical content is required alongside, or to underpin, in-service learning, for effective learning to take place. In addition, the broader service-learning literature indicates that reflection forms an important component of in-service or practice-based learning, since it supports meaning making of the experience (Mitchell et al., 2015). None of the programmes using in-service learning explicitly utilised reflection as a component of this approach and this may account for some of the reduced efficacy of this teaching method.

Knowledge and understanding. In 88 of the studies, outcomes related to improvements in knowledge were evaluated. Overwhelmingly (85%), studies reported staff increased their knowledge following training. In the 17% of studies adopting a qualitative methodology, ten studies used staff self-report post-training (via interviews, focus groups, or open-ended questionnaire) as the sole data collection method. All of these studies found staff

Page 20 of 55

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

felt attendance at training had increased their knowledge. Of the five studies also using observation of practice as a data collection method, two reported no change or mixed results for knowledge gains, suggesting this may offer a more robust method for assessing knowledge gains qualitatively than self-report alone. The remaining studies assessed knowledge gains using validated measures or un-validated questionnaires. Greater credibility has been given to studies using validated measures of knowledge within the synthesis, but all studies have contributed to the findings.

A training method that consistently led to no or weak knowledge gains was learning through a written resource (either hard copy or on-line), as assessed in three studies all using non-validated quantitative knowledge questionnaires, suggesting this should not be used as a primary mode of learning. This is a finding that is replicated in research in other health fields (Bluestone et al., 2013; Cervero & Gaines, 2014). A common feature of other studies reporting limited or no knowledge gains post-training, was utilisation of in-service learning as the main mode of delivery, with two of the three studies using this method reporting variable, limited or no learning had taken place. The main reasons cited as barriers to effective in-service learning were, time available for the learner and mentor while working in-service, poor mentor engagement and a task-focussed organisational culture (Skaalvik, Normann, & Henriksen, 2010; Skog, Negussie, & Grafstrom, 2000; Smythe et al., 2014). This indicates that it may be the poor setting conditions rather than in-service learning itself that is a barrier to learning via this method. However, of the studies reporting no knowledge gains, none had also evaluated learner reaction to training and, therefore, these issues were not explored systematically within the studies, to gain greater insight to them from a learner perspective.

Simulated, experiential and role-play based learning produced weak or variable outcomes for knowledge gains in two out of the five studies adopting these approaches. A

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

noticeable difference between the studies using these methods with positive knowledge gains and those with mixed or no gains, was the inclusion, in studies with positive results, of learner debriefing and feedback. This indicates it may be the way in which experiential/simulated learning was applied in these studies, rather than the method itself, which influenced learning efficacy. The existing body of research into simulated learning in non-dementia education within health and social care indicates simulation can be highly effective in knowledge and skills development (Stocker, Burmester, & Allen, 2014). However, a critical component of learning within simulation, role-play or experiential situations lies within the structured debriefing process (Dedy, Bonrath, Zevin, & Grantcharov, 2013; Gjeraa, Møller, & Østergaard, 2014; Gordon, Darbyshire, & Baker, 2012; Kottewar, Bearelly, Bearelly, Johnson, & Fleming, 2014; Stocker et al., 2014), which appears to be missing from training involving simulation within a number of the dementia educational studies.

Attitudes and beliefs. Thirty-three of the studies evaluated impact of training on staff attitudes or beliefs. Studies most commonly adopted a quantitative approach to measurement (81%), with the remaining studies adopting qualitative approaches including interviews, focus groups and analysis of learners' written assignments or reflective essays. Just over half (52%) of quantitative studies used a validated measure, with the remainder using a non-validated scale to assess attitude change. Of the 33 studies, 27% (n=9) reported no significant change in attitude post-training; one found improvements in attitudes for qualified nurses but not non-direct care staff (Long, 2013) and the remaining 23 studies (70%) reported positive attitude change. In four studies (Beville, 2002; Elliot & Adams, 2012; Mahendra, Fremont, & Dionne, 2013; van Zuilen et al., 2008) there were significant methodological weaknesses in the measures used to assess attitude change or in the data

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

analysis techniques adopted and, therefore, their results were not used to identify issues of training efficacy in this component of analysis.

A feature of six of the eight studies, in which attitude change was not observed, was training delivery in one or more individual sessions of two-hours or less duration, although multiple individual sessions were delivered to provide training programmes that ranged from 2-12 hours in total. In contrast, 18 of the 23 studies (78%) reporting positive attitude change included individual training sessions of longer than two-hours and were generally longer in overall programme duration (total programme lengths of four hours to ten days); indicating that staff need consolidated time engaging with sessions of training in order to effect attitude change. A similar finding has been reported in other studies within healthcare services (Bjerrum, Tewes, & Pederson, 2012).

Confidence, competence and self-efficacy. Thirty-seven studies reported on the impact of training on staff confidence, competence and self-efficacy in delivering dementia care. Over half (54%) adopted a quantitative methodology, 14% a qualitative design and 32% mixed-methods. Studies were largely moderate (n=23) and high quality (n=10), with five of the ten high quality studies utilising a validated measure of confidence, competence or self-efficacy. In the remaining studies confidence, competence and self-efficacy were measured using a non-validated measure, questionnaire or survey or through focus groups and interviews. Of the 37 studies, 81% reported improvements to staff confidence, competence, competence or self-efficacy, although significant positive change was found in only one-third of studies where this was measured using a validated tool (McCabe et al., 2014; Surr, Smith, Crossland, & Robins, 2016). Significant improvements were found in some sub-scales of competence, statistically but not clinically significant change was found in a further two (Elvish et al., 2014; Goyder, 2011) and no change in the remaining two (Finnema et al., 2005; Smythe et al., 2014). This suggests that non-validated measures may be unreliable in

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

assessing changes in these areas. This supports findings in broader educational research, which have consistently found self-report estimations of competency to be invalid (De Rijdt et al., 2013). However, despite these issues, the studies do provide information regarding whether health and social care staff feel they are more confident or competent in caring for people with dementia, whether or not this is evidenced through data collected using a validated measure. Staff reports of increased confidence, might in turn lead to additional positive outcomes, for example, improved role satisfaction or reduced stress and burden.

Twenty-six (70%) of the studies reporting improvements to staff confidence, competence or self-efficacy utilised small or large group face-to-face learning, suggesting active participation and discussion appears important for these aspects of learning. The value of interaction in learning was also evidenced in the studies using e-learning. Studies evaluating interactive web-based resources were largely found to increase confidence, competence or self-efficacy, although in one study (Irvine, Beaty, Seeley, & Bourgeois, 2013) this effect was seen only for non-clinical staff. However, in a separate study, a written, non-interactive, web-based resource did not increase GP confidence in administering driving assessments to people with dementia (Moorhouse & Hamilton, 2014). Therefore, despite caution needing to be applied to results due to issues about methodological robustness of studies, (inter)active learning appears more effective than passive approaches, in supporting improved confidence or self-efficacy, a finding that replicates that found in other workplace learning fields (Bluestone et al., 2013).

Studies also indicated that a combination of theory (through classroom-based learning) and practice (in-service learning) was more likely to produce positive results in improving staff confidence, competence or self-efficacy. This finding also reflects those found in other fields of professional education (De Rijdt et al., 2013; Tolsgaard et al., 2014).

Behaviour Change

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Sixty studies reported on changes to staff behaviour or practices as an outcome of training. The majority of studies were rated as moderate (47%) or high quality (43%) with only 10% of low quality. Data collection methods included review of practice documents such as care records (40%), observation of staff behaviours and practice (27%), and staff selfreport (40%), with some utilising more than one method. Self-report can provide useful information about perceived behaviour change, and may permit the exploration of the influence of potential confounding factors, thus providing contextual information about why behaviour change may or may not have occurred. However, it is also open to potential bias. Nearly two-fifths (37%) of studies relied solely on subjective staff self-report to assess behaviour change and thus the results must be treated with caution, although 27% of these studies were rated as high quality and a further 64% moderate quality. Areas of change reported in studies included communication, anti-psychotic prescribing and administration, person-centred care/general care practice improvements, restraint and implementation of a specific care process or tool. Studies using self-report methods only were more likely to assess broader, less specific care improvements (e.g. delivery of care that is more personcentred, general practice changes) than studies using more objective measures. They were also more likely to report wholly or some positive outcomes for behaviour change (86% and 14% respectively), compared to studies using at least one objective measure (68% wholly positive, 26% mixed results, 5% no change on any outcomes).

The most commonly adopted teaching method in over half of studies (58%) where positive outcomes for behaviour change were reported, was the inclusion of structured application of learning into practice. Approaches to achieving this included, in-practice activities or projects to be implemented as part of, between or after training sessions; expert clinical supervision; application of a participatory action research cycle; provision of tools or decision support software and development of staff as Champions who support

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

implementation of training in practice. Conversely, a common feature of over half (53%) of studies where mixed or no change outcomes were reported was use of a purely classroombased approach to learning. While the content of this was varied and included active and passive approaches, application of learning in practice as part of the training itself was not present. This indicates the importance for behaviour change, of supporting staff to implement learning in structured ways as part of training, ahead of expectations around application in everyday practice.

A further common feature of studies reporting positive behaviour change was the inclusion of structured assessment or care planning tools, or care delivery approaches within training. This suggests staff may more readily adapt behaviours and practices when provided with a specific tool or method to guide change in a structured way. Overall, the importance of structured approaches to changing practice behaviours is indicated in the literature. The use of a structured tool or set process within dementia care training provides learners with a goal or desired behaviour or way of practicing to be achieved. This finding is replicated in research conducted in other areas of education and training (Bluestone et al., 2013), which in particular identify goal setting as an effective method for driving behaviour change (Goodwin, Ostuzzi, Khan, Hotopf, & Moss-Morris, 2016; Porcheret et al., 2014).

Of the 15 studies that reported mixed or no-change results for behaviour change, only four also evaluated reaction and seven learning. Two of the studies that examined reaction reported mixed staff reactions, with lack of relevance/applicability of the training to practice being identified as a key issue by staff attending. This indicates staff may be less likely to make efforts to modify their behaviours or may be unable to do so, if the training does not provide them with relevant methods to support this. Of the studies also evaluating learning, six reported some improvements in knowledge and one improvement in staff confidence. One reported no significant change in staff attitudes. This suggests that while training may

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

increase staff knowledge this does not automatically translate into behaviour change in practice.

Outcomes, Results or Impact

Fifty of the studies reported on the impact of education and training on outcomes or results, for people with dementia (76%), staff (32%) and family caregivers/relatives (8%). Studies tended to be of medium (46%) to high (50%) quality with only 4% rated as low quality. Only 8% utilised a qualitative design, with 82% adopting a quantitative methodology and 10% using mixed methods.

Outcomes for people with dementia. Thirty-nine studies reported outcomes for people with dementia, however, in one study (Kellett, Moyle, McAllister, King, & Gallagher, 2010) data was collected solely via staff report using qualitative interviews, meaning it was excluded from the analysis due to unreliability of results due to bias. Most commonly, the outcomes evaluated in studies were impact on behaviours of people with dementia such as agitation, anxiety and aggression (66%), quality of life (21%), depression (14%), communication (11%) and activities of daily living (11%).

In 68% of studies a positive improvement in all or some outcomes was found and in all of these studies the training adopted face-to-face teaching methods, with active participation for example through discussing examples from their own practice and engagement in problem solving forming a common feature. This finding is aligned with existing understanding regarding practice learning approaches most likely to lead to practice change: where situated learning, whereby learners collaboratively discuss genuine problems arising from their professional practice, provides optimal learning conditions (Webster-Wright, 2009). In a number of studies, this was supported by clinical supervision or inservice learning with mentorship, which offered learners the opportunity to gain feedback on and discuss their own practice. A similar finding is reported in broader in-service learning

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

research, where clinically integrated teaching has been found to support improvements across a range of outcomes (Bluestone et al., 2013).

A second common feature was training facilitation by an experienced trainer who was usually also an experienced clinician. On the other hand, a feature of 30% of the training in studies where no change results were found was the use of a member of staff or in-house trainer to deliver some or all of the training intervention. Therefore, trainer qualities appear significant in influencing outcomes for people with dementia, beyond the level of participant reaction to training. While the current educational literature discusses areas such as instructor expertise and credibility with regard to student engagement, motivation (Imlawi, Gregg, & Karimi, 2015), satisfaction and learning or educational outcomes (Paechter, Maier, & Macher, 2010), there is limited discussion of if and how instructor expertise impacts on the behavioural and outcome levels of the Kirkpatrick framework.

Outcomes for family members. Only four studies evaluated outcomes for family members/carers, in the form of satisfaction with care of their relative. In two of these, the training included a component focussed on working positively with and engaging families. The two studies using qualitative methods reported greater satisfaction with care, but also identified areas where families felt further improvements could be made. The two studies adopting quantitative measures of satisfaction found no significant changes between pre and post-training. Given the low numbers of studies in this area, it is not feasible to draw any further conclusions regarding training characteristics most likely to lead to improved family member outcomes.

Outcomes for staff. Sixteen studies reported on outcomes related to staff, falling into three categories: job satisfaction and accomplishment (44%); stress, strain and burden (56%); and exhaustion, burnout and health complaints (38%) with some studies evaluating outcomes across multiple categories. Studies showed training was more likely to lead to no change

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

than to positive outcomes across all categories. Studies that reported some positive outcomes for staff utilised training that was more likely to be longer in total duration (8+ hours in total) and comprised multiple individual sessions of 90-mins to a full day duration. Studies reporting no change results utilised training which was more likely to be shorter overall (less than 7.5 hours) and have shorter individual session length (1-hour or less). This suggests that increasing staff job satisfaction and reducing stress or exhaustion is more likely to occur if training permits greater depth of staff engagement in terms of the overall training programme, and individual session length. This finding mirrors that of the broader professional development literature which advocates an educational model of ongoing, situated learning within a workplace context, as opposed to a 'training' model of discrete sessions, decontextualized from real-world practice (Webster-Wright, 2009). Given the earlier finding that staff attitudes are more likely to be improved in studies with longer individual session duration, this suggests there may be some interaction between staff attitudes towards people with dementia and their feelings of role satisfaction, stress or exhaustion, a finding supported by research in this area (Brodaty, Draper, & Low, 2003; Willemse et al., 2014). This suggests some inter-relationship is likely to be present between the Kirkpatrick levels, for staff related outcomes.

A further feature of over half of the studies reporting positive staff outcomes was inclusion in the training of a structured tool, manual or practice guidelines. This suggests that staff may feel more satisfied and less stressed, if they are provided with materials that help to give a clear structure to follow in their practice.

Studies evaluating impact across all levels

Only three studies evaluated all levels of the framework (Clare et al., 2013; McCurry, LaFazia, Pike, Logsdon, & Teri, 2012; Speziale, Black, Coatsworth-Puspoky, Ross, & O'Regan, 2009). Two of these studies were high quality and one moderate and all used a

 RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

quantitative design. The training programmes they utilised were similar in delivery methods (classroom and classroom + practice-based work), were of at least 3 hours total duration and had individual session length of 1-2 hours (where reported). All reported positive learner reactions to the training, improvements in knowledge post-training, improvements in behaviour post-training and at least some improved outcomes for people with dementia. However, none of the studies discussed the inter-relationship between factors at each Kirkpatrick level in contributing to efficacy of the training programme and their quantitative design limits their ability to offer learner or service-based explanations for potential efficacy.

Discussion

The review revealed a wide range of published international research, evaluating efficacy of dementia training for staff working across a wide range of health and social care settings. As was found in previous systematic literature reviews in this area (Alushi et al., 2015; Beeber et al., 2010; Brody & Galvin, 2013; Eggenberger et al., 2013; Elliott et al., 2012; Fossey et al., 2014; Kuske et al., 2007; McCabe et al., 2007; Perry et al., 2011; Raymond et al., 2014; Spector et al., 2013; Zientz et al., 2007), both the quality of evidence and reported efficacy of training interventions varied. This review identified that research has predominantly been conducted in care home settings, largely with qualified nurses or nurse aides/care assistants. There were smaller numbers of studies in other health and social care settings and with other staff groups. Existing research has evaluated training delivered using a variety of teaching methods, although predominantly face-to-face learning was adopted often alongside other methods. However, a limitation of many of the studies was use of a pre-post design, or a 'no training' control group in randomised studies. Few studies compared the efficacy of different training methods against each other. Therefore, where positive results were found this only provided an indicator of efficacy compared to receiving no training and thus provided limited understanding of whether the methods used were

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

optimal for delivering particular outcomes. There was a general lack of attempt to address potential methodological bias in the majority of studies, with many using qualitative selfreport and non-validated measures or questionnaires to assess change on outcomes, sometimes only collected immediately post-training, meaning comparison with pre-training or longer term practice were not possible. On the other hand, studies using validated measures often did not collect additional data; therefore, failing to examine what learners felt were particularly useful or unhelpful components of training that could help to explain quantitative results. A further limitation of the majority of studies was their focus on only one or two Kirkpatrick levels of efficacy, meaning they were unable to examine potential inter-relationships between factors. Only three studies evaluated all levels of the framework (Clare et al., 2013; McCurry et al., 2012; Speziale et al., 2009), but none have explored the inter-relationship between factors. We therefore, have provided some discussion of the potential inter-relationships across Kirkpatrick levels, based on our synthesis of the evidence and inferences we can draw from studies evaluating two or more levels. However, our conclusions remain tentative given the lack of exploration of these relationships within published studies. Therefore, as with previous systematic reviews, this study indicates the need to conduct further robust research that examines the efficacy of dementia training, taking into account the multi-faceted range of outcomes and inter-relationships between factors that are likely to influence these. The use of mixed methods, which can provide robust evidence of efficacy, but also the possibility to explain findings through gaining learner and facilitator perspectives, would appear advantageous. Future research should also engage more clearly with existing educational research in workplace learning across other professions. While there are some factors which may be particular to a specific professional workforce, such as that providing dementia care, our review has identified similarities of

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

experience across educational research of different workforce groups and thus suggests benefits to dementia educators of engaging with this body of literature.

Despite the limitations of existing research, this review has identified a number of key features that seem to exist in effective dementia training and which support understanding of approaches to effective professional development and workplace education more broadly. Training/education most likely to be effective:

- Is relevant and realistic to the role, experience and practice of learners rather than a one-size-fits-all training programme;
- Includes active participation;
- Underpins practice-based learning with theoretical or knowledge-based content;
- Ensures experiential and simulation-based learning includes adequate time for debriefing and discussion;
- Is delivered by an experienced trainer/facilitator who is able to adapt it to the needs of each group;
- Does not involve reading written materials (paper or web-based) or in-service learning as the sole teaching method;
- Is of a total duration of 8+ hours with individual training sessions of at least 90 minutes;
- Includes active, small or large group face-to-face learning either alone or in addition to another learning approach;
- Includes learning activities that support the application of training into practice;
- Provides staff with a structured tool, method or practice guideline to underpin care practice.

These features will be useful to consider in designing future dementia training and educational programmes. Given the similarities within our findings, to those reported in

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

other areas of adult and workplace education, they may also provide the basis for more general principles to inform the design and delivery of adult and workplace education and professional development, across health and social care and in other professional sectors such as teaching and business.

Limitations

Limitations to this review are, the inclusion only of papers published in English since 2000, may have excluded non-English language and older studies that might further contribute to understanding of effective dementia training and education. While using CIS and open inclusion criteria has permitted a broad examination of the evidence base, caution is however needed in interpreting the results. The limited number of studies in settings other than care homes and hospitals means application of findings to other settings providing dementia care, such as community and primary care requires caution.

Conclusion

Despite methodological weaknesses and variability in methods adopted there are some common features of training/education programmes that appear more efficacious and these may be adopted as underpinning guidelines for the design of new dementia training and educational programmes. They may also have relevance for adult professional development and workplace learning across a broad range of workplace settings. However, further robust research on the ingredients that lead to efficacy of dementia education and training is urgently required, to avoid continued utilisation of programmes that may have limited positive benefits.

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

References

- All-Party Parliamentary Group on Dementia. (2009). *Prepared to Care. Challenging the dementia skills gap.* London: The Stationary Office.
- Allen, J. M., & Wright, S. E. (2014). Integrating theory and practice in the pre-service teacher education practicum. *Teachers and Teaching*, 20(2), 136-151. doi:10.1080/13540602.2013.848568
- Alnes, R. E., Kirkevold, M., & Skovdahl, K. (2013). The influence of the learning climate on learning outcomes from Marte Meo counselling in dementia care. *Journal of Nursing Management, 21*(1), 130-140. doi:10.1111/j.1365-2834.2012.01436.x
- Alushi, L., Hammond, J. A., & Wood, J. H. (2015). Evaluation of dementia education programs for pre-registration healthcare students—A review of the literature. *Nurse Education Today*.
- Alzheimer's Association. (2015). Alzheimer's Disease Facts and Figures. *Alzheimer's and Dementia, 11*(3), 332+.
- Alzheimer's Disease International. (2010). World Alzheimer report: the global economic impact of dementia. London: Alzheimer's Disease International.
- Alzheimer's Society. (2009). Counting the cost. Caring for people with dementia on hospital wards. London: Alzheimer's Society.
- Arghode, V., & Wang, J. (2016). Exploring trainers' engaging instructional practices: a collective case study. *European Journal of Training and Development*, 40(2), 111-127. doi:10.1108/EJTD-04-2015-0033
- Bates, R. (2004). A critical analysis of evaluation practice: the Kirkpatrick model and the principle of beneficence. *Evaluation and Program Planning*, *27*, 341-347.
 doi:10.1016/j.evalprogplan.2004.04.011

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Beeber, A. S., Zimmerman, S., Fletcher, S., Mitchell, C. M., & Gould, E. (2010). Challenges and strategies for implementing and evaluating dementia care staff training in longterm care settings. *Alzheimer's Care Today*, 11(1), 17-39.

Benner, P. (1982). From novice to expert. American Journal of Nursing, 82(3), 402-407.

- Bettio, F., Simonazzi, A., & Villa, P. (2006). Change in Care Regimes and Female Migration: the Care Drain in the Mediterranean. *Journal of European Social Policy*, *16*(3), 271-285. doi:10.1177/0958928706065598
- Beville, P. K. (2002). Virtual Dementia Tour© helps sensitize health care providers. American Journal of Alzheimer's Disease & Other Dementias, 17(3), 183-190.
- Bjerrum, M., Tewes, M., & Pederson, P. (2012). Nurses' self-reported knowledge about and attitude to nutrition – before and after a training programme. *Scandinavian Journal of Caring Sciences*, 26(1), 81-89. doi:10.1111/j.1471-6712.2011.00906.x
- Bluestone, J., Johnson, P., Fullerton, J., Carr, C., Alderman, J., & BonTemp, J. (2013).
 Effective in-service training design and delivery: evidence from an integrative literature review. *Human Resources For Health*, *11*, 51. doi:10.1186/1478-4491-11-51
- Bourgeault, I., Atanackovic, J., Rashid, A., & Parpia, R. (2010). Relations Between
 Immigrant Care Workers and Older Persons in Home and Long-term Care. *Canadian Journal on Aging*, 29(1), 109-118. doi:10.1017/S0714980809990407.
- Brodaty, H., Draper, B., & Low, L. F. (2003). Nursing home staff attitudes towards residents with dementia: strain and satisfaction with work. *Journal of Advanced Nursing*, 44(6), 583-590.
- Brody, A. A., & Galvin, J. E. (2013). A review of interprofessional dissemination and education interventions for recognizing and managing dementia. *Gerontology & Geriatrics Education*, 34(3), 225-256. doi:10.1080/02701960.2013.801342

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Caldwell, K., Henshaw, L., & Taylor, G. (2005). Developing a framework for critiquing health research. *Journal of health, social and environmental issues, 6*(1), 45-54.
- Cangiano, A., & Shutes, I. (2010). Ageing, Demand for Care and the Role of Migrant Care Workers in the UK. *Journal of Population Ageing*, *3*, 1-2.
- Cervero, R. M., & Gaines, J. K. (2014). Effectiveness of continuing medical education: updated synthesis of systematic reviews (A. C. f. C. M. Education Ed.). Chicago, IL: Accreditation Council for Continuing Medical Education.
- Clare, L., Whitaker, R., Woods, R. T., Quinn, C., Jelley, H., Hoare, Z., ... Wilson, B. A. (2013). AwareCare: a pilot randomized controlled trial of an awareness-based staff training intervention to improve quality of life for residents with severe dementia in long-term care settings. *International Psychogeriatrics*, 25(1), 128-139. doi:10.1017/S1041610212001226
- Corlett, J. (2000). The perceptions of nurse teachers, student nurses and preceptors of the theory-practice gap in nurse education. *Nurse Education Today, 20*(6), 499-505. doi:10.1054/nedt.1999.0414
- Corwin, M., Owen, D., & Perry, C. (2008). Student service learning and dementia: bridging classroom and clinical experiences. *Journal of Allied Health*, *37*(4), e278-e295.
- Creaser, M., Hargreaves, J., & Ong, A. (2014). *Readiness to meet demand for skills: a study of five growth industries*. Adelaide, Southern Australia: National Centre for Vocational Education Research.
- Critical Appraisal Skills Programme (CASP). (2014). CASP Checklists. Retrieved from http://www.casp-uk.net/#!checklists/cb36
- Dall'Alba, G., & Sandberg, J. (2006). Unveiling Professional Development: A Critical Review of Stage Models. *Review of Educational Research*, 76(3), 383-412. doi:10.3102/00346543076003383

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Darling-Hammond, L., Barron, B., Pearson, P. D., Schoenfeld, A. H., Stage, E. K., Zimmerman, T. D., . . . Chen, M. (2008). *Powerful learning. What we know about teaching for understanding*. San Francisco, CA: Jossey Bass.
- De Rijdt, C., Stes, A., van der Vleuten, C., & Dochy, F. (2013). Influencing variables and moderators of transfer of learning to the workplace within the area of staff development in higher education: Research review. *Educational Research Review*, 8, 48-74. doi:10.1016/j.edurev.2012.05.007
- Dedy, N. J., Bonrath, E. M., Zevin, B., & Grantcharov, T. P. (2013). Teaching nontechnical skills in surgical residency: a systematic review of current approaches and outcomes. *Surgery*, 154(5), 1000-1008. doi:10.1016/j.surg.2013.04.034
- Department of Health. (2009). *Living well with dementia: A National Dementia Strategy*. London: Department of Health.
- Department of Health. (2012). Prime Minister's challenge on dementia. delivering major improvements in dementia care and research by 2015. London: Department of Health.
- Department of Health. (2013). Delivering high quality, effective, compassionate care: Developing the right people with the right skills and the right values. A mandate from the Government to Health Education England: April 2013 to March 2015. London: Department of Health.
- Department of Health. (2014). Delivering high quality, effective, compassionate care: Developing the right people with the right skills and the right values. A mandate from the Government to Health Education England: April 2014 to March 2015. London: Department of Health.
- Department of Health. (2016). Delivering high quality, effective, compassionate care: Developing the right people with the right skills and the right values. A mandate from

 RER

the Government to Health Education England: April 2016 to March 2017 London: Department of Health.

- Dixon-Woods, M., Cavers, D., Agearwal, S., Annandale, E., Arthur, A., Harvey, J., . . .
 Sutton, A. J. (2006). Conducting a critical interpretive synthesis of the literature on access to healthcare by vulnerable groups. *BMC Medical Research Methodology, 6*, 35. doi:10.1186/1471-2288-6-35
- Dixon-Woods, M., Cavers, D., Agerwal, S., Annandale, E., Arthur, A., Harvey, J., . . . Sutton,
 A. J. (2006). Conducting a critical interpretive synthesis of the literature on access to
 healthcare by vulnerable groups. *BMC Medical Research methodology*, *6*, 35.
 doi:10.1186/1471-2288-6-35
- Dreyfus, S. E. (1982). Formal models vs. human situational understanding: Inherent limitations on the modelling of business expertise. *Office Technology and People, 1*, 133-165.
- Dychtwald, K., Erikson, T. J., & Morison, R. (2006). *Workforce crisis: How to beat the coming shortage of skills and talent*. Boston, MA.: Harvard Business School Press.
- Eggenberger, E., Heimerl, K., & Bennett, M. I. (2013). Communication skills training in dementia care: a systematic review of effectiveness, training content, and didactic methods in different care settings *International Psychogeriatrics*, 25(3), 345-358. doi:10.1017/S1041610212001664
- Elliot, R., & Adams, J. (2012). Using a practice development project to improve standards of care for people with dementia. *Nursing Older People, 24*(8), 28.
- Elliott, K.-E. J., Scott, J. L., Stirling, C., Martin, A. J., & Robinson, A. (2012). Building capacity and resilience in the dementia care workforce: A systematic review of interventions targeting worker and organizational outcomes. *International Psychogeriatrics, 24*(6), 882-894. doi:10.1017/S1041610211002651

Page 38 of 55

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Elvish, R., Burrow, S., Cawley, R., Harney, K., Graham, P., Pilling, M., . . . Keady, J. (2014).
'Getting to Know Me': the development and evaluation of a training programme for enhancing skills in the care of people with dementia in general hospital settings. *Aging & Mental Health*, 18(4), 481-488. doi:10.1080/13607863.2013.856860

RER

Finnema, E., Dröes, R. M., Ettema, T., Ooms, M., Adèr, H., Ribbe, M., & Tilburg, W. (2005). The effect of integrated emotion-oriented care versus usual care on elderly persons with dementia in the nursing home and on nursing assistants: a randomized clinical trial. *International Journal of Geriatric Psychiatry*, 20(4), 330-343. doi:10.1002/gps.1286

Fossey, J., Masson, S., Stafford, J., Lawrence, V., Corbett, A., & Ballard, C. (2014). The disconnect between evidence and practice: a systematic review of person-centred interventions and training manuals for care home staff working with people with dementia. *International Journal of Geriatric Psychiatry*, 29(8), 797-807. doi:10.1002/gps.4072

Freeman, S., Eddy, S. L., McDonough, M., Smith, M. K., Okoroafor, N., Jordt, H., & Wenderoth, M. P. (2014). Active learning increases student performance in science, engineering, and mathematics. *PNAS*, *111*(23), 8410-8415. doi:10.1073/pnas.1319030111

Gallagher, P. (2004). How the metaphor of a gap between theory and practice has influenced nursing education. *Nurse Education Today*, *24*, 263-268.doi:10.1016/j.nedt.2004.01.006

Gauld, D., & Miller, P. (2004). The qualifications and competencies held by effective workplace trainers. *Journal of European Industrial Training*, 28(1), 8-22.
doi:10.1108/03090590410513866

- Ghosh, P., Satawadi, R., Joshi, J. P., Ranjam, R., & Singh, P. (2012). Towards more effective training programmes: a study of trainer attributes. *Industrial and Commercial Training*, 44(4), 194-202. doi:10.1108/00197851211231469
- Giangreco, A., Carugati, A., & Sebastiano, A. (2008). Are we doing the right thing? Food for thought on training evaluation and its context. *Personnel Review*, 39(2), 162-177. doi:10.1108/00483481011017390
- Gjeraa, K., Møller, T. P., & Østergaard, D. (2014). Efficacy of simulation-based trauma team training of non-technical skills. A systematic review. Acta Anaesthesiologica Scandinavica, 58(7), 775-787. doi:10.1111/aas.12336
- Goodwin, L., Ostuzzi, G., Khan, N., Hotopf, M. H., & Moss-Morris, R. (2016). Can We
 Identify the Active Ingredients of Behaviour Change Interventions for Coronary Heart
 Disease Patients? A Systematic Review and Meta-Analysis. *PloS One, 11*(4),
 e0153271. doi:10.1371/journal.pone.0153271
- Gordon, M., Darbyshire, D., & Baker, P. (2012). Non-technical skills training to enhance patient safety: a systematic review. *Medical Education*, 46(11), 1042-1054. doi:10.1111/j.1365-2923.2012.04343.x
- Goyder, J. (2011). Staff Training using STAR (Staff Training in Assisted Living Residences):A Pilot Study in UK Residential Care Homes (Thesis). (D. Clin Psy), King's College London.
- Hatlevik, I. K. R. (2011). The theory-practice relationship: reflective skills and theoretical knowledge as key factors in bridging the gap between theory and practice in initial nursing education. *Journal of Advanced Nursing*, 68(4), 868-877. doi:10.1111/j.1365-2648.2011.05789.x
- Health and Social Care Board. (2016). *The dementia learnign and development framework*. Belfast: Health & Social Care Board.

Page 40 of 55

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Health Education England Skills for Health and Skills for Care. (2015). *Dementia Core Skills Education and Training Framework*. London: Health Education England.
- Holton, E. F. (1996). The flawed four-level evaluation model. *Human Resource Development Quarterly*, 7(1), 5-21.
- Imlawi, J., Gregg, D., & Karimi, J. (2015). Student engagement in course-based social networks: The impact of instructor credibility and use of communication. *Computers* & *Education*, 88, 84-96. doi:10.1016/j.compedu.2015.04.0150360-1315
- Irvine, A. B., Beaty, J. A., Seeley, J. R., & Bourgeois, M. (2013). Use of a Dementia Training Designed for Nurse Aides to Train Other Staff. *Journal of applied gerontology*, 32(8), 936-951. doi:10.1177/0733464812446021
- Jackson, A., Bluteau, P., & Furlong, J. (2013). Interprofessional Working in Practice Avoiding a Theory-Practice Gap. *International Journal of Practice Learning in Health and Social Care, 1*(1), 90-92. doi:10.11120/pblh.2013.00001
- Jensen, J. L., Kummer, T. A., & Godoy, P. d. M. (2014). Improvements from a flipped classroom may simply be fruits of active learning. *CBE - Life Sciences Education*, 14(1), ar5. doi:10.1187/cbe.14-08-0129
- Kangasniemi, M., Kallio, H., & Pietila, A.-M. (2014). Towards environmentally responsible nursing: a critical interpretive synthesis. *Journal of Advanced Nursing*, 70(7), 1465-1478.
- Kellett, U., Moyle, W., McAllister, M., King, C., & Gallagher, F. (2010). Life stories and biography: A means of connecting family and staff to people with dementia. *Journal* of Clinical Nursing, 19(11-12), 1707-1715. doi:10.1111/j.1365-2702.2009.03116.x
- Kessels, J. P. A. M., & Korthagen, F. A. J. (1996). The relationship between theory and practice: back to the classics. *Educational Researcher*, *25*(3), 17-22.

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Khamarko, K., Koester, K. A., Bie, J., Baron, R. B., & Myers, J. J. (2012). Developing Effective Clinical Trainers. *SAGE Open, 2*(2). doi:10.1177/2158244012448486
- Kirkpatrick, D. L. (1979). Techniques for evaluating training programmes. *Training and Development Journal*(June), 178-192.
- Kirkpatrick, D. L. (1984). Techniques for evaluating training programmes. *Training and Development Journal, 32*(9), 6-9.

Knapp, M., Prince, M., Albanese, E., Banerjee, S., Dhanasiri, S., Fernandez, J. L., ... Stewart, R. (2007). *Dementia UK: Report to the Alzheimer's Society*. London: Alzheimer's Society.

- Knowles, M. S. (1970). *The modern practice of adult education: andragogy versus pedagogy*. New York: Association Press.
- Knowles, M. S., Holton, E. F., & Swanson, R. A. (2015). *The Adult Learner. The definitive classic in adult education and human resource development 8th edition*. London: Routledge.
- Kottewar, S. A., Bearelly, D., Bearelly, S., Johnson, E. D., & Fleming, D. A. (2014).
 Residents' end-of-life training experience: a literature review of interventions. *Journal of Palliative Medicine*, *17*(6), 725-732. doi:10.1089/jpm.2013.0353
- Kuske, B., Hanns, S., Luck, T., Angermeyer, M. C., Behrens, J., & Riedel-Heller, S. G. (2007). Nursing home staff training in dementia care: a systematic review of evaluated programs. *International Psychogeriatrics*, 19(5), 818-841. doi:10.1017/S1041610206004352
- Kyndt, E., & Baert, H. (2013). Antecedents of Employees' Involvement in Work-Related Learning: A Systematic Review. *Review of Educational Research*, *83*(2), 273-313. doi:10.3102/0034654313478021

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Landers, M. G. (2000). The theory-practice gap in nursing: the role of the nurse teacher. Journal of Advanced Nursing, 32(6), 1550-1556. doi:10.1046/j.1365-2648.2000.01605.x
- Long, C. O. (2013). Pain management education in long-term care: it can make a difference. *Pain Management Nursing*, *14*(4), 220-227. doi:10.1016/j.pmn.2011.04.005

Mahendra, N., Fremont, K., & Dionne, E. (2013). Teaching future providers about dementia: The impact of service learning. *Seminars and Speech and Language*, 34(1), 5. doi:10.1055/s-0033-1337390

Mansouri, M., & Lockyer, J. (2007). A meta-analysis of continuing medical education effectiveness. *Journal of continuing education in the health professions, 27*(1), 6-15.

McCabe, M. P., Bird, M., Davison, T. E., Mellor, D., MacPherson, S., Hallford, D., & Seedy, M. (2014). An RCT to evaluate the utility of a clinical protocol for staff in the management of behavioral and psychological symptoms of dementia in residential aged-care settings. *Aging & Mental Health*, *19*(9), 1-9.

doi:10.1080/13607863.2014.967659

- McCabe, M. P., Davison, T. E., & George, K. (2007). Effectiveness of staff training programs for behavioral problems among older people with dementia. *Aging and Mental Health*, 11(5), 505-519. doi:10.1080/13607860601086405
- McCurry, S. M., LaFazia, D. M., Pike, K. C., Logsdon, R. G., & Teri, L. (2012).
 Development and evaluation of a sleep education program for older adults with dementia living in adult family homes. *American Journal of Geriatric Psychiatry*, 20(6), 494-504. doi:10.1097/JGP.0b013e318248ae79
- Mitchell, T. D., Richard, F. D., Battistoni, R. M., Rost-Banik, C., Netz, R., & Zakoske, C. (2015). Reflective Practice that Persists: Connections Between Reflection in Service-

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Learning Programs and in Current Life. *Michigan Journal of Community Service Learning*(Spring), 49-63.

Moorhouse, P., & Hamilton, L. M. (2014). Not If, But When: Impact of a Driving and Dementia Awareness and Education Campaign for Primary Care Physicians. *Canadian Geriatrics Journal*, 17(2), 70-75. doi:10.5770/cgj.17.109

- Opfer, V. D., & Pedder, D. (2011). Conceptualizing teacher professional education. *Review of Educational Research*, 81(3), 376-407. doi:10.3102/0034654311413609
- Paechter, M., Maier, B., & Macher, D. (2010). Students' expectations of, and experiences in e-learning: Their relation to learning achievements and course satisfaction. *Computers & Education*, 54, 222-229. doi:10.1016/j.compedu.2009.08.005
- Pereles, L., Davis, D., Lockyer, J., Robinson, B. E., Ryan, D., & Spivak, B. (2003). The use of the opinion leader in continuing medical education. *Medical Teacher*, 25(4), 438-441. doi:10.1080/0142159031000137472

Perry, M., Drašković, I., Lucassen, P., Vernooij-Dassen, M., van Achterberg, T., & Rikkert, M. O. (2011). Effects of educational interventions on primary dementia care: A systematic review. *International Journal of Geriatric Psychiatry*, 26(1), 1-11. doi:10.1002/gps.2479

Porcheret, M., Main, C., Croft, P., McKinley, R., Hassell, A., & Dziedzic, K. (2014).
Development of a behaviour change intervention: a case study on the practical application of theory. *Implementation Science*, *9*, 42. doi:10.1186/1748-5908-9-42

Pulsford, D., Jackson, G., O'Brien, T., Yates, S., & Duxbury, J. (2013). Classroom-based and distance learning education and training courses in end-of-life care for health and social care staff: a systematic review. *Palliative Medicine*, *27*(3), 221-235. doi:10.1177/0269216311429496

Page 44 of 55

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Raymond, M., Warner, A., Davies, N., Baishnab, E., Manthorpe, J., & Iliffe, S. (2014).
 Evaluating educational initiatives to improve palliative care for people with dementia: a narrative review. *Dementia (London, England), 13*(3), 366-381.
 doi:10.1177/1471301212474140
- Resnick, B., Cayo, J., Galik, E., & Pretzer-Aboff, I. (2009). Implementation of the 6-week educational component in the Res-Care intervention: process and outcomes. *Journal of continuing education in nursing*, *40*(8), 353-360. doi:10.3928/00220124-20090723-04
- Roth, W., M., Mavin, T. J., & Dekker, S. W. A. (2014). The theory-practice gap:
 epistemology, identity and education. *Education and Training*, 56(6), 521-536.
 doi:10.1108/ET-11-2012-0117
- Royal College of Psychiatrists. (2011). *Report of the National Audit of Dementia Care in General Hospitals*. London: Healthcare Quality Improvement Partnership.
- Scully, N. J. (2011). The theory-practice gap and skill acquisition: An issue for nursing education. *Collegian*, 18(2), 93-98. doi:10.1016/j.colegn.2010.04.002
- Severinsson, E. I. (1998). Bridging the gap between theory and pratice: a supervision programme for nursing students. *Journal of Advanced Nursing*, *27*, 1269-1277.
- Skaalvik, M. W., Normann, H. K., & Henriksen, N. (2010). Student experiences in learning person-centred care of patients with Alzheimer's disease as perceived by nursing students and supervising nurses. *Journal of Clinical Nursing*, *19*(17/18), 2639-2648. doi:10.1111/j.1365-2702.2010.03190.x
- Skog, M., Negussie, B., & Grafstrom, M. (2000). Learning dementia care in three contexts: practical training in day-care, group dwelling and nursing home. *Journal of Advanced Nursing*, 32(1), 148-157.

RER

- Smith, P. J. (2003). Workplace learning and flexible delivery. *Review of Educational Research*, *73*(1), 53-88.
- Smythe, A., Jenkins, C., Harries, M., Atkins, S., Miller, J., Wright, J., . . . Oyebode, J. (2014). Evaluation of dementia training for staff in acute hospital settings. *Nursing Older People*, 26(2), 18-24. doi:10.7748/nop2014.02.26.2.18.e527
- Sogunro, O. A. (2015). Motivating factors for adult learners in higher education. International Journal of Higher Education, 4(1), 22-37. doi:10.5430/ijhe.v4n1p22
- Spector, A., Orrell, M., & Goyder, J. (2013). A systematic review of staff training interventions to reduce the behavioural and psychological symptoms of dementia. *Ageing Research Reviews*, 12(1), 354-364. doi:10.1016/j.arr.2012.06.005
- Speziale, J., Black, E., Coatsworth-Puspoky, R., Ross, T., & O'Regan, T. (2009). Moving Forward: Evaluating a Curriculum for Managing Responsive Behaviors in a Geriatric Psychiatry Inpatient Population. *Gerontologist*, 49(4), 570-576. doi:10.1093/geront/gnp069
- Stocker, M., Burmester, M., & Allen, M. (2014). Optimisation of simulated team training through the application of learning theories: a debate for a conceptual framework.
 BMC Medical Education, 14, 69. doi:10.1186/1472-6920-14-69
- Surr, C. A., Smith, S. J., Crossland, J., & Robins, J. (2016). Impact of a person-centred dementia care training programme on hospital staff attitudes, role efficacy and perceptions of caring for people with dementia: A repeated measures study. *International Journal of Nursing Studies, 53*, 144-151. doi:10.1016/j.ijnurstu.2015.09.009
- Tamkin, P., Yarnall, J., & Kerrin, M. (2002). Kirkpatrick and beyond: A review of models of training evaluation. Brighton: The Institute for Employment Studies.

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

- Taylor, D. C. M., & Hamdy, H. (2013). Adult learning theories: Implications for learning and teaching in medical education: AMEE guide No. 83. *Medical Teacher*, 35, e1561e1572.
- Tolsgaard, M. G., Rasmussen, M. B., Tappert, C., Sundler, M., Sorensen, J. L., Ottesen, B., . .
 Tabor, A. (2014). Which factors are associated with trainees' confidence in performing obstetric and gynecological ultrasound examinations? *Ultrasound in Obstetrics and Gynecology*, *43*(4), 444-451. doi:10.1002/uog.13211
- Traynor, V., Cumming, A., & Britten, N. (2015). Developing an inter-disciplinary crosssetting dementia care competency framework. New South Wales: University of Wollongong, NSW.
- Tullo, E., & Allan, L. (2011). What should we be teaching medical students about dementia? *International psychogeriatrics / IPA*, 23(7), 1044-1050. doi:10.1017/S1041610211000536
- Upton, D. J. (1999). How can we achieve evidence-based practice if we have a theorypractice gap in nursing today? *Journal of Advanced Nursing*, *29*(3), 549-555. doi:10.1046/j.1365-2648.1999.00922.x
- US Department of Health and Human Services. (2013). National Plan to Address Alzheimer's Disease: 2013 update. Available on-line from

http://aspe.hhs.gov/daltcp/napa/NatlPlan2013.shtml.

van Zuilen, M. H., Mintzer, M. J., Milanez, M. N., Kaiser, R. M., Rodriguez, O., Paniagua, M. A., . . . Roos, B. A. (2008). A competency-based medical student curriculum targeting key geriatric syndromes. *Gerontology & Geriatrics Education, 28*(3), 29-45. doi:10.1300/J021v28n03 03

Vollmar, H. C., Mayer, H., Ostermann, T., Butzlaff, M. E., Sandars, J. E., Wilm, S., & Rieger, M. A. (2010). Knowledge transfer for the management of dementia: a cluster-

RER

55	NEN	
	EFFECTIVE DEMENTIA EDUCATION AND TRAINING	47
	randomised trial of blended learning in general practice. Implementation Science,	
	5(1), 1. doi:10.1186/1748-5908-5-1	
	Wales, C. C. f. (2016). Good Work: A dementia learning and development framework for	
	Wales. Cardiff: Care Council for Wales.	
	Wall, P., Andrus, P. M., & Morrison, P. (2014). Bridging the Theory Practice Gap through	L
	Clinical Simulations in a Nursing UnderGraduate Degree Program in Australia.	
	International Journal of Learning, Teaching and Educational Research, 8(1), 127-	
	135.	
	Webster-Wright, A. (2009). Reframing Professional Development Through Understanding	5
	Authentic Professional Learning. Review of Educational Research, 79(2), 702-739.	
	WHO/Alzheimer's Disease International. (2012). Dementia: A Public Health Priority.	
	London: WHO.	
	Wilkinson, D. M., Smallidge, D., Boyd, L. D., & Giblin, L. (2015). Students' Perceptions of	of
	Teaching Methods That Bridge Theory to Practice in Dental Hygiene Education.	
	Journal of Dental Hygiene, 89(5), 330-337.	
	Willemse, B. M., De Jonge, J., Smit, D., Visser, Q., Depla, M. F. I. A., & Pot, A. M. (2014)	I).
	Staff's person-centredness in dementia care in relation to job characteristics and job	0-
	related well-being: a cross-sectional survey in nursing homes. Journal of Advanced	!
	Nursing, 71(2), 404-416. doi:10.1111/jan.12505	
	Wilson, J. H. (2014). Investing in English Skills: The Limited English Proficient Workford	e in
	U.S. Metropolitan Areas. Washington, D.C.: The Metropolitan Policy Programme	at
	Brookings.	
	Zaher, E., & Ratnapalan, S. (2012). Practice-based small group learning programs:	
	Systematic review. Canadian Family Physician, 58(6), 637-642.	
	Systematic review. Canadian Family Physician, 58(6), 637-642.	

http://mc.manuscriptcentral.com/rer

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Zientz, J., Rackley, A., Chapman, S. B., Hopper, T., Mahendra, N., Kim, E. S., & Cleary, S. (2007). Evidence-based practice recommendations for dementia: educating caregivers on Alzheimer's disease and training communication strategies. *Journal of Medical Speech-Language Pathology*, 15(1), liii-lxiv.

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Table 1

Characteristics of Included Studies

8	Description	N (%)
9 —	Country of study	
10	US	58 (38)
10	UK UK	30 (20)
12	Australia	17(11)
14	Canada	18 (12)
15	Netherlands	5(2)
16	Norway	5(3) 5(3)
17	Norway Swadar	(3)
18	Sweden	(4)
19	France	3(2)
20	Rest of Europe	8 (5)
21	Other	4 (3)
22	Setting	
23	Care Homes	75 (49)
24	Hospitals	14 (9)
25	Higher Education	18 (12)
26	Primary Care	12 (8)
27	Community	7 (5)
28	Assisted living	5 (3)
29	Day care	2(1)
31	More than one service setting	19 (13)
32	Staff group training aimed at	- (-)
33	Nurse Aides/Assistants	56 (37)
34	Unknown staff groups within a particular service setting	53 (35)
35	Oualified Nurses	51 (34)
36	General Practitioners/Doctors/Physicians	23(0)
37	University students on health/social care programmes	23(0)
38	Managers	15(10)
39		13 (10)
40	Activities staff	13 (9)
41	Social workers	13 (9)
42	Ancillary staff (porters, laundry etc)	11(/)
43	Allied Health Professionals	9 (6)
44	Pharmacists	3 (2)
45	Unknown	2 (1)
40	Methodology	
47	Quantitative	97 (64)
40 /Q	Qualitative	21 (14)
49 50	Mixed methods	34 (22)
51	Quality	
52	High	52 (34)
53	Medium	79 (52)
54	Low	21(14)
55		(- ·)
56		
57		
58		

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING

Table 2

Teaching and Learning Approaches Adopted

Teaching and learning approach	Number of studies n (%)*
Small/large group face-to-face	69 (45)
Small/large group face-to-face plus	51 (34)
In-service/practice-based learning	25 (16)
Mentorship/supervision	6 (4)
Experiential learning/simulation	7 (5)
Practical exercise/project	5 (3)
Written materials	3 (2)
Psychoeducation	1 (1)
On-line learning	1 (1)
DVD	1 (1)
Drama	1 (1)
Peer support	1 (1)
Individual/Group DVD	9 (6)
Written resource	6 (4)
On-line	5 (3)
In-service	4(3)
Train-the-trainer	3 (2)
Peer support/learning set	2(1)
Simulation/experiential	2(1)
Drama	2(1)
Counselling	2(1)
Psychoeducation	1(1)
*Number of studies is greater than 152 since s	some studies compared more than one type of
training	

RER

2 3 4 5	EFFECTIVE DEMENTIA I Table 3	EDUCATION AND TRAINING
6 7 8 9	Adapted quality rating crite	ria
10 11 12 13	Quality criteria Are	Specific
14 15 16 17	1. The research aims and questions/hypotheses clearly stated?	• Does the author(s) clearly sta
18 19 20 21	2. Ethical issues addressed?	 Does the author(s) state that Does the author(s) demonstrative (E.g. informed consent, configurate)
22 23 24 25 26 27 28 29 30 31 32 33 34	3. The methodology/study design appropriate to the research question and rationale for choice evident?	 etc.). Does the author explicitling Is the chosen methodolo quantitative or mixed methodolo quantitative or mixed methodolo quantitative research are search aims and questing For qualitative research provide the author (s) justifies a study?
35 36 37 38 39 40 41 42 43 44	4. The sample size, selection and description appropriate?	 Does the authors(s) clean Does the sample size app Does the author(s) adequ to care receiver etc.) so t Does the author(s) descr
45 46 47		http://n

Specific questions to consider when rating Rating uthor(s) clearly state what they plan to research? 0 = No1 = Partially2 = Yesuthor(s) state that ethical approval was sought? 0 = No1 = Partiallyuthor(s) demonstrate an awareness of the ethical issues raised by the study? 2 = Yesned consent, confidentiality, responding to upset or distress, withdrawal he author explicitly state what research methodology they have chosen? 0 = No1 = Partially butchosen methodology appropriate to the research question? E.g. qualitative or with ative or mixed methods approach? Where qualitative – grounded theory, weaknesses/missing thnography etc info alitative research - does the chosen methodology appear appropriate to the 2 = Yesch aims and questions? Is this fully justified? antitative research - Does the author(s) clearly state the design of the study? he author(s) justify the research design used? (E.g. longitudinal, cross al etc.) Does the author(s) identify the main variables investigated in the he authors(s) clearly state how the study sample size was identified? 0 = No1 = Partially buthe sample size appear to be large enough/appropriate? with he author(s) adequately describe the sample (E.g. Gender, age, relationship weaknesses/missing receiver etc.) so that the reader can determine transferability of findings? info he author(s) describe the context of where the samples were recruited from?

Page 52 of 55

FFECTIVE DEMENTIA ED	UCATION AND TRAINING	
5. Are the method(s) of data collection appropriate, reliable and valid?	 Does the author(s) describe the method of recruitment used? (E.g. the sampling method, recruitment etc.) Does the author(s) identify the inclusion criteria? For quantitative studies: Does the author(s) justify that the measure is suitable for this population? Does the author(s) use measures that measure the desired constructs? Does the author(s) indicate whether the measures used have good psychometric properties? (E.g. test-retest reliability, inter-rater-reliability, internal reliability and internal consistency (Cronbach's alpha)). Does the author(s) indicate that the measures used have demonstrated validity? For qualitative studies Were the methods used appropriate for the participants, valid and likely to be free of bias? Does the author justify why particular data collection approaches were 	2 = Yes 0 = No 1 = Partially but with weaknesses/missing info 2 = Yes
 The method(s) of data analysis reliable and valid? 	 used e.g. interviews, focus groups For quantitative studies Does the author(s) state which statistic tests were used? Does the author(s) use statistical tests that appear to be appropriate to the nature of the data collected? (E.g. Does the data meet the assumptions of the test). Were the statistical tests used appropriate to the research question? Does the author(s) consider the impact of extraneous variables and control for these within the analysis process? Does the author(s) provide evidence of statistical findings? (E.g. Data within the text, tables etc.). Does the author(s) state the levels of significance? For qualitative studies Does the authors(s) state what approach they used to data analysis? 	0 = No 1 = Partially 2 = Yes
	http://mc.manuscriptcentral.com/rer	

RER

 RER

7. The findings and discussion clearly stated and appropriate?	 Does this approach appear to be suitable to the data gathered? Does the approach appear to have been implemented in a structured/robust manner? Does the author(s) provide details of how findings were validated? Does the author(s) explicitly state their findings? Does the author(s) present the statistical/qualitative data in a clear manner? For quantitative studies Does the author(s) clearly differentiate between significant and nonsignificant findings? For qualitative studies Does the author(s) clearly identify key themes or issues arising from the data? Does the author(s) present data to support the themes presented Does the author(s) link their findings back to the research aims? Does the author(s) consider the clinical usefulness of their findings? Does the author(s) consider the clinical usefulness of their findings? Does the author(s) consider the clinical usefulness of their findings? Does the author(s) consider the clinical usefulness of their findings? Does the author(s) identify the limitations of the research? (E.g. Sample size, recruitment strategies, method of data collection, analysis etc.) Does the author(s) identify the strengths of the research? (E.g. Its usefulness etc.) 	0 = No 1 = Partially 2 = Yes
Total	mangs:	Range 0-14

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING



Figure 1: PRISMA diagram of included and excluded studies

RER

EFFECTIVE DEMENTIA EDUCATION AND TRAINING



Figure 2: Scores on quality review tool individual criteria by overall quality of study

