



LEEDS
BECKETT
UNIVERSITY

Citation:

Weakley, J and Till, K and Read, DB and Phibbs, PJ and Roe, G and Darrall-Jones, J and Jones, BL (2017) The effects of superset configuration on kinetic, kinematic, and perceived exertion in the barbell bench press. Journal of strength and conditioning research / National Strength & Conditioning Association. ISSN 1064-8011 DOI: <https://doi.org/10.1519/JSC.0000000000002179>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4272/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

**The effects of superset configuration on kinetic, kinematic, and perceived exertion in the
barbell bench press**

Running Head: Superset configuration and kinetic, kinematic, and perceived exertion

Jonathon JS Weakley^{ab}, Kevin Till^{ab}, Dale B Read^{ab}, Padraic J Phibbs^{ab},

Gregory AB Roe^{ab}, Joshua Darrall-Jones^{ab}, and Ben L Jones^{ab}

^aInstitute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, West
Yorkshire, United Kingdom

^bYorkshire Carnegie Rugby Union Club, Kirkstall Training Ground, Leeds Rugby Academy,
Leeds, West Yorkshire, United Kingdom

Corresponding Author:

Jonathon Weakley

Room G03, Macaulay Hall

Institute for Sport, Physical Activity and Leisure

Centre for Sports Performance

Headingley Campus, Leeds Beckett University

W. Yorkshire, LS6 3GZ

Phone: (0044) 1138121801

Email: j.j.weakley@leedsbeckett.ac.uk

1 ABSTRACT

2 Training that is efficient and effective is of great importance to an athlete. One method of
3 improving efficiency is by incorporating supersets into resistance training routines. However,
4 the structuring of supersets is still unexplored. Therefore, the purpose of this study was to
5 assess the effects of agonist-antagonist (A-A), alternate peripheral (A-P), and similar
6 biomechanical (SB) superset configurations on rate of perceived exertion (RPE), kinetic and
7 kinematic changes during the bench press. 10 subjects performed resistance training protocols
8 in a randomized-crossover design, with magnitude-based inferences assessing
9 changes/differences within and between protocols. Changes in RPE were *very likely* and
10 *almost certainly* greater in the A-P and SB protocols when compared with the A-A, while all
11 superset protocols had *very likely* to *almost certain* reductions in mean velocity and power
12 from baseline. Reductions in mean velocity and power were almost certainly greater in the
13 SB protocol, with differences between the A-A and A-P protocols being *unclear*. Decreases
14 in peak force were *likely* and *almost certain* in the A-A and SB protocols respectively, with
15 changes in A-P being *unclear*. Differences between these protocols showed likely greater
16 decreases in SB peak forces when compared to A-A, with all other superset comparisons
17 being *unclear*. This study demonstrates the importance of exercise selection when
18 incorporating supersets into a training routine. It is suggested that the practitioner uses A-A
19 supersets when aiming to improve training efficiency and minimize reductions in kinetic and
20 kinematic output of the agonist musculature while completing the barbell bench press.

21

22 **Key words:** Perceived exertion; velocity; power; force

23

24

1 INTRODUCTION

2

3 Resistance training is a well-established method of improving muscular strength, power, and
4 hypertrophy (9, 28). Resistance training has been shown to be a safe method of training in
5 athletes (14, 24), and can reduce the risk of injury when appropriately implemented (15).

6 However, athletes are often unable to commit large periods of time to resistance training due
7 to other training requirements (e.g. skill development, conditioning, team practice) and the
8 large amount of variability within a training week (40). Therefore, it is important that training
9 is time efficient. One method that has been shown to enhance resistance training efficiency
10 while maintaining efficacy is the application of resistance training supersets (SS) (29). This
11 method refers to the grouping of exercises so that two different exercises are completed
12 consecutively, followed by a rest period (e.g. a squat and bench press followed by a rest
13 interval). By strategically removing recovery between sets, reductions in training time have
14 been shown to occur while training volume (e.g. total amount of weight lifted (Kg)) is
15 maintained (30, 41).

16

17 Within the literature (31), three forms of SS configuration have been defined, with agonist-
18 antagonist (A-A) pairings (e.g. bent-over row and bench press) being the most commonly
19 investigated (4, 5). Nevertheless, alternate peripheral (A-P) (e.g. back squat and bench press)
20 and similar biomechanical (SB) movements (e.g. dumbbell bench press and barbell bench
21 press) have also been examined (7, 41). However, these different exercise groupings have not
22 been compared despite evidence suggesting conflicting resistance training outcomes (e.g. in
23 the bench press throw exercise, A-A pairings caused acute increases of 4.7% in mean power
24 output (4), while SB demonstrated acute decreases in peak velocity of 10.7% (33)).

25 Consequently, the effect of SS configuration on resistance training is still unknown.

1
2 While SS are known to improve training efficiency (41), neuromuscular function may be
3 impacted when using this method. Previous research (27, 28) has shown the importance of
4 maintaining high levels of kinetic (e.g. force) and kinematic (e.g. velocity and power) outputs
5 when aiming to develop muscle size, strength, and power. However, Weakley et al. (41) has
6 demonstrated that SS may cause greater losses in these variables when compared to
7 traditional (i.e. one exercise set followed by a rest) resistance training. It was shown that SS
8 can cause large reductions in lower body power (e.g. countermovement jump height) 24
9 hours post training (effect size (ES) \pm 90% confidence interval (90% CI); SS: -0.29 ± 0.19 vs.
10 traditional: 0.01 ± 0.19), with these changes attributed to increased metabolic perturbation
11 causing reduction in force-generating capabilities. Furthermore, these reductions in
12 neuromuscular function were also accompanied by increased rating of perceived exertion
13 (RPE) (41). These increases in RPE have shown near perfect relationships with metabolic
14 responses (41) and may provide additional insight into kinematic changes when resistance
15 training (34). However, it has previously been postulated that by manipulating SS exercise
16 order and exercise selection, kinetic, kinematic, and RPE outcomes may be positively
17 affected (4). These changes have been suggested to occur due to changes in motor unit
18 recruitment (3), varying rates of localized muscle fatigue (7), and / or alterations in the
19 triphasic neural pathways (4). Nevertheless, further research is still required to assess the
20 influence of varying SS configurations on neuromuscular outcomes.

21
22 Existing research has assessed the effects of differing forms of SS on bench press kinetic and
23 kinematic outputs (4, 30, 32), RPE (5, 41), and resistance training volume (5, 30). However,
24 no study has compared the effects of these different forms of SS configuration on bench press
25 outcomes when resistance training. Furthermore, it is unknown whether these different

1 arrangements of SS alter perceptions of perceived exertion. Therefore, due to the importance
2 of kinetic and kinematic outputs when training to develop muscle size, strength, and power
3 (27, 28), the aim of this study was to assess the effects of differing SS arrangement on
4 kinetic, kinematic, and RPE changes during the bench press exercise.

5

6 **METHODS**

7 **Experimental approach to the problem**

8 To assess the effects of three different forms of SS arrangement on kinetic, kinematic, and
9 RPE outcomes in the barbell bench press, 10 sub-elite adult rugby union players completed
10 four exercise protocols in a randomized-crossover design, with at least 7 days between each
11 testing session. The four resistance training protocols consisted of one set of the barbell
12 bench press followed by a five minute recovery period then either; 1) control (CON): three
13 further sets of the barbell bench press with a two minute recovery period between sets; 2) A-
14 A: three SS of the bent-over row and the barbell bench press with two minute rest periods; 3)
15 A-P: three SS of the back squat and the barbell bench press with two minute rest periods; 4)
16 SB: three SS of the dumbbell bench press and the barbell bench press with two minute rest
17 periods. All repetitions were completed without assistance. Exercises were completed at 65%
18 of three repetition maximum (3RM) as this intensity has been found to be near maximal when
19 performing SS (33) and has previously been utilized in literature investigating this training
20 method (41). A linear position transducer (GymAware, Kinetic Performance Technology,
21 Canberra, Australia) was used to record kinetic and kinematic outcomes due to their
22 importance while resistance training (28), while RPE was reported 15 minutes after the
23 completion of each protocol owing to its validity as an internal measure of training load and
24 relationship with metabolic outcomes (17, 41).

25

1

2 **Subjects**

3 10 male rugby players (mean \pm standard deviation (SD); age: 20.9 ± 9.6 years; height: 183.2
4 ± 6.1 cm; body mass: 90.2 ± 9.6 kg; 3RM barbell bench press: 114.3 ± 10.3 kg; 3RM back
5 squat: 139.7 ± 27.9 kg; 3RM bent-over row: 101.1 ± 12.0 kg; 3RM dumbbell bench press:
6 87.5 ± 11.6 kg), who had at least two years resistance training experience (3.9 ± 1.2 years)
7 were recruited from a university rugby union team from the United Kingdom. Testing took
8 place in February (which is within the second half of the university rugby playing calendar).

9 All subjects confirmed that they did not have any current injuries, have not or do not
10 consume any medications or supplements that could influence performance, and that they
11 were not suffering from any diseases. Subjects were explained the design of the study,
12 provided an opportunity to ask questions and then provided informed written consent. All
13 experimental procedures were approved by Leeds Beckett University's ethics committee and
14 written assent was provided by all subjects.

15

16 **Experimental Procedures**

17 All testing was conducted at the same time of day one week apart, with subjects being asked
18 to refrain from physical activity for the 48 hours prior to all testing procedures. Subjects were
19 instructed to maintain normal dietary habits in the 24 hours prior to testing, with caffeine not
20 being consumed in the 12 hours before. All subjects were screened prior to acceptance into
21 the study (38), followed by anthropometric and 3RM strength assessment of the barbell bench
22 press, back squat, bent-over row, and dumbbell bench press. Subjects were randomized to
23 complete four testing sessions (i.e. CON, A-A, A-P, and SB) with session order designated
24 through computer-generated random numbering (37). All exercise protocols consisted of a
25 standardized warm-up, which consisted of dynamic movements and exercise specific

1 stretches, and a set of 10 repetitions of the bench press at 65% of 3RM performed as
2 explosively as possible prior to the experimental trials (i.e. CON, A-A, A-P, and SB). This
3 indicated the subject's baseline performance on that day. After a five-minute recovery period,
4 subjects then completed their scheduled protocol. During all repetitions and sets of the barbell
5 bench press, a linear position transducer (Kinetic Performance Technology, Canberra,
6 Australia) was attached which calculated peak force (i.e. the highest force value recorded
7 between two sampling points) and mean velocity and power (i.e. the overall velocity and
8 power output across the whole concentric range of motion) (19). These variables were used
9 due to their high level of reliability and validity (i.e. coefficient of variation <5% (6)) and
10 previous use in the barbell bench press (10). Following all exercise protocols RPE was
11 reported using a modified Borg category ratio-10 scale (16)

13 *3RM strength assessments testing*

14 The assessment of 3RM strength was completed as this is regularly used within similar
15 cohorts and has been used in the prescription of SS training methods (39-41). 3RM strength
16 testing of the barbell bench press, back squat, bent-over row, and dumbbell bench press was
17 completed during a familiarisation session after acceptance into the study. These exercises
18 were chosen due to the subjects' familiarity with these movements and their previous use in
19 rugby union research (11, 39, 40). 3RM strength of each exercise was assessed using the
20 following protocols which have previously been used to assess strength (39-41). The bench
21 press was completed with hand position at a self-selected width which was recorded and
22 replicated across conditions. The bar was lowered to the chest and returned to a locked-out
23 position to complete the repetition. The back squat was completed with the bar resting on the
24 upper trapezium with subjects required to lower themselves so that the top of the thigh was
25 parallel with the floor; as determined by the lead researcher. The bent-over row was

1 completed with an overhand grip which raised the bar to the lower sternum; while the torso
2 was maintained parallel to the ground. The dumbbell bench press begun with the subject
3 lying flat on a bench with the arms holding both dumbbells so that the elbow was at a 90-
4 degree angle. The arms were extended so that the dumbbells were directly over the subject's
5 chest and then returned to the start position.

6 7 *Superset and control protocols*

8 All protocols used the barbell bench press as an outcome measure, with SS protocols
9 requiring the completion of an exercise immediately prior, while the CON protocol
10 completed a single set of the bench press. All exercises (i.e. barbell bench press, back squat,
11 bent-over row, and dumbbell bench press) were loaded with a weight that was 65% of 3RM.
12 This intensity was selected for the protocols as it has previously been established that when
13 completing SS, intensities above this cause notable losses in repetition completion (i.e.
14 12.5%) (33). Furthermore, this is consistent with recent literature (41) which has investigated
15 the physiological responses of rugby union players. Each repetition required subjects to
16 complete a two second eccentric action, while the concentric portion of each exercise was
17 required to be as "forceful and as powerful as possible" (18, 28). Furthermore, each exercise
18 consisted of three sets of 10 repetitions, apart from the barbell bench press which had a fourth
19 set that was completed at baseline.

20 21 *Kinematic and Kinetic Assessment*

22 Assessment of mean velocity, mean power, and peak forces of the bench press were recorded
23 with a GymAware® optical encoder which sampled at 50Hz (Kinetic Performance
24 Technology, Canberra, Australia). The optical encoder, which was placed directly below the
25 barbell bench press exercise, contains a retractable cord that was attached to the barbell

1 during each set for each subject. Velocity and displacement are calculated from the rotation
2 of a pulley system within the optical encoder upon the movement of the barbell during the
3 exercise (2). The encoder, which has previously been assessed as highly valid for reporting of
4 velocity, power, and force (6), provides approximately one electrical impulse every three
5 millimetres of barbell displacement with each value time stamped with a one-millisecond
6 resolution. To assist in the calculation of the variables (i.e. power and force), barbell mass
7 and additional weight are inputted by the lead researcher before each trial (2, 12).

8

9 *RPE Measures*

10 Subjects were asked to rate their perceived exertion 15 minutes after each resistance training
11 protocol after being asked the question “How was your workout?”. Subjects were supplied
12 the modified-Borg Scale and verbally indicated an answer which was recorded. This has
13 previously been shown to have a high level of reliability in differing resistance training
14 protocols (36).

15

16 **Statistical analyses**

17 Data are presented as either mean \pm SD or percentage/effect size (ES) \pm 90% confidence
18 intervals (90% CI) where specified. Prior to analysis, all data were log-transformed to reduce
19 bias arising from non-uniformity error, and then analysed for practical significance using
20 magnitude-based inferences (21). The chance of the RPE, mean concentric velocity, power,
21 or peak force being lower, similar, or greater than the smallest worthwhile change/difference
22 (SWC/D) (i.e. 0.2 x between subject difference) was calculated using an online spreadsheet
23 (23), with all between group comparisons of effects being further analysed using a separate
24 spreadsheet (22). The probability that the magnitude of change was greater than the SWC/D
25 was rated as <0.5%, *almost certainly* not; 0.5-5%, *very unlikely*; 5-25%, *unlikely*; 25-75%,

1 possibly; 75-95%, *likely*; 95-99.5%, *very likely*; >99.5%, *almost certainly* (21). Where the
2 90% Confidence Interval (CI) crossed both the upper and lower boundaries of the SWC
3 ($ES \pm 0.2$), the magnitude of change was described as *unclear* (21). ES thresholds were set at
4 < 0.2 (*trivial*), 0.2-0.6 (*small*), 0.6-1.2 (*large*), and 1.2-2.0 (*very large*) (21).

6 RESULTS

7
8 Data are presented for mean ($\pm SD$) concentric velocity, power, and concentric peak force
9 (Table 1) of the barbell bench press in the CON and three SS protocols. Also provided are the
10 ES ($\pm 90\% CI$), inference of change from baseline, and between condition comparison of ES
11 change from baseline to set three (ES $\pm 90\% CI$ and inference) across all conditions.

12
13 ***Insert table 1 here***

14
15 RPE (mean $\pm SD$) in the CON (2.9 ± 0.8), A-A (4.0 ± 0.5), A-P (5.1 ± 0.8), SB (8.2 ± 0.7),
16 were found. All three SS protocols had *almost certainly* greater mean RPE values than the
17 CON, while the A-P condition had a *very likely* greater RPE when compared to the A-A
18 protocol. The SB protocol had an *almost certainly* greater RPE compared to both the A-A and
19 A-P protocols.

20

21

22

1 DISCUSSION

2 The purpose of the current study was to assess and compare the effects of three different SS
3 configurations on kinetic, kinematic, and RPE responses across 3 sets during the bench press
4 exercise. Of the three SS configurations, mean concentric velocity and power were reduced to
5 the greatest extent from baseline in the SB condition with *almost certainly* greater reductions
6 in this protocol when compared to the A-A and A-P pairings. Furthermore, when compared to
7 baseline, changes in peak force were *unclear* in the traditional and A-P condition, while *likely*
8 and *almost certain* reductions were reported in the A-A and SB complexes, respectively.
9 Comparisons between protocols demonstrated *likely* greater reductions in peak force in SB
10 when compared with A-A. However, differences between SB and A-P were *unclear*. Finally,
11 results revealed that the A-A pairing had *very likely* and *almost certainly* lower RPE than the
12 A-P and SB pairings, respectively.

13
14 The current study establishes the importance of SS configuration on resistance training
15 responses. Velocity loss from baseline to the final set occurred across all conditions, with
16 *moderate* changes in the CON (ES \pm 90% CI; -0.62 ± 0.29) and A-A (ES \pm 90% CI; $-0.91 \pm$
17 0.41), and *large* (ES \pm 90% CI; -1.45 ± 0.88) to *very large* (ES \pm 90% CI; -6.17 ± 0.57)
18 reductions in the A-P and SB. While *possible* and *likely* differences were evident when CON
19 was compared with the A-A and A-P protocols, respectively, it is of note that *unclear*
20 differences were evident between the two latter conditions. This *unclear* result is partially
21 attributed to the large amount of uncertainty around the mean loss in velocity in the A-P
22 pairing. It is speculated that this uncertainty is due to varying tolerance of the large metabolic
23 cost of completing the back squat immediately followed by the bench press (40). With near
24 perfect relationships between measures of fatigue (e.g. lactate and ammonia accumulation) (r
25 = 0.95-0.97) and velocity loss in the barbell bench press (34), subjects with greater lower

1 body work capacity may have shown improved tolerance to this form of SS configuration.
2 Therefore, a high level of work capacity may be required to reduce loss of kinematic
3 outcomes in SS complexes that incorporate the lower body and its large component of lean
4 body mass.

5
6 The *very large* losses of velocity reported in the SB protocol were *almost certainly* greater
7 than reductions in all other protocols. Previous research (18) has shown that repeated high-
8 intensity muscle contractions can reduce substrate availability and increase metabolic
9 accumulation. These changes can cause acute reductions in muscle performance which have
10 been shown to impede moderate term (i.e. 8 weeks) strength and power development (28).
11 However, greater reductions in barbell velocity across these time frames have been related to
12 increased muscle hypertrophy (28). This promotes the idea that SS complexes that pair
13 exercises of a similar nature may not be ideal for the development of strength and power (8,
14 28, 31), yet may assist in the development of lean body mass due to responses related to
15 increased peripheral fatigue (31, 35). However, longitudinal research investigating this is still
16 required to elucidate these mechanisms and changes.

17
18 Accompanying the previously reported decreases in velocity, were almost identical losses in
19 power in the SS protocols (see table 1). However, the *moderate* (A-A), *large* (A-P), and *very*
20 *large* (SB) decreases in power from baseline do not mirror the much smaller changes in peak
21 force production. This suggests that the implementation of SS resistance training causes
22 reduction in power output primarily due to decreases in velocity rather than force, with these
23 changes in velocity most likely attributed to mitigated shortening speeds of muscle fibers and
24 the slowing of muscle relaxation (1). While reductions in peak forces do occur, albeit to a
25 lesser extent than mean velocity and power, these smaller changes may be due to an increase

1 in motor unit efficiency in the triceps brachii (3). Artur et al. (3) have demonstrated the
2 additional recruitment of higher threshold motor units when fatigue is induced prior to the
3 bench press exercise. However, it should be noted that this increased motor unit recruitment
4 may not promote long-term improvements in force and power (25, 26, 28). The practitioner
5 and sports scientist may therefore be cautious in the selection of SB pairings due to the large
6 amounts of localized fatigue, declines in kinetic and kinematic variables, and the mechanisms
7 that are utilized to maintain performance.

8
9 The current study also acknowledges the importance of SS configuration on perceived
10 intensity, with the A-A pairing reporting lower RPE when compared with A-P and SB. While
11 this study is not the first to assess SS configuration and RPE (5), it is the first to compare
12 supersets that extend beyond the agonist and antagonist format. It is speculated that the
13 greater range of movement and muscle mass utilized in the A-P SS (13), and the increased
14 localized muscle fatigue and subsequent substrate depletion in the SB protocol (8), increased
15 RPE and impacted performance. This indicates that perceived intensity may not only be
16 related to exercise choice or intensity, but total work completed (i.e. displacement of the
17 external load) and the total volume each muscle group completes (20). Therefore, when
18 implementing SS, consideration should be given to the movement pattern and range of
19 motion each exercise requires and how this can impact upon perceived measures of training
20 load.

21
22 While this study is the first to assess and compare the varying effects of different SS
23 configuration, it is not without its limitations that might reduce transferability to application.
24 First, while the definition of different forms of SS have been referred to in numerous
25 publications (3, 8, 31), the formulation of each SS complex within a defined category can

1 occur in a large variety of ways (41). This includes exercise selection (e.g. multi-
2 joint/isolated), order, intensity (as a percentage of 1RM), range of motion, and recovery
3 allotted between subsequent sets. It is therefore crucial that the practitioner and sport scientist
4 consider these training variables when designing relevant SS interventions. Secondly, due to
5 the nature of the research question, the outcomes of this study were being assessed within a
6 singular exercise (i.e. barbell bench press). Previous research has assessed neuromuscular and
7 metabolic responses across acute and short-term time periods (41). However, further research
8 is required to assess the outcomes of longitudinal research investigating varying
9 configurations of SS complexes. Finally, it is acknowledged that kinetic and kinematic
10 responses demonstrated in the current study may be exercise and loading specific. Complexes
11 of a similar nature may vary due to muscle mass utilized (34), terminal concentric velocity of
12 individual exercises (18), contribution of the stretch-shortening cycle (13), and the relative
13 “sticking region” of each exercise (34). These findings therefore need to be interpreted with
14 caution for exercises other than the barbell bench press.

15
16 In conclusion, the current study demonstrates the kinetic, kinematic and perceived exertion
17 responses to differing SS structures. The findings suggest that SS configuration can induce
18 varying levels of acute fatigue, with movements that are of a SB nature resulting in the
19 largest decline in performance. Additionally, resistance training that utilizes A-A and A-P
20 may demonstrate similar changes in performance outcomes. However, due to the large
21 amount of deviation in individual responses to A-P, as well as the *very likely* smaller change
22 in rate of perceived exertion in the A-A condition, protocols that integrate “*pull-push*” SS
23 complexes may be favourable in time constrained periods. Finally, TRAD training structures
24 that incorporate increased recovery may be most beneficial when training objectives are to

1 maximize movement velocity and power. Future research should endeavour to apply the
2 above findings and assess how these results can enhance training outcomes.

3

4 **Practical Application**

5 The ability to spend prolonged amounts of time training is often not feasible for an athlete.
6 Therefore, structures that enhance training efficiency by decreasing training time should be
7 considered. The implementation of SS is a unique training method that can enhance
8 resistance training efficiency, with A-A and A-P configurations both appearing to minimize
9 declines in kinetic and kinematic outcomes when compared with SB. Consequently, these
10 configurations should be utilized when aiming to reduce recovery interval frequency, or when
11 large amounts of training volume are required (e.g. muscular endurance/hypertrophy
12 mesocycles). Furthermore, the use of A-A SS may mitigate increases in perceived exertion
13 which may be beneficial during these training phases. However, the selection of SS
14 configuration may depend upon the desired training outcome and it should be noted that
15 traditional resistance training methods (i.e. singular sets) were superior in the maintenance of
16 velocity, power, and force. Therefore, when resistance training quality (i.e. high levels of
17 kinetic and kinematic outputs) is of the highest importance (e.g. during the development of
18 power) traditional methods may be of the greatest benefit.

19

20

21

22

23

24

25

26

27

1 References

- 2 1. Allen DG, Lamb GD, and Westerblad H. Skeletal muscle fatigue: cellular
3 mechanisms. *Physiol Rev* 88: 287-332, 2008.
- 4 2. Argus CK, Gill ND, Keogh JW, and Hopkins WG. Acute effects of verbal feedback
5 on upper-body performance in elite athletes. *J Strength Cond Res* 25: 3282-3287,
6 2011.
- 7 3. Artur G, Adam M, Przemyslaw P, Stastny P, James T, and Adam Z. Effects of pre-
8 exhaustion on the patterns of muscular activity in the flat bench press. *J Strength*
9 *Cond Res (ahead of press)* 10.1519/JSC.0000000000001755.
- 10 4. Baker D and Newton RU. Acute effect on power output of alternating an agonist and
11 antagonist muscle exercise during complex training. *J Strength Cond Res* 19: 202-
12 205, 2005.
- 13 5. Balsamo S, Tibana RA, Nascimento DdC, de Farias GL, Petruccelli Z, de Santana
14 FdS, Martins OV, de Aguiar F, Pereira GB, de Souza JC, and Prestes J. Exercise order
15 affects the total training volume and the ratings of perceived exertion in response to a
16 super-set resistance training session. *Int J Gen Med* 5: 123-127, 2012.
- 17 6. Banyard HG, Nosaka K, Sato K, and Haff GG. Validity of Various Methods for
18 Determining Velocity, Force and Power in the Back Squat. *Int J Sports Physiol*
19 *Perform*: 1-25, 2017.
- 20 7. Brennecke A, Guimaraes TM, Leone R, Cadarci M, Mochizuki L, Simao R, Amadio
21 AC, and Serrao JC. Neuromuscular activity during bench press exercise performed
22 with and without the preexhaustion method. *J Strength Cond Res* 23: 1933-1940,
23 2009.
- 24 8. Castanheira RP, Ferreira-Junior JB, Celes R, Rocha-Junior VA, Cadore EL, Izquierdo
25 M, and Bottaro M. Effects of synergist vs. Non-synergist split resistance training
26 routines on acute neuromuscular performance in resistance trained men. *J Strength*
27 *Cond Res (ahead of press)* DOI: 10.1519/JSC.0000000000001762.
- 28 9. Cormie P, McGuigan MR, and Newton RU. Adaptations in athletic performance after
29 ballistic power versus strength training. *Med Sci Sports Exerc* 42: 1582-1598, 2010.
- 30 10. Cronin JB, Mcnair PJ, and Marshall RN. Force-velocity analysis of strength-training
31 techniques and load: implications for training strategy and research. *J Strength Cond*
32 *Res* 17: 148-155, 2003.
- 33 11. Darrall-Jones JD, Jones B, and Till K. Anthropometric and Physical Profiles of
34 English Academy Rugby Union Players. *J Strength Cond Res* 29: 2086-2096, 2015.
- 35 12. Drinkwater EJ, Galna B, McKenna MJ, Hunt PH, and Pyne DB. Validation of an
36 optical encoder during free weight resistance movements and analysis of bench press
37 sticking point power during fatigue. *J Strength Cond Res* 21: 510-517, 2007.
- 38 13. Drinkwater EJ, Moore NR, and Bird SP. Effects of changing from full range of
39 motion to partial range of motion on squat kinetics. *J Strength Cond Res* 26: 890-896,
40 2012.
- 41 14. Faigenbaum AD, Kraemer WJ, Blimkie CJ, Jeffreys I, Micheli LJ, Nitka M, and
42 Rowland TW. Youth resistance training: updated position statement paper from the
43 national strength and conditioning association. *J Strength Cond Res* 23: S60-S79,
44 2009.
- 45 15. Faigenbaum AD and Myer GD. Resistance training among young athletes: safety,
46 efficacy and injury prevention effects. *Bri J Sports Med* 44: 56-63, 2010.
- 47 16. Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, Doleshal P,
48 and Dodge C. A new approach to monitoring exercise training. *J Strength Cond Res*
49 15: 109-115, 2001.

- 1 17. Gearhart RF, Jr., Goss FL, Lagally KM, Jakicic JM, Gallagher J, Gallagher KI, and
2 Robertson RJ. Ratings of perceived exertion in active muscle during high-intensity
3 and low-intensity resistance exercise. *Journal of strength and conditioning research*
4 16: 87-91, 2002.
- 5 18. Gonzalez-Badillo JJ and Sanchez-Medina L. Movement velocity as a measure of
6 loading intensity in resistance training. *Int J Sports Med* 3: 347-352, 2010.
- 7 19. Harris NK, Cronin J, Taylor K-L, Boris J, and Sheppard J. Understanding position
8 transducer technology for strength and conditioning practitioners. *Strength Cond J* 32:
9 66-79, 2010.
- 10 20. Hiscock DJ, Dawson B, Clarke M, and Peeling P. Can changes in resistance exercise
11 workload influence internal load, countermovement jump performance and the
12 endocrine response? *J Sports Sci*: 1-7, 2017.
- 13 21. Hopkins W, Marshall S, Batterham A, and Hanin J. Progressive statistics for studies
14 in sports medicine and exercise science. *Med Sci Sports Exerc* 41: 3, 2009.
- 15 22. Hopkins WG. A spreadsheet for combining outcomes from several subject groups.
16 *Sportsci* 10: 50-53, 2006.
- 17 23. Hopkins WG. Spreadsheets for analysis of controlled trials with adjustment for a
18 predictor. *Sportsci* 10: 46-50 2006.
- 19 24. Kraemer WJ, Adams K, Cafarelli E, Dudley GA, Dooly C, Feigenbaum MS, Fleck SJ,
20 Franklin B, Fry AC, and Hoffman JR. American College of Sports Medicine position
21 stand. Progression models in resistance training for healthy adults. *Med Sci Sports*
22 *Exerc* 34: 364-380, 2002.
- 23 25. Newham D, McCarthy T, and Turner J. Voluntary activation of human quadriceps
24 during and after isokinetic exercise. *J App Physiol* 71: 2122-2126, 1991.
- 25 26. Nyland JA, Caborn DN, Shapiro R, and Johnson DL. Fatigue after eccentric
26 quadriceps femoris work produces earlier gastrocnemius and delayed quadriceps
27 femoris activation during crossover cutting among normal athletic women. *Knee Surg*
28 *Sports Traumatol Arthrosc* 5: 162-167, 1997.
- 29 27. Pareja-Blanco F, Rodriguez-Rosell D, Sanchez-Medina L, Gorostiaga EM, and
30 Gonzalez-Badillo JJ. Effect of movement velocity during resistance training on
31 neuromuscular performance. *Int J Sports Med* 35: 916-924, 2014.
- 32 28. Pareja - Blanco F, Rodríguez - Rosell D, Sánchez - Medina L, Sanchis - Moysi J,
33 Dorado C, Mora - Custodio R, Yáñez - García J, Morales - Alamo D, Pérez - Suárez
34 I, and Calbet J. Effects of velocity loss during resistance training on athletic
35 performance, strength gains and muscle adaptations. *Scand J Med Sci Sports* (ahead
36 of press) DOI: 10.1111/sms.12678.
- 37 29. Robbins DW, Young WB, Behm DG, and Payne WR. Effects of agonist-antagonist
38 complex resistance training on upper body strength and power development. *Journal*
39 *of sports sciences* 27: 1617-1625, 2009.
- 40 30. Robbins DW, Young WB, and Behm DG. The effect of an upper-body agonist-
41 antagonist resistance training protocol on volume load and efficiency. *J Strength*
42 *Cond Res* 24: 2632-2640, 2010.
- 43 31. Robbins DW, Young WB, Behm DG, and Payne WR. Agonist-antagonist paired set
44 resistance training: a brief review. *J Strength Cond Res* 24: 2873-2882, 2010.
- 45 32. Robbins DW, Young WB, Behm DG, Payne WR, and Klimstra MD. Physical
46 performance and electromyographic responses to an acute bout of paired set strength
47 training versus traditional strength training. *J Strength Cond Res* 24: 1237-1245,
48 2010.

- 1 33. Sabido R, Peñaranda M, and Hernández-Davó JL. Comparison of acute responses to
2 four different hypertrophy-oriented resistance training methodologies. *Eur J Hum*
3 *Mov* 37: 109-121, 2016.
- 4 34. Sanchez-Medina L and González-Badillo JJ. Velocity loss as an indicator of
5 neuromuscular fatigue during resistance training. *Med Sci Sports Exerc* 43: 1725-
6 1734, 2011.
- 7 35. Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to
8 resistance training. *J Strength Cond Res* 24: 2857-2872, 2010.
- 9 36. Singh F, Foster C, Tod D, and McGuigan MR. Monitoring different types of
10 resistance training using session rating of perceived exertion. *Int J Sports Physiol*
11 *Perform* 2: 34-45, 2007.
- 12 37. Suresh KP. An overview of randomization techniques: An unbiased assessment of
13 outcome in clinical research. *J Hum Reprod Sci* 4: 8-11, 2011.
- 14 38. Thompson PD, Arena R, Riebe D, and Pescatello LS. ACSM's New Preparticipation
15 Health Screening Recommendations from ACSM's Guidelines for Exercise Testing
16 and Prescription, Ninth Edition. *Curr Sports Med Rep* 12: 215-217, 2013.
- 17 39. Weakley J, Till K, Darrall-Jones J, Roe G, Phibbs P, Read D, and Jones B. The
18 influence of resistance training experience on the between-day reliability of
19 commonly used strength measures in male youth athletes. *J Strength Cond Res*,
20 DOI:10.1519/JSC.0000000000001883.
- 21 40. Weakley JJS, Till K, Darrall-Jones J, Roe GAB, Phibbs PJ, Read DB, and Jones BL.
22 Strength and Conditioning Practices in Adolescent Rugby Players: Relationship with
23 Changes in Physical Qualities. *J Strength Cond Res* (ahead of press) DOI:
24 10.1519/JSC.0000000000001828.
- 25 41. Weakley JJS, Till K, Read DB, Roe GAB, Darrall-Jones J, Phibbs PJ, Jones BL
26 (2017) The effects of traditional, superset, and tri-set resistance training structures on
27 perceived intensity and physiological responses *Eur J App Physiol* (ahead of press)
28 DOI: 10.1007/s00421-017-3680-3

29

30

Table 1. Within and between condition comparison of barbell bench press mean velocity ($\text{m}\cdot\text{s}^{-1}$), power (W), and peak force (N) across traditional and superset resistance training protocols

ACCEPTED

	Baseline M ± SD	Set 1 M ± SD	Baseline – Set 1 ES ± 90% CI Inference	Set 2 M ± SD	Baseline – Set 2 ES ± 90% CI Inference	Set 3 M ± SD	Baseline – Set 3 ES ± 90% CI Inference	Baseline – Set 3 Between condition comparison ES ± 90% CI / Inference
<u>Mean Velocity (m·s⁻¹)</u>								
Control	0.70 ± 0.10	0.70 ± 0.10	0.01 ± 0.21 <i>Likely trivial</i>	0.67 ± 0.10	-0.25 ± 0.24 <i>Possibly ↓</i>	0.64 ± 0.09	-0.62 ± 0.29 <i>Very likely ↓</i>	-0.29 ± 0.46 / A-A <i>possibly</i> greater ↓ -0.83 ± 0.90 / A-P <i>likely</i> greater ↓ -5.55 ± 0.94 / SB <i>almost certainly</i> greater ↓
Agonist-antagonist	0.70 ± 0.13	0.68 ± 0.15	-0.18 ± 0.19 <i>Possibly ↓</i>	0.64 ± 0.18	-0.51 ± 0.32 <i>Likely ↓</i>	0.59 ± 0.19	-0.91 ± 0.41 <i>Very likely ↓</i>	-0.54 ± 0.94 / A-A and A-P <i>Unclear</i> -5.26 ± 0.67 / SB <i>almost certainly</i> greater ↓
Alternate peripheral	0.70 ± 0.09	0.67 ± 0.13	-0.23 ± 0.39 <i>Possibly ↓</i>	0.61 ± 0.15	-0.89 ± 0.59 <i>Very likely ↓</i>	0.57 ± 0.16	-1.45 ± 0.88 <i>Very likely ↓</i>	-4.72 ± 1.00 / SB <i>almost certainly</i> greater ↓
Similar Biomechanical	0.74 ± 0.09	0.54 ± 0.15	-2.63 ± 0.74 <i>Almost certain ↓</i>	0.43 ± 0.15	4.66 ± 1.03 <i>Almost certain ↓</i>	0.36 ± 0.07	-6.17 ± 0.57 <i>Almost certain ↓</i>	
<u>Mean Power (W)</u>								
Control	509 ± 66	510 ± 68	-0.02 ± 0.22 <i>Unclear</i>	492 ± 62	-0.26 ± 0.26 <i>Possibly ↓</i>	485 ± 118	-0.36 ± 0.54 <i>Possibly ↓</i>	-0.63 ± 0.67 / A-A <i>possibly</i> greater ↓ -1.24 ± 1.14 / A-P <i>likely</i> greater ↓ -7.37 ± 0.89 / SB <i>almost certainly</i> greater ↓
Agonist-antagonist	507 ± 87	488 ± 101	-0.22 ± 0.21 <i>Possibly ↓</i>	458 ± 109	-0.59 ± 0.35 <i>Very likely ↓</i>	425 ± 121	-1.02 ± 0.45 <i>Almost certain ↓</i>	-0.61 ± 1.11 / A-A and A-P <i>Unclear</i> -6.74 ± 0.85 / SB <i>almost certainly</i> greater ↓
Alternate peripheral	501 ± 62	482 ± 92	-0.31 ± 0.52 <i>Unclear</i>	440 ± 100	-1.02 ± 0.73 <i>Very likely ↓</i>	407 ± 111	-1.63 ± 1.05 <i>Very likely ↓</i>	-6.13 ± 1.23 / SB <i>almost certainly</i> greater ↓
Similar Biomechanical	539 ± 51	398 ± 117	-3.08 ± 1.16 <i>Almost certain ↓</i>	303 ± 101	-5.83 ± 1.26 <i>Almost certain ↓</i>	250 ± 47	-7.76 ± 0.76 <i>Almost certain ↓</i>	
<u>Peak Force (N)</u>								
Control	1258 ± 126	1271 ± 146	0.10 ± 0.25 <i>Possibly trivial</i>	1295 ± 170	0.28 ± 0.33 <i>Possibly ↓</i>	1286 ± 170	0.21 ± 0.42 <i>Unclear</i>	-0.59 ± 0.83 / CON and A-A <i>unclear</i> -0.73 ± 0.85 / A-P <i>likely</i> greater ↓ -1.39 ± 0.59 / SB <i>almost certainly</i> greater ↓
Agonist-antagonist	1381 ± 191	1365 ± 225	-0.08 ± 0.13 <i>Likely trivial</i>	1341 ± 209	-0.21 ± 0.18 <i>Possibly ↓</i>	1309 ± 212	-0.38 ± 0.22 <i>Likely ↓</i>	-0.14 ± 1.03 / A-A and A-P <i>Unclear</i> -0.80 ± 0.85 / SB <i>likely</i> greater ↓
Alternate peripheral	1349 ± 225	1355 ± 215	0.02 ± 0.19 <i>Likely trivial</i>	1313 ± 215	-0.16 ± 0.22 <i>Possibly ↓</i>	1235 ± 211	-0.52 ± 0.77 <i>Unclear</i>	-0.66 ± 0.86 / A-P and SB <i>unclear</i>
Similar Biomechanical	1447 ± 139	1329 ± 193	-0.85 ± 0.38 <i>Very likely ↓</i>	1310 ± 145	-0.99 ± 0.33 <i>Almost certain ↓</i>	1286 ± 1285	-1.18 ± 0.46 <i>Almost certain ↓</i>	

M ± SD: Mean ± standard deviation. ES: Effect size. 90% CI: 90% confidence interval. ↑: increase. ↓: decrease. CON: control. A-A: agonist-antagonistic pairing; A-P: alternate peripheral pairing; S-B: similar biomechanical pairing.