

---

Citation:

Tee, JC and Till, K and Jones, B (2017) Effects of an intensified competition period on neuromuscular function in youth rugby union players. Sport Performance and Science Reports.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4372/>

Document Version:

Article (Published Version)

---

Creative Commons: Attribution 4.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

# Effects of an intensified competition period on neuromuscular function in youth rugby union players

Jason C. Tee<sup>1</sup>, Kevin Till<sup>1</sup>, Ben Jones<sup>1</sup>

<sup>1</sup> Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, West Yorkshire, United Kingdom

Neuromuscular Performance | Recovery | Game | Rugby union

## Headline

Playing rugby union matches causes a number of fatigue responses, including reduced lower body neuromuscular function (NMF) (commonly measured by counter movement jump (CMJ))(1). The time course of this response following match play is well established in professional (2) and academy (3) level rugby union players, who take at least 60 hours for NMF to recover. No data exist for high school level rugby union players, but these players are often exposed to multiple game tournaments and festivals (2 games in 3 days, or 3 games in 5 days) within their competition structures.

**Aim.** The aim of this case study is to document the NMF response to playing three rugby union matches within five days. This will provide useful information to practitioners who must manage fatigue and recovery of youth rugby union players who play multiple games within short time periods.

## Methods

**Athletes.** Fifteen male rugby union players (age  $17.2 \pm 0.8$  years; height  $178 \pm 8$  cm; body mass  $87.1 \pm 15.2$  kg; 1 repetition maximum (RM) Bench Press  $105 \pm 20$  kg; 5RM Squat  $139 \pm 24$  kg; 40 m sprint  $5.5 \pm 0.2$  s) competing in a high school rugby festival over a 5-day period volunteered to take part in the study. Ethics approval was granted by Leeds Beckett University, and permission to use the previously collected data was granted by the players' school.


**Design.** A within-group repeated measures design was used to examine the magnitude of change in NMF throughout the festival. The players participated in three 70-minute rugby matches over the course of five days. The matches kicked off at 13:15 on match day 1, and at 15:00 on match days 2 and 3. Tactical substitutions were not employed, and players were only substituted in the case of injury. As a result, the starting fifteen players played the majority (>90%) of the playing time available. Immediately after each match players performed a combined cold-water immersion and active recovery protocol. Players entered a waist deep swimming pool (approx. temp.  $10^{\circ}\text{C}$ ) and spent approximately 5 minutes walking or jogging in the water. A single recovery day was scheduled between each match. Low intensity training sessions, consisting of 35 minutes of stretching, walking, jogging and skill execution (catch/pass etc.) activity took place at 11:00 each morning on the days between matches. NMF was measured by determining CMJ flight time each day two hours prior to competition or one hour prior to training. Two players were injured during the second match and were unable to take further part in the study. As such, the data set for the final two days is for just 13 players.

**Methodology.** The CMJ's were performed on an electronic timing mat system (Just Jump System, Probotics, Inc., Huntsville, AL). CMJs were conducted according to the procedures described by Twist et al. (2012) (4). Players were familiar with this methodology as they regularly performed counter movement jumps in pre-season testing sessions and during training. The between day reliability of the CMJ flight time measure in this population was acceptable ( $\text{CV} = 2.14\%$ ).

**Statistical analysis.** Changes in group CMJ flight time were analysed for practical significance using magnitude-based inferences<sup>5</sup>. The threshold for change considered to be practically important (the smallest worthwhile change; SWC) was set at  $0.2 \times$  between subject standard deviation (SD), based on Cohen's d effect size (ES) principle. The probability that the magnitude of change was greater than the SWC was rated as <0.5%, almost certainly not; 0.5-5%, very unlikely; 5-25%, unlikely; 25-75%, possibly; 75-95%, likely; 95-99.5%, very likely; >99.5%, almost certainly<sup>4</sup>. Where the 90 % Confidence Interval (CI) crossed both the upper and lower boundaries of the SWC ( $\text{ES} \pm 0.2$ ), the magnitude of change was described as unclear (5). Individual changes in CMJ flight time were considered clearly meaningful if the individual mean change  $\pm$  the typical error of measurement (TEM) clearly exceeded the SWC (6). Individual TEM was calculated as the within-subject standard deviation (7).

## Discussion

Rugby union is a physically demanding sport (8). Full recovery of NMF following match participation requires at least 60 hours(2,3). The effects of training or competing again within



**Fig. 1.** Percentage change in counter movement jump flight time over a five-day rugby festival. Bold lines indicate group mean with 90% confidence intervals. Grey lines indicate individual participant scores. Greyed out area indicates the thresholds for smallest worthwhile change. Analysis indicates that there were likely negative effects of match participation on flight time on recovery day 1 and recovery day 2. However, 48 hours following the second game the effects were unclear.

this 60-hour window are unclear. Preliminary results suggest that high training loads within the recovery period extend the time required for full recovery (9,10).

Rugby union injuries occur most frequently in the final quarter of matches (11) indicating that fatigue may be a risk factor for rugby injury. Research in professional football indicates that injury risk is higher when players have less than 5 days recovery between matches (12). Youth rugby players are often exposed to multiple game tournaments and festivals (2 games in 3 days, or 3 games in 5 days) within their competition structures. Youth rugby players also often play for multiple teams (e.g. schools, clubs and academies) leading to multiple match exposures within the same week across different competition structures (13). Therefore it is important to understand how participation in periods of intensified competition may affect fatigue indicators. To the author's knowledge, this is the first study to examine the effects of an intensified competition period on NMF in youth rugby union players.

The results of this case study indicate that the majority of players did not display any residual neuromuscular fatigue 48 hours following participation in the first match. Furthermore, no clear group effect of playing two matches in three days was evident 48 hours following participation in Match 2 (Figure 1). This is similar to previous research that showed that NMF returned to baseline 24 hours after playing five games in four days in junior rugby league players (14). However, there was a great deal of variability in fatigue response between players, and four players displayed clearly reduced NMF

on match day 3 (Figure 2). This demonstrates that at least some players were meaningfully fatigued as a result of the intensified competition exposure. The fatigue response results from the metabolic stress of repeated high-intensity actions, the mechanical trauma of repeated collisions, or a combination of both (2). Therefore the observed variability in fatigue response may be the result of both differing positional requirements (8), and differences in individual within match playing involvements (number of tackles, ball carries etc.). Due the likelihood of reduced match performance (14,15) and possibly increased injury risk (11), this phenomenon warrants close monitoring in the applied setting.

## Practical Applications

- CMJ testing can be used to determine which individual players are exhibiting signs of neuromuscular fatigue during periods of intensified rugby union competition.
- Provides reference data regarding the magnitude and time course of changes in NMF for players involved in periods of intensified competition.
- Illustrates the importance of implementing between game recovery strategies for players involved in periods of intensified competition.

## Limitations

- Fatigue is multi-factorial in nature<sup>16</sup>, and this research has only measured one indirect indicator of fatigue (NMF). Future research should aim to include a range of fatigue measures including biochemical markers of muscle damage, perceptual fatigue and upper body NMF.
- No measure of internal or external load was collected, making it impossible to link NMF outcomes to physical exertions.
- It would have been better to use a force plate rather than a timing mat to measure NMF. Mean power and peak force are demonstrably more sensitive measures of NMF (7), and the inclusion of these may have revealed more meaningful group effects.
- The effect of the recovery methods employed in this study cannot be quantified because no control group was available.

**Acknowledgments.** The staff and players of King Edward VII School are thanked for their support of this project.


## Dataset

Dataset available on SportPerfSci.com

**Twitter:** Follow Jason Tee @JasonCTee, Kevin Till @ktconditioning and Ben Jones @23Benjones.

## References

1. Twist C, Highton J. Monitoring Fatigue and Recovery in Rugby Players. In: Twist C, Worsfold P, editors. *The Science of Rugby*. Routledge; 2015. p. 68-82.
2. West DJ, Finn C, Cunningham DJ, Shearer DA, Jones MR, Harrington B, Crewther BT, Cook CJ, Kilduff LP. The neuromuscular function, hormonal, and mood responses to a professional rugby union match. *J Strength Cond Res*. 2014 Jan;28(1): 194-200.
3. Roe G, Till K, Darrall-Jones J, Phibbs P, Weakley J, Read D, Jones B. Changes in markers of fatigue following a com-



**Fig. 2.** Individual change from baseline in counter movement jump flight time on match day 2 (a) and match day 3 (b) of a five day rugby festival. Error bars indicate individual typical error of measurement. Dashed grey lines indicate the thresholds for smallest worthwhile change. Between subjects responses were highly variable. On match day 2 (Figure 2a) no clearly negative changes were apparent, while one player had substantially improved his jump flight time. On match day 3 (figure 2b), four players displayed clearly reduced jump flight time from pre-tournament values.

- petitive match in elite academy rugby union players. *South African Journal of Sports Medicine*. 2017 Jan;28(1):1-4.
4. Twist C, Waldron M, Highton J, Burt D, Daniels M. Neuromuscular, biochemical and perceptual post-match fatigue in professional rugby league forwards and backs. *J Sports Sci*. 2012 Feb;30(4): 359-367.
  5. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc*. 2009 Jan;41(1):3-13.
  6. Hopkins WG. How to interpret changes in an athletic performance test. *Sportscience*. 2004;8(1):1-7.
  7. Roe G, Darrall-Jones J, Till K, Phibbs P, Read D, Weakley J, Jones B. Between-Days Reliability and Sensitivity of Common Fatigue Measures in Rugby Players. *Int J Sports Physiol Perform*. 2016 Jul;11(5):581-586.
  8. Deutsch MU, Kearney GA, Rehrer NJ. Time - motion analysis of professional rugby union players during match-play. *J Sports Sci*. 2007 Feb;25(4):461-472.
  9. McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular, endocrine, and perceptual fatigue responses during different length between-match microcycles in professional rugby league players. *Int. J Sports Physiol Perform*. 2010 Sep;5(3):367-83
  10. Higgins TR, Climstein DA, Cameron M. Evaluation of hydrotherapy, using passive tests and power tests, for recovery across a cyclic week of competitive rugby union. *J Strength Cond Res*. 2013 Apr;27(4):954-65
  11. Williams S, Trewartha G, Kemp S, Stokes K. A meta-analysis of injuries in senior men's professional Rugby Union. *Sports Med*. 2013 Oct;43(10):1043-55.
  12. Bengtsson H, Ekstrand J, Waldén M, Häggglund M. Muscle injury rate in professional football is higher in matches played within 5 days since the previous match: a 14-year prospective study with more than 130 000 match observations. *Br J Sports Med*. 2017 Nov;[Epub ahead of print]
  13. Phibbs PJ, Jones B, Roe G, Read D, Darrall-Jones J, Weakley J, Rock A, Till K. Organised chaos in late specialisation team sports: Weekly training loads of elite adolescent rugby union players. *J Strength Cond Res*. 2017 Apr;[Epub ahead of print]
  14. Johnston RD, Gabbett TJ, and Jenkins DG. Influence of an intensified competition on fatigue and match performance in junior rugby league players. *J Sci Med Sport*. 2013 Sep;16(5):460-5.
  15. Johnston RD, Gibson NV, Twist C, Gabbett TJ, MacNay SA, MacFarlane NG. Physiological responses to an intensified period of rugby league competition. *J Strength Cond Res*. 2013 Mar; 27(3): 643-654.
  16. Enoka RM, Duchateau J. Translating Fatigue to Human Performance. *Med Sci Sports Exerc*. 2016 Nov;48(11):2228-38.

**Copyright:** The articles published on Science Performance and Science Reports are distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.