



LEEDS
BECKETT
UNIVERSITY

Citation:

Jones, BL and Till, K and Manley, A (2015) Advancing the profiling of athletes: Incorporating coach-athlete collaboration when interpreting fitness testing data. In: The British Association of Sport and Exercise Sciences (BASES) conference, 01 December 2015 - 02 December 2015, St George's Park, Burton upon Trent, UK.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4721/>

Document Version:

Conference or Workshop Item (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

**A Multidisciplinary Approach to the Profiling and Interpretation of Fitness Testing
Data: A Case Study Example**

Ben Jones^{1,2,3}, Kevin Till^{1,2,3}, Andrew J. Manley¹, Michael R. McGuigan⁴

Running Title: Multidisciplinary Approach to the Interpretation of Testing Data

¹Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, West Yorkshire, United Kingdom

²Leeds Rhinos Rugby League Club, Headingley Stadium, Leeds, LS6 3BR

³Yorkshire Carnegie Rugby Union Club, Headingley Stadium, Leeds, LS6 3BR

⁴Sports Performance Research Institute New Zealand, Auckland University of Technology, Auckland, New Zealand

Dr Ben Jones – B.Jones@leedsbeckett.ac.uk

Dr Kevin Till – K.Till@leedsbeckett.ac.uk

Dr Andrew Manley – A.J.Manley@leedsbeckett.ac.uk

Prof. Michael McGuigan - Michael.McGuigan@aut.ac.nz

Corresponding Author:

Dr. Ben Jones

Room 112, Fairfax Hall

Institute for Sport, Physical Activity and Leisure

Centre for Sports Performance

Headingley Campus, Leeds Beckett University

W.Yorkshire, LS6 3QS

Phone: (044-11) 01132-832600 Ext: 24009

“When profiling athletes, it is proposed that multiplying a modified Z-score by a factor based on relative importance of each characteristic may prove a useful tool for practitioners.”

ABSTRACT

Fitness testing is commonly used to assess athletes' strengths and weaknesses. However, limitations exist including lack of position-specific comparative data, and little consideration for athletes' or coaches' needs. This article proposes a collaborative method for interpreting fitness testing data to address the practical limitations identified above. Data obtained from three male professional academy rugby league players (age 17.7 ± 0.6 years; height 180.1 ± 6.7 cm; body mass 88.8 ± 4.6 kg) were used for the purpose of this article. Following the fitness testing battery, in collaboration with each athlete, three professional coaches ranked the relative importance of each test (1 = least important; 7 = most important) for each player considering playing position, stage of physical and technical / tactical development, and future goals. Data were presented as absolute values, Z-scores and modified (benchmark; mean + 2 *SDs*) Z-scores in comparison to previously published data. The different approaches of data presentation identified different goals for each athlete. This article provides a novel approach for presenting and interpreting fitness testing data, considering athlete and coach needs. When profiling athletes, it is proposed that multiplying a modified Z-score by a factor based on relative importance of each characteristic may prove a useful tool for practitioners. In practice this process allows athletes and coaches to specifically focus on individual weaknesses, which are identified as being important for an athlete's particular role in their sport or their stage of development.

Key Words: Rugby, Strength, Speed, Athletic Development

INTRODUCTION

Fitness testing of athletes is common practice within strength and conditioning. Strength and conditioning coaches typically implement a testing battery to determine the characteristics of an athlete (6, 7, 26), provide objective data during talent identification and development (20, 22, 23), evaluate the effectiveness of training (25) or nutritional interventions (10), and develop/evaluate training programs based on athletes' weaknesses (11, 17). The aim of a testing battery is to encompass a range of tests to ensure that all athlete attributes (e.g., speed, agility, fitness, strength) are evaluated. Data can then be compared against population-specific reference data, facilitating the interpretation of an athlete's scores. Despite this common approach, there appear to be some key limitations. Reference data are typically presented as the mean of a squad (6, 7), or at best, positional subgroups (e.g., in rugby, this may be forwards and backs; (21, 24, 26)). Furthermore, the presentation of mean data by positional group (i.e., forwards or backs) does not identify specific targets for an individual. For example, within rugby union, specific playing positions (i.e., prop, hooker, second row, flanker, number eight) within a positional group (i.e., forwards), all have differing roles during match play (5); thus it is assumed that their respective physiology would also differ. Presenting data as the mean for a group does not provide a target or benchmark for elite athletes, as often a larger degree of inter-athlete and even intra-positional variability will exist (18). The lack of position-specific data are likely due to challenges recruiting sufficient participant numbers from one positional group to establish an appropriate sample size.

There appears to be little consideration for athletes' or coaches' goals when strength and conditioning data are presented in the literature. Common within the discipline of sport and exercise psychology is performance profiling, which was first introduced by Butler (2-4). Performance profiling stems from personal construct psychology (13-15), which in the broadest sense, is concerned with understanding the ways individuals perceive and behave

within the world. Performance profiling encourages researchers and practitioners to regard an individual's perception or meaning of their performance as an essential source of information for identifying and understanding areas requiring improvement (8). Typically, this approach is lacking from the strength and conditioning literature, as even when individual case studies are presented, the goals of the athlete or indeed coach are often not considered (19). The development of performance profiling was to surpass traditional psychometric assessments that impose desirable constructs on an athlete (i.e., questionnaires), allowing participants to be more active in the decision-making or goal setting process (8). Participants would rank the importance of each attribute assessed. Attributes are then multiplied by the ranking score to provide greater context and relative importance within the interpretation (12). This concept would appear advantageous in the development of multidisciplinary support for athletes. Traditionally, a strength and conditioning coach would test an athlete and present data back to the coach or athlete to identify their respective strengths and weaknesses. Collaborative interpretation of the data, as used during performance profiling, would progress interdisciplinary to multidisciplinary practice.

The purpose of this article is to propose a collaborative method of interpreting fitness testing data, in relation to reference data with the addition of an athlete and coaches' ratings of the perceived importance of individual attributes.

METHODS

Approach to the Problem

Data from three academy rugby league players from an English Super League club were used to present this multidisciplinary approach when interpreting fitness testing data. The data were collected at the end of preseason in February 2015. Players were assessed on anthropometric (height, body mass, sum of four site skinfolds) and physical (10 and 20 m

sprint, 10 m momentum, vertical jump, Yo-Yo intermittent recovery test level 1 [IRT-1], one repetition max [1-RM] back squat, bench press and prone row) qualities. In collaboration with each player, three experienced rugby coaches then ranked the importance of the tests performed, considering playing position, stage of physical and technical / tactical development, and future goals. Data were then presented in various ways to highlight a novel way of athlete profiling, supporting the interpretation of the data.

Subjects

Three male professional academy rugby league players were included in the study. Player one was a positional prop (age 17 years; height 186.5 cm; body mass 93.2 kg). Player two was a positional hooker (age 18 years; height 173.2 cm; body mass 89.0 kg) and Player three was a positional winger (age 18 years; height 180.5 cm; body mass 84.1 kg). Players were involved in a professional rugby league academy, undertaking 3-4 gym-based sessions and 2-3 field-based sessions per week during preseason. All experimental procedures were approved by the institutional ethics committee with assent and parental consent provided along with permission from the rugby league club.

Procedures

The data were obtained from fitness testing sessions completed across two testing sessions, as previously described by Till et al., (26). The procedures were replicated to allow a comparison with the reference data previously published (26). All testing was undertaken by the lead researcher, and consisted of a standardised warm up including jogging, dynamic movements and stretches prior to testing, followed by full instruction and demonstrations of the assessments. The first testing session assessed speed (10 and 20 m sprint) and endurance (Yo-Yo IRT-1) capacity. Three days later, the second testing session was undertaken and

incorporated gym-based testing including anthropometric (height, body mass, sum of four site skinfolds), countermovement jump and 1-RM strength (back squat, bench press, prone row) measures.

Height was measured to the nearest 0.1 cm using a Seca Alpha stand (Seca, Birmingham, UK), and body mass, wearing only shorts, was measured to the nearest 0.1 kg using calibrated Seca Alpha (model 770) scales. Sum of four site skinfolds (biceps, triceps, subscapular, suprailliac) were measured using calibrated skinfold callipers (Harpenden, British Indicators, West Sussex, UK) in accordance to Hawes and Martin (9). A countermovement jump, with both hands positioned on the hips, was used to assess lower body power using a just jump mat (Probotics, Huntsville, AL, USA). Jump height was measured to the nearest 0.1 cm from the highest of three attempts recorded (26). Players were allowed 60 seconds rest between each countermovement jump. Sprint speed was assessed over 10 and 20 m using timing gates (Brower Timing Systems, IR Emit, Draper, UT, USA), as described by Till et al., (26). Players started 0.5 m behind the initial timing gate and were instructed to set off in their own time and run maximally past the 20 m timing gate. Times were recorded to the nearest 0.01 seconds with the quickest of the three times used for the sprint score. To calculate player momentum ($\text{kg}\cdot\text{sec}^{-1}$), 10 m sprint velocity (distance / sprint time) was multiplied by body mass (kg).

Endurance capacity was assessed via the Yo-Yo IRTL-1. Players completed 20 m shuttles, keeping to a series of beeps, followed by a 10-second rest interval. Running speed increased progressively throughout until the players reached volitional exhaustion or until players missed two beeps, resulting in the test being terminated. $\dot{V}O_2$ was then predicted ($\dot{V}O_2$ ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) = IRTL-1 distance (m) \times 0.0084 + 36.4; (1)) to allow comparisons with reference data (26), although it is accepted that a degree of error exists when predicting $\dot{V}O_2$ *per se* (16). Strength assessment included a 1-RM back squat, bench press, and prone row, exercises with which all players were familiar as they were regularly used as part of their

training program. Players performed a warm-up protocol of 8, 5, and 3 repetitions of individually-selected loads before 3 attempts of their 1-RM with 3-minute rest between attempts. For the squat, all players had to squat down until the top of the thigh was parallel with the ground, and then return to a standing position to record a 1RM score (26). For the bench press, athletes lowered the barbell to touch the chest and then pushed the barbell until elbows were locked out (26). For the prone row, the players lay face down on a bench which was determined so players' arms were locked out at the bottom position and then had to pull the barbell towards their chest, touching the bench.

In collaboration with each player, three rugby coaches, who worked with the players 3-4 times per week, ranked; body mass, explosive power (i.e., countermovement jump), acceleration (i.e., 10 m sprint), sprint ability (i.e., 20 m sprint), momentum, fitness (i.e., Yo-Yo test) and strength (i.e., squat, bench press and prone row) from the most (weighting of 7) to least (weighting of 1) important quality for each individual player. Prior to this, an experienced accredited (UKSCA) strength and conditioning coach explained what each test measured (e.g., the physiology), and how this likely translated to rugby *performance*. The rugby coaches did not have any specific education in strength and conditioning or athletic development, therefore the strength and conditioning coach provided examples of rugby specific key performance indicators (i.e., the tackle, successful collision, line bust, finishing a line break [by out running the opposition]) to align the tests to. The aim of this collaborative method was to ensure that the strength and conditioning coach, rugby coach and player were all working on the same goals to develop their rugby performance. The coaches and player reached a consensus for each of these qualities prior to their importance score being used. Coaches valued the input from each player, and understood the importance within the goal setting process. The ranking scores are shown in Table 1.

Statistical Analyses

Data are presented for individual players. Z-scores (statistical measurement of a score's relationship to the mean in relation to the standard deviation (*SD*)) are then presented with 0 representing the mean provided by Till et al. (26). A Z-score of 1 represents a player whose test scores were 1 *SD* above the mean (26), and a Z-score of -1 represents a player whose test scores were 1 *SD* below the mean (26). Modified Z-scores were then calculated to compare the players to benchmark data. The modified Z-score (i.e., 0) was determined as 2 *SDs* (i.e., 98th percentile) above the mean. Thus, when modified Z-scores are presented, a Z-Score of -1 represents a player whose test scores were 1 *SD* away from the benchmark data (mean plus 2 *SDs*; (26)). To incorporate the coach-athlete ranking the modified Z-score was simply multiplied by the coach-athlete ranking.

RESULTS

The following section presents the players data in various ways, to demonstrate the proposed method. Physical profiles of three players, their respective Z-score away from benchmark data (mean plus 2 *SDs* above data from Till et al. (26)), coach-athlete ranking score based on importance and total score (Z-Score away from benchmark data multiplied by coach-athlete ranking score) are shown in Table 1.

1 **Table 1. The physical profiles of three players, presented as Z-Scores away from benchmark data, coach-athlete ranking score based on importance**
 2 **and total score (Z-Score away from bench mark data multiplied by coach-athlete ranking score).**
 3

	Reference data (26)		Player 1 (Prop)				Player 2 (Hooker)				Player 3 (Winger)			
	Mean	Benchmark Reference Data	Test Score	Z-Scores away from Benchmark	Coach-Athlete Ranking	Total Score	Test Score	Z-Scores away from Benchmark	Coach-Athlete Ranking	Total Score	Test Score	Z-Scores away from Benchmark	Coach-Athlete Ranking	Total Score
Body Mass (kg)	88.8 ± 9.9	108.6	93	-1.6	7	-11.0	89	-2.0	2	-4.0	84	-2.5	1	-2.5
CMJ (cm)	52.5 ± 5.5	63.5	48.1	-2.8	5	-14.0	50.2	-2.4	3	-7.3	52.2	-2.1	4	-8.2
10 m Sprint (secs)	1.82 ± 0.07	1.68	1.86	-2.6	2	-5.1	1.79	-1.6	6	-9.4	1.76	-1.1	6	-6.9
20 m Sprint (secs)	3.11 ± 0.12	2.87	3.07	-1.7	1	-1.7	3.02	-1.3	1	-1.3	3.00	-1.1	7	-7.6
10m Momentum (kg·sec ⁻¹)	488 ± 47	582	500	-1.7	6	-10.5	497	-1.8	5	-9.0	477	-2.2	5	-11.2
Est. $\dot{V}O_2$ (ml·kg·min ⁻¹)	48.5 ± 2.9	54.3	48.0	-2.2	3	-6.5	53.0	-0.4	7	-3.2	52.0	-0.8	2	-1.6
Squat (kg)	138.4 ± 19.6	177.6	155	-1.2	4	-4.6	140	-1.9	4	-7.7	135	-2.2	3	-6.5
Bench (kg)	113.3 ± 16.4	146.1	135	-0.7	4	-2.7	105	-2.5	4	-10.0	105	-2.5	3	-7.5
Prone Row (kg)	97.6 ± 12.4	122.4	105	-1.4	4	-5.6	97.5	-2.0	4	-8.0	85	-3.0	3	-9.0


4 Benchmark reference data are based on the 98th percentile of data presented by Till et al., (26). CMJ = countermovement jump

5

6 For Player one, countermovement jump, 10 m speed and estimated $\dot{V}O_{2\max}$ were all
7 below mean reference data. Player two had lower countermovement jump and upper body
8 strength in comparison to mean reference data (26). Player three had lower body mass, 10 m
9 momentum and strength scores in comparison to mean reference data (26).

10 The coaches ranked body mass, 10 m momentum and countermovement jump as the
11 three most important attributes for Player one, while 20 m speed, 10 m speed and estimated
12 $\dot{V}O_{2\max}$ were ranked as the least important. For Player two, estimated $\dot{V}O_{2\max}$, 10 m speed and
13 10 m momentum were ranked as most important, whereas 20 m speed, body mass and
14 countermovement jump were deemed least important. For Player three, 20 m, 10 m speed and
15 10 m momentum were ranked as the most important, in contrast to body mass, estimated $\dot{V}O_{2\max}$
16 and strength, which were ranked as least important.

17 Data in comparison to their peers and mean data are shown in Figure 1A. Only data
18 with comparable units (i.e., kg) can be presented, therefore other variables are not depicted.
19 Figure 1B presents all the characteristics for players as Z-scores against mean reference data
20 (26).



21

22 **Figure 1. A) Presentation of body mass and strength data for three male academy rugby league players against mean reference data (26). B)**
 23 **Presentation of fitness characteristics for three male academy rugby league players against mean reference data (26). C) Presentation of fitness**
 24 **characteristics for three male academy rugby league players against benchmark mean reference data (26). D) Presentation of fitness characteristics**
 25 **for three academy male rugby league players against benchmark mean reference data (26) multiplied by coach-athlete ranking score.**

26 Figure 1C depicts player characteristics against benchmark (mean plus 2 *SDs*), as
27 opposed to mean reference data (26). When reviewing player characteristics, for example body
28 mass, player three is furthest away from benchmark data, whereas player one is closest. In
29 contrast, for 20 m sprint, Player one is furthest away from benchmark data, whereas player
30 three is closest.

31 Figure 1D presents player characteristics, incorporating the coach-athlete weighting.
32 The 0.0 is still the benchmark (i.e., mean plus 2 *SDs*; the 98th percentile), although the distance
33 away now considers how the coach-athlete ranked the importance of the characteristics for
34 each player. When reviewing the same example characteristics (i.e., body mass and 20 m
35 sprint), Figure 1D identifies different player weaknesses, for body mass, Player one is furthest
36 away from benchmark data, as opposed to Player three who was further away in Figure 1C.
37 Player three is now closest to the benchmark. This is due to the importance of body mass for
38 Player one and relatively lower importance for Player three. A similar trend is apparent for 20
39 m sprint, as Player three is now furthest away from benchmark data, as opposed to Player one
40 who was the furthest away in Figure 1C. This is again due to the importance of 20 m speed for
41 Player three and relatively lower importance for Player one.

42

43 **DISCUSSION**

44 This article considers the importance of coach-athlete ranked player characteristics
45 when presenting athlete profile data. The purpose of athlete profiling is to identify strengths
46 and weaknesses, to assist in the development of player characteristics for performance or
47 development. This report shows that when data are presented against benchmarks, considering
48 the importance of the characteristics when collaboratively ranked by the athlete and coaches
49 (Figure 1D), different athlete weaknesses can be highlighted, as opposed to the presentation of
50 data as absolute values (Figure 1A), against mean data (Figure 1B) or benchmark data on their

51 own (Figure 1C). Reference data presented in the literature is typically representative of various
52 positional playing groups to present data from a large sample. Despite the importance of a large
53 sample size, the application of this type of data are limited for elite athletes.

54 The limitation of presenting data in absolute units is that it is not possible to detail all
55 of an athlete's physical qualities (due to varying units of measurement such as kg, seconds, cm,
56 etc.). Thus, despite the presentation of an athlete's data against others' or mean data, it does
57 not present all their characteristics, provide a benchmark or target, and may not account for
58 individual goals, attributes or desired characteristics. The use of Z-scores using mean data
59 allows measures with varying units to be presented on the same figure (i.e., Figure 1B), and
60 thus the chance to view all physical attributes for the target athlete. Limitations of this
61 presentation method still exist: it remains a comparison to mean reference data, and does not
62 provide a robust benchmark as it is unlikely that athletes will strive to achieve mean levels of
63 characteristics associated with athletic performance. As such, modifying the Z-score to provide
64 athletes with a benchmark may be a more appropriate method of data presentation (Figure 1C),
65 although a limitation of this current method is that it does not account for individual goals,
66 attributes or desired characteristics. Given the large degree of within- and between-positional
67 differences that exist, the use of mean data, even when modified to provide a benchmark, is
68 questionable for elite athletes. As such, multiplying the data by a collaborative coach-athlete
69 rating (based on a hierarchy of most to least important) provides the athlete with individual,
70 identifiable and accessible traits to develop (Figure 1D). Athletes should, therefore, focus on
71 attributes that are identified as furthest away from their benchmark, and in turn, considered by
72 coaches to be important correlates of peak performance.

73

74

75 PRACTICAL APPLICATIONS

76 Strength and conditioning coaches should look to present the performance
77 characteristics of athletes against benchmark data, but also collaboratively with a coach and
78 athlete to consider the importance of each characteristic for individuals. In practice this process
79 allows athletes and coaches to specifically focus on individual weaknesses which are identified
80 as being important for an athlete's particular role in their sport or stage of development.

81 The examples presented from this Case Study show how data presentation may affect
82 the specific goals of players. For example, when observing the Z-score for Player one (prop),
83 countermovement jump and 10 m speed are the only tests that appear below average. Body
84 mass on the other hand appears above average. Given body mass development was the main
85 goal (achieved the highest score for the coach-athlete ranking) for this player, he may have not
86 been aware of this given he was already 'above average'. Therefore when the modified Z-score
87 was multiplied by the weighting factor, it is clear the athlete needs to focus on this attribute. A
88 similar example is presented for Player three (winger), who was below average for body mass,
89 and above average for 20 m speed. When the modified Z-score is multiplied by the coach-
90 athlete ranking weighting factor, 20 m speed is identifiable as an attribute to develop. Focusing
91 on specific attributes as opposed to multiple attributes would likely maximise the intended
92 adaptation by reducing any interference. Again this supports the ability of being able to
93 objectively identify goals from a large fitness testing data set, as shown in this Case Study
94 example.

95 When between athletes comparisons are made, 20 m speed from this Case Study is also
96 a good example. From the modified Z-score (Figure 1C), the order appears Player three, Player
97 two and then Player one (best to worse). Based on the coach-athlete ranking 20 m speed is low
98 on the priority for Player one, given his playing position. As such, when multiplying the
99 modified Z-score with the coach-athlete ranking, the order appears inverse, as this now

100 considers players goals (Player one, Player two and then Player three; best to worse, Figure
101 1D).

102 This article proposes a useful and novel method, based on Z-scores and coach-athlete
103 rating for practitioner to identify a target for individual athletes from a comprehensive testing
104 battery which may produce a large amount of information. The proposed method, with the use
105 of effective and simple spread sheet use can up a time effective way of processing data
106 objectively. Practitioners need to be aware that buy-in from the player and coach is needed for
107 this method to work effectively.

108

109

110 **REFERENCES**

- 111 1. Bangsbo J, Iaia FM, and Krstrup P. The Yo-Yo intermittent recovery test : a useful
112 tool for evaluation of physical performance in intermittent sports. **Sports Medicine.**
113 38: 37-51, 2008.
- 114 2. Butler R, Kremer J, and Crawford W. Psychological preparation of Olympic boxers.
115 **The Psychology of Sport: Theory and Practice:** 74-84, 1989.
- 116 3. Butler RJ. The performance profile: Theory and application. TSP 6, 2010.
- 117 4. Butler RJ, Smith M, and Irwin I. The performance profile in practice. **Journal of**
118 **Applied Sports Psychology.** 5: 48-63, 1993.
- 119 5. Cahill N, Lamb K, Worsfold P, Headey R, and Murray S. The movement characteristics
120 of English Premiership rugby union players. **Journal of Sports Science.** 31: 229-237,
121 2013.
- 122 6. Darrall-Jones JD, Jones B, and Till K. Anthropometric and Physical Profiles of English
123 Academy Rugby Union Players. **Journal of Strength and Conditioning Research.**
124 29: 2086-2096, 2015.
- 125 7. Emmonds S, Till K, Jones B, Mellis M, and Pears M. Anthropometric, Speed and
126 Endurance Characteristics of English Academy Soccer Players: Retrospective
127 comparisons with future career progression. **International Journal of Sports Science**
128 **and Coaching,** In press.
- 129 8. Gucciardi DF and Gordon S. Revisiting the performance profile technique: Theoretical
130 underpinnings and application. **The Sport Psychologist.** 23: 93-117, 2009.
- 131 9. Hawes MR and Martin AD. Human body composition. Kinanthropometry and exercise
132 physiology laboratory manual: tests, procedures and data 1: 7-46, 2001.

- 133 10. Ispoglou H, King R, Polman R, and Zanker C. Daily L-leucine supplementation in
134 novice trainees during a 12-week weight training program. **International Journal of**
135 **Sports Physiology and Performance**. 6: 38-50, 2011.
- 136 11. Jones B, Till K, Barlow M, Lees M, O'Hara J, and Hind K. Anthropometric and three-
137 compartment body composition differences between Super League and Championship
138 rugby league players: Considerations for the 2015 season and beyond. **PloS One**. 10,
139 2015.
- 140 12. Jones G. The role of performance profiling in cognitive behavioral interventions in
141 sport. **Sport Psychologist**. 7: 160-160, 1993.
- 142 13. Kelly G. The nature of personal constructs. *The psychology of personal constructs 1:*
143 105-183, 1955.
- 144 14. Kelly G. *The Psychology of Personal Constructs: Vol. 2; Clinical Diagnosis and*
145 *Psychotherapy*. WW Norton, 1955.
- 146 15. Kelly G. 1955. *The Psychology of Personal Constructs*. New York: Norton, 1965.
- 147 16. Krustup P, Mohr M, Nybo L, Jensen JM, Nielsen JJ, and Bangsbo J. The Yo-Yo IR2
148 test: Physiological response, reliability, and application to elite soccer. **Medicine and**
149 **Science in Sports and Exercise** 38: 1666-1673, 2006.
- 150 17. McGuigan MR, Cormack SJ, and Gill ND. Strength and power profiling of athletes:
151 selecting tests and how to use the information for program design. **Strength &**
152 **Conditioning Journal**. 35: 7-14, 2013.
- 153 18. Morehen JC, Routledge HE, Twist C, Morton JP, and Close GL. Position specific
154 differences in the anthropometric characteristics of elite European Super League rugby
155 players. **European Journal of Sports Science**. 20: 1-7, 2015.

- 156 19. Till K, Cobley S, O'Hara J, Chapman C, and Cooke C. An individualized longitudinal
157 approach to monitoring the dynamics of growth and fitness development in adolescent
158 athletes. **Journal of Strength and Conditioning Research.** 27: 1313-1321, 2013.
- 159 20. Till K, Cobley S, O'Hara J, Cooke C, and Chapman C. Considering maturation status
160 and relative age in the longitudinal evaluation of junior rugby league players.
161 **Scandinavian Journal of Medicine and Science in Sports.** 24: 569-576, 2014.
- 162 21. Till K, Cobley S, O'Hara J, Chapman C, and Cooke C. Anthropometric, physiological
163 and selection characteristics in high performance UK junior rugby league players.
164 **Talent Development and Excellence.** 2: 193-207, 2010.
- 165 22. Till K, Cobley S, O'Hara J, Morley D, Chapman C, and Cooke C. Retrospective
166 analysis of anthropometric and fitness characteristics associated with long-term career
167 progression in Rugby League. **Journal of Science and Medicine in Sport.** 18: 310-
168 314, 2015.
- 169 23. Till K and Jones B. Monitoring anthropometry and fitness using maturity groups within
170 youth rugby league. **Journal of Strength and Conditioning Research.** 29: 730-736,
171 2015.
- 172 24. Till K, Jones B, Darrall-Jones J, Emmonds S, and Cooke C. Longitudinal development
173 of anthropometric and physical characteristics with academy rugby league players.
174 **Journal of Strength and Conditioning Research.** 29; 1713-1722, 2015.
- 175 25. Till K, Jones B, Emmonds S, Tester E, Fahey J, and Cooke C. Seasonal changes in
176 anthropometric and physical characteristics within English academy rugby league
177 players. **Journal of Strength and Conditioning Research.** 28: 2689-2696, 2014.
- 178 26. Till K, Tester E, Jones B, Emmonds S, Fahey J, and Cooke C. Anthropometric and
179 physical characteristics of English academy rugby league players. **Journal of Strength
180 and Conditioning Research.** 28: 319-327, 2014.