
Citation:

Price, OJ and Ansley, L and Levai, IK and Molphy, J and Cullinan, P and Dickinson, JW and Hull, JH (2016) Eucapnic Voluntary Hyperpnea Testing in Asymptomatic Athletes. American journal of respiratory and critical care medicine, 193 (10). pp. 1178-1180. ISSN 1535-4970 DOI: <https://doi.org/10.1164/rccm.201510-1967LE>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/5091/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

EUCAPNIC VOLUNTARY HYPERPNEA TESTING IN ASYMPTOMATIC ATHLETES

Oliver J. Price¹ PhD, Les Ansley⁴ PhD, Irisz Levai⁵ MD, John Molphy⁶ PhD, Paul Cullinan² PhD,
John W. Dickinson^{*5} PhD, James H. Hull^{*2,3} MRCP, PhD

¹Carnegie School of Sport, Leeds Beckett University, Leeds, United Kingdom (UK).

²National Heart and Lung Institute, Imperial College London, London, UK.

³Department of Respiratory Medicine, Royal Brompton Hospital, London, UK.

⁴Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria
University, Newcastle upon Tyne, UK.

⁵Sports Therapy, Physical Activity and Health Research Group, School of Sport and Exercise
Sciences, University of Kent, Chatham Maritime, UK.

⁶Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool,
UK.

***Authors contributed equally**

Corresponding author:

Dr Oliver J Price, PhD

Carnegie School of Sport, Leeds Beckett University, Leeds,

LS6 3QT, UK

Tel: 0113 812 0001

Email: o.price@leedsbeckett.ac.uk

Word Count: 1115.

Keywords: Asthma, Athletes, Bronchoprovocation, Diagnosis, Exercise-induced bronchoconstriction.

To the Editor:

The prevalence of exercise-induced bronchoconstriction (EIB) is consistently reported to be greater in athletic individuals than in the general population (1). The reason for this difference remains to be fully determined, but may be explained by the development of airway hyper-responsiveness arising from repeated episodes of exercise hyperpnea when performed in noxious environments (2). Equally however, it is important that the prevalence of EIB is not over-estimated (i.e. false positive) by the application of overly sensitive diagnostic test methodologies. The aim of this study was to determine the normative response to a eucapnic voluntary hyperpnoea (EVH) challenge, in cohort of entirely asymptomatic athletes.

Traditionally EIB was diagnosed using an exercise test, accompanied by a spirometric assessment of expiratory airflow. A positive result was typically defined by a 10% pre-post challenge reduction in forced expiratory volume in one second (FEV₁), based largely on population studies evaluating the 'normative' response to exercise (3). Although this approach is logical, there are several limitations when employing this methodology in competitive athletes; these include the difficulties inherent to standardizing and controlling both the effective cardiorespiratory workload and environmental conditions (4). On this basis, several surrogate means for securing a diagnosis have been recommended (1), including both direct and indirect bronchoprovocation tests. Of these, EVH testing is cited as one of the best means to confirm or refute a diagnosis of EIB; principally due to its simplicity and the fact that it mimics the desiccating stimulus driving the development of EIB (4).

The diagnostic threshold for a positive EVH test was originally established from a cohort of asthmatic army recruits ($n = 90$) and 'normal' healthy controls ($n = 30$). A 10% fall in FEV₁ was recommended as the cut-off on the basis of optimising the relationship between specificity (90%) and sensitivity (63%) and approximates the threshold most commonly utilised with exercise testing (5, 6). The published data in athletes, is however limited and confounded by a selection bias with inclusion of individuals with a prior diagnosis of airways disease, history of respiratory symptoms and/or those prescribed asthma medication (3). Indeed, it is our experience that despite having normal baseline lung function and no respiratory symptoms, the majority of competitive athletes completing an EVH

challenge experience a fall in FEV₁ following EVH, frequently close to or beyond the 10% diagnostic cut-off. To describe this further we undertook a retrospective analysis of EVH tests performed in a large cohort of entirely asymptomatic athletes without a prior diagnosis of asthma or use of asthma medication. In accordance with previous methods (4) the EVH protocol consisted of breathing a dry compressed gas mixture (21% O₂, 5% CO₂, balance N₂) at a target ventilation equivalent to 85% maximum voluntary ventilation for 6 min. Spirometry was performed in triplicate at baseline and in duplicate at 3, 5, 7, 10, 15 min post EVH.

All athletes assessed ($n = 224$) were competitive at elite (i.e. either national or international standard) ($n = 161$) or recreational level (i.e. training/competing ≥ 6 hours/week) ($n = 63$) from a variety of sporting disciplines: athletics (i.e. competing in track and field events) ($n = 71$); rugby ($n = 61$); badminton ($n = 4$); boxing ($n = 28$); soccer ($n = 22$); hockey ($n = 13$); swimming ($n = 9$); rowing ($n = 8$); and biathlon ($n = 8$). All had normal predicted lung function values with no evidence of airway obstruction at rest (Table 1). The majority of athletes (98%) met accepted minimal target ventilation (i.e. minute ventilation $\geq 60\%$ MVV) (4). The mean (\pm SD) maximum fall in FEV₁ was calculated as $-7.6 \pm 6.7\%$ (Figure. 1) with the vast majority of athletes (98.2%) presenting with bronchoconstriction (i.e. reduction in FEV₁) at all time-points post EVH. The mean fall in FEV₁ was greater in elite ($-8.0 \pm 7.2\%$) than in recreational athletes ($-4.2 \pm 2.0\%$) ($P < 0.01$). In the very few athletes eliciting bronchodilation post challenge (1.8%, $n = 4$), the ‘improvement’ in FEV₁ was only minor (i.e. approximately 1-2% increase post EVH). When athletes who failed to achieve their target ventilation were excluded from the analysis ($n = 5$), the findings remained unchanged ($P > 0.05$) (data not shown). Likewise, when those with a severe fall in FEV₁ ($> 30\%$, $n = 4$) were excluded, the mean fall was not significantly altered ($-7.2 \pm 5.9\%$) ($P > 0.05$).

This study reports, for the first time, what may be considered the pattern of response to an EVH test in a cohort of athletes. Approximately 20% ($n = 44$) of this entirely asymptomatic athletic population would be deemed positive for a diagnosis of EIB based on the accepted 10% cut-off value. Although markers of airway inflammation or other pathological profiling for ‘asthma’ was not performed, the findings highlight that a fall in FEV₁ $> 10\%$ is encountered in a significant proportion of entirely

78 healthy asymptomatic athletes and thus in many cases may actually represent a variation of the
79 ‘normative’ airway response following exposure to the highly potent stimulus of EVH. Indeed, if an
80 abnormal response is based on a mean + 2 SD change, as has been previously used to define a cut-off
81 for EIB when employing an exercise test (8) our data suggests that a 15% cut-off would be a more
82 appropriate threshold. Of note, in all athletes with >15% fall in FEV₁, a sustained reduction in lung
83 function (i.e. minimum of two consecutive time-points) was observed, thus consistent with current
84 EIB American Thoracic Society committee guidelines (1).

85 The decision to initiate treatment for EIB should clearly be decided following the synthesis of clinical
86 findings and objective test results, however the selection of a ‘correct’ cut-off value for detection of a
87 condition is vital to guarantee diagnostic accuracy and ensure clinical care is optimised. We have
88 previously highlighted the poor clinical reproducibility of EVH in athletes when a 10% FEV₁ cut-off
89 threshold is employed (7). Whilst some may consider that a low cut-off threshold ensures that EIB is
90 ‘detected’ and the health and performance of an athlete is optimised, it is equally important to balance
91 this consideration with both the deleterious impact of unnecessary beta-2 agonist prescription and the
92 potential for distraction from other potentially important causes of exertional dyspnoea (9).

93 The findings from this study provide evidence that caution should be applied in the interpretation of a
94 mild post challenge reduction in lung function (i.e. 10-15% fall in FEV₁), certainly when applying
95 EVH to screen athletic squads. Further work is required to evaluate differences between athletes with
96 mild (e.g. 10-15% FEV₁) and more severe (e.g. >30% FEV₁) EIB and in comparison with
97 commensurate exercise challenge data. Employing inflammatory biomarker analysis and applying
98 supplementary test methodologies (e.g. impulse oscillometry) would provide additional value in this
99 setting.

100 *Oliver J. Price¹ PhD*

101 *Les Ansley⁴ PhD*

102 *Irisz Levai⁵ MD*

103 *John Molphy⁶ MSc*

104 *Paul Cullinan² PhD*

105 *John W. Dickinson⁵ PhD*

ACKNOWLEDGEMENTS

Nil.

FUNDING STATEMENT

Nil relevant.

COMPETING INTERESTS

The authors have no real or perceived conflict of interest in respect of this manuscript.

CONTRIBUTION STATEMENT

Conception and design: OP, JD, JH; Analysis and interpretation: OP, JD, JH; Drafting the manuscript for important intellectual content: OP, LA, IL, JM, PC, JD, JH.

GUARANTOR STATEMENT

OP confirms full responsibility for the content of the manuscript.

REFERENCES

1. Parsons JP, Hallstrand TS, Mastrorade JG, Kaminsky DA, Rundell KW, Hull JH, Storms WW, Weiler JM, Cheek FM, Wilson KC. An official American Thoracic Society clinical practice guideline: exercise-induced bronchoconstriction. *American journal of respiratory and critical care medicine* 2013; 187: 1016-1027.
2. Price OJ, Ansley L, Menzies-Gow A, Cullinan P, Hull JH. Airway dysfunction in elite athletes—an occupational lung disease? *Allergy* 2013; 68: 1343-1352.
3. Dryden DM, Spooner CH, Stickland MK, Vandermeer B, Tjosvold L, Bialy L, Wong K, Rowe BH. Exercise-induced bronchoconstriction and asthma. *Evidence Report/Technology Assessment* 2010; 189.
4. Anderson S, Argyros G, Magnussen H, Holzer K. Provocation by eucapnic voluntary hyperpnea to identify exercise induced bronchoconstriction. *British journal of sports medicine* 2001; 35: 344-347.
5. Hurwitz KM, Argyros GJ, Roach JM, Eliasson AH, Phillips YY. Interpretation of eucapnic voluntary hyperventilation in the diagnosis of asthma. *CHEST Journal* 1995; 108: 1240-1245.
6. Argyros GJ, Roach JM, Hurwitz KM, Eliasson AH, Phillips YY. Eucapnic voluntary hyperventilation as a bronchoprovocation technique. *Chest* 1996; 109: 1520-1524.
7. Price OJ, Ansley L, Hull JH. Diagnosing Exercise-Induced Bronchoconstriction With Eucapnic Voluntary Hyperpnea: Is One Test Enough? *The Journal of Allergy and Clinical Immunology: In Practice* 2014.
8. Helenius IJ, Tikkanen HO, Haahtela T. Occurrence of exercise induced bronchospasm in elite runners: dependence on atopy and exposure to cold air and pollen. *British journal of sports medicine* 1998; 32: 125.
9. Walsted NE, Hull JH, Backer V. High Prevalence of Exercise-Induced Laryngeal Obstruction in Athletes. *Medicine and science in sports and exercise* 2013; 45: 2030-2035.

Table 1. Clinical characteristics and baseline lung function.

Variables	
Sex (M:F)	178 : 46
Age (years)	23 \pm 4
Height (cm)	179.6 \pm 10.2
Weight (kg)	83.7 \pm 17.5
BMI (kg•m ⁻²)	22.0 \pm 4.0
FEV ₁ (L)	4.52 \pm 0.78
FEV ₁ (% predicted)	101.9 \pm 11.2
FVC (L)	5.36 \pm 1.03
FVC (% predicted)	102.2 \pm 12.5
FEV ₁ /FVC (%)	85.1 \pm 7.6
Target ventilation (L/min)	135.7 \pm 23.4
Achieved ventilation (L/min)	121.0 \pm 25.2
Predicted ventilation (%)	89.6 \pm 14.4
Total fall in FEV ₁ (%)	-7.6 \pm 6.7%
-Elite athlete; fall in FEV ₁	-8.0 \pm 7.2%
-Recreational athlete; fall in FEV ₁	-4.2 \pm 2.0%

Data presented as mean \pm SD. *n* = 224.




Figure 1. Frequency distribution of the maximum reduction in FEV₁ in asymptomatic athletes post EVH. Broken horizontal line (*black*) represents current diagnostic threshold (i.e. $\geq 10\%$ fall in FEV₁) and broken horizontal line (*red*) represents proposed revised diagnostic threshold (i.e. $\geq 15\%$ fall in FEV₁). Data presented as Mean \pm SD.