
Citation:

Weaving, D and Sawczuk, T and Williams, S and Scott, T and Till, K and Beggs, CB and Johnston, RD and Jones, B (2018) The peak duration-specific locomotor demands and concurrent collision frequencies of European Super League rugby. *Journal of Sports Sciences*, 37 (3). pp. 322-330. ISSN 0264-0414 DOI: <https://doi.org/10.1080/02640414.2018.1500425>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/5226/>

Document Version:

Article (Accepted Version)

This is an Accepted Manuscript of an article published by Taylor & Francis in *Journal of Sports Sciences* on 19 July 2018, available online: <http://www.tandfonline.com/10.1080/02640414.2018.1500425>

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

1 **The peak duration-specific locomotor demands and concurrent collision**
2 **frequencies of European Super League rugby.**

3 **Running Title:** Peak locomotor and collision demands of the European Super League

4 **Word Count:** 4330

5 Dan Weaving^{1,2}, Thomas Sawczuk^{1,3}, Sean Williams⁴, Tannath Scott^{5,6}, Kevin
6 Till^{1,2,7}, Clive Beggs¹, Rich D Johnston⁸, Ben Jones^{1,2,7,9}

7 ¹Institute for Sport, Physical Activity and Leisure, Leeds Beckett University,
8 Leeds, West Yorkshire, United Kingdom.

9 ²Leeds Rhinos Rugby League Club, Leeds, West Yorkshire, United Kingdom

10 ³Queen Ethelburgas Collegiate, York, United Kingdom

11 ⁴University of Bath, Bath, United Kingdom

12 ⁵Brisbane Broncos Rugby League club, Brisbane, Australia

13 ⁶School of Human Movement and Nutrition Sciences, University of
14 Queensland, Brisbane, Australia

15 ⁷Yorkshire Carnegie Rugby Union club, Leeds, United Kingdom

16 ⁸School of Exercise Science, Australian Catholic University, Brisbane,
17 Australia

18 ⁹The Rugby Football League, Leeds, United Kingdom

19

20 **Abstract**

21 Understanding the most demanding passages of European Super League
22 competition can optimise training prescription. We established positional and match
23 half differences in peak relative distances ($\text{m}\cdot\text{min}^{-1}$) across durations, and the number
24 of collisions, high-speed- and very-high-speed-distance completed in the peak 10 min
25 period. Moving-averages (10 s, 30 s, 1 min, 5 min, 10 min) of instantaneous speed
26 ($\text{m}\cdot\text{s}^{-1}$) were calculated from 25 professional rugby league players during 25 matches
27 via microtechnology. Maximal $\text{m}\cdot\text{min}^{-1}$ was taken for each duration for each half.
28 Concurrently, collisions (n), high-speed- (5 to 7 $\text{m}\cdot\text{s}^{-1}$; m) and very-high-speed-
29 distance (> 7 $\text{m}\cdot\text{s}^{-1}$; m) were coded during each peak 10 min. Mixed-effects models
30 determined differences between positions and halves. Aside from peak 10 s, *trivial*
31 differences were observed in peak $\text{m}\cdot\text{min}^{-1}$ between positions or halves across
32 durations. During peak 10 min periods, adjustables, full- and outside-backs ran more
33 at high-speed and very-high-speed whilst middle- and edge-forwards completed more
34 collisions. Peak $\text{m}\cdot\text{min}^{-1}$ is similar between positional groups across a range of
35 durations and are maintained between halves of the match. Practitioners should
36 consider that whilst the overall peak locomotor ‘intensity’ is similar, how they achieve
37 this differs between positions with forwards also exposed to additional collision bouts.

38

39 **Introduction**

40 Rugby league is played professionally in the European Super League (ESL) and in
41 Australasia within the National Rugby League (NRL). It is a team-sport characterised
42 by prolonged intermittent bouts of locomotor and collision activity (Waldron, Twist,
43 Highton, Worsfold & Daniels, 2011; Gabbett, Jenkins & Abernethy, 2012; Johnston,
44 Gabbett & Jenkins, 2014; Twist et al., 2014). Practitioners prescribe numerous
45 training modes to develop the wide range of physical qualities (e.g. muscular strength,
46 speed) that are needed to succeed in competition (Gabbett et al., 2012; Till,
47 Scantlebury & Jones, 2017; Weaving et al., 2017). However, in order to improve the
48 likelihood of positive outcomes it is important to manage the accumulation and
49 distribution of the external and internal loads prescribed to players (Impellizzeri,
50 Rampinini & Marcora, 2005; Soligard et al., 2016; Vanrenterghem, Nedergaard,
51 Robinson & Drust, 2017). By understanding the most intense periods of competition,
52 practitioners can improve their prescription of the external load (i.e. running,
53 accelerating, collisions) across training modes and ensure players are appropriately
54 exposed to these demands in training (Impellizzeri et al., 2005; Weaving et al., 2017).

55 Microtechnology units incorporating global positioning systems (GPS) chips and other
56 inertial measurement devices are now widely used to quantify both the locomotor
57 (Johnston et al., 2014) and collision demands (Gabbett et al., 2012; Hulin, Gabbett,
58 Johnston & Jenkins, 2017) of professional rugby league competition. Across a whole
59 match, players typically cover between 5000 and 8000m (Waldron et al., 2011; Twist
60 et al., 2014; Johnston et al., 2014) and are subjected to 30-65 collision events (Hulin
61 et al., 2017) dependent on position (Gabbett et al., 2012). Whilst whole-game data
62 are useful to understand the accumulation of load and how it varies by position,
63 quantifying the rate in which this activity accumulates (i.e. 'intensity') is important for
64 understanding the specificity of training.

65 Relative distance ($\text{m}\cdot\text{min}^{-1}$) is a frequently reported measure used to quantify the
66 overall rate of locomotor activity during competition (Waldron et al., 2011; Twist et al.,
67 2014; Johnston et al., 2014). In a systematic review, Johnston et al. (2014) reported
68 23 positional relative distances from 9 manuscripts across the NRL ($n = 7$) and ESL
69 ($n = 2$) competitions. The mean data across these studies suggests the whole-game
70 relative distance to be $\sim 94.7 \pm 6.1 \text{ m}\cdot\text{min}^{-1}$. However, the utility of this information as
71 a basis to prepare players is questionable because it under-represents periods in the
72 game where players complete greater relative distances for prolonged periods of time
73 (i.e. $> 5 \text{ min}$) (Delaney et al., 2015). Technical-tactical training is a commonly
74 prescribed modality in professional rugby league training programmes (Gabbett et al.,
75 2011; Lovell et al., 2013; Weaving et al., 2017). Therefore, identifying the maximal
76 relative distances across a range of time periods should provide useful information
77 for technical-tactical coaches to evaluate their training prescription (Robertson &
78 Joyce, 2015).

79 Delaney et al. (2015) used a moving-average of the instantaneous sampled speed
80 ($5\text{Hz m}\cdot\text{s}^{-1}$) during NRL competition. Using this approach, the authors were able to
81 determine the between-position differences in peak relative distances completed
82 across 1 to 10 min moving average periods. Logically, as the duration of activity
83 decreased, the peak relative distance for a given duration increased (Delaney et al.
84 2015). Interestingly, however, substantial differences in total distance between player
85 positions have been observed using whole-game data (Waldron et al., 2011; Twist et
86 al., 2014; Johnston et al., 2014). Delaney et al. (2015) reported that full-backs
87 completed substantially greater peak relative distances across the range of durations
88 compared with players in other positions (i.e. halves, outside backs, edge-forwards
89 and hit-up-forwards), who covered similar peak relative distances. For example, the
90 mean maximal 10 min relative distance reported across a NRL season for full-backs
91 was $105 \pm 10 \text{ m}\cdot\text{min}^{-1}$, with halves ($93 \pm 10 \text{ m}\cdot\text{min}^{-1}$), middle forwards ($90 \pm 10 \text{ m}\cdot\text{min}^{-1}$)

92 ¹), edge forwards ($95 \pm 7 \text{ m}\cdot\text{min}^{-1}$) and outside backs ($97 \pm 14 \text{ m}\cdot\text{min}^{-1}$) covering
93 substantially reduced relative distances. Due to the previously reported differences in
94 whole-game relative distances (including high-speed) between the two competitions
95 (Twist et al., 2014), this would seem important to establish in the ESL.

96 Given the interplay that occurs between locomotor and collision activity in rugby
97 league, one limitation of the above study (Delaney et al., 2015) is that the collision
98 activities completed by players during periods of peak locomotor intensity were not
99 reported. Hit-up-forwards have less playing time (Johnston et al., 2014), despite
100 Delaney et al. (2015) demonstrating little practical difference in the peak running
101 demands for this position compared to positions which complete the full match.
102 Increased collision activity (Gabbett et al., 2012) and body mass (Darrall-Jones et al.,
103 2015; Jones et al., 2015) compared to other positions are possible mechanisms for
104 this reduced involvement. However, whilst the frequency of collision activity of whole-
105 match NRL competition has previously been detailed (Gabbett, Jenkins & Abernethy,
106 2011; Gabbett et al., 2012; Cummins & Orr, 2015), concurrent data relating to collision
107 activity embedded within the peak locomotor (i.e. relative distances) distances
108 covered during ESL competition is currently unavailable. However, provision of such
109 data would provide practitioners with extremely useful information with which to
110 generate a holistic understanding of the most demanding passages of play for the
111 positional groups. These data could then be used as collective markers of “intensity”
112 to assist practitioners to plan the incremental progression of both collision and
113 locomotor activity during physical preparation (i.e. pre-season) and return-to-play
114 protocols.

115 Based on the information above, we designed the current study with the specific aim
116 of: 1) establishing the positional differences in duration-specific peak relative
117 distances covered during ESL competition; 2) establishing the positional differences
118 in high-speed-distance ($5 \text{ to } 7 \text{ m}\cdot\text{s}^{-1}$), very-high-speed-distances ($> 7 \text{ m}\cdot\text{s}^{-1}$) and the

119 number of concurrent collisions within the peak 10 min relative distances of ESL
120 rugby; and 3) establishing the within-position differences in these demands between
121 halves of the match.

122 **Method**

123 **Participants**

124 Data were collected from 25 male professional rugby league players (age = $27.3 \pm$
125 4.8 yrs, body mass = 96.0 ± 12.6 kg and height = 184.5 ± 6.8 cm) from the same ESL
126 club during 25 matches during the 2017 ESL regular season (18 wins, 7 losses; mean
127 \pm SD score margin: 4 ± 21 points). Players were coded for position at the start of each
128 match, with the number of match observations and individual player appearances for
129 each position including: fullbacks (5 players; $n = 25$), outside backs (centres and
130 wings; 9 players; $n = 96$), adjustables (half-back, five-eighth; hooker; 6 players; $n =$
131 72), middle-forwards (middle- and loose-forward; 10 players; $n = 92$) and edge-
132 forward (6 players; $n = 48$). The mean \pm SD number of matches per player was $16 \pm$
133 6 . When a player changed position within a half their data was omitted from the
134 dataset ($n = 7$). Players provided informed consent and ethics approval was gained
135 from the institutions review board.

136 Microtechnology (Optimeye S5, Catapult Innovations, Melbourne, Victoria) was
137 positioned in a customised padded pouch sewn into the players shirt which was
138 positioned in the centre of the upper back. To reduce the influence of inter-unit error,
139 each player was provided with the same device for the period of data collection. The
140 test-retest reliability of Catapult 10Hz devices to measure instantaneous speed
141 across a range of starting velocities has been reported to be acceptable (coefficient
142 of variation: 2.0 to 5.3%) (Varley et al., 2012). The number of satellites and horizontal
143 dilution of precision (HDOP) during data collection were (mean \pm SD) 15 ± 2 and 0.8

144 ± 0.6 , respectively. Greater than 6 connected satellites and HDOP values less than 1
145 are considered ideal for GPS data collection (Malone, Lovell, Varley & Coutts, 2016).

146 *Duration-specific peak relative distance ($m \cdot min^{-1}$)*

147 During matches, each players period of involvement in the game was coded in real-
148 time using proprietary software (Catapult Openfield v1.14; firmware: 7.27) (Weaving,
149 Whitehead, Till & Jones, 2017; Barrett, 2017). A Greenwich mean time (GMT) 'time-
150 stamp' was created to determine the 'start' and 'end' time of each players involvement
151 in each half. This was also completed for interchange players to ensure that only
152 match time were included in the analysis and to ensure appropriate coding of their
153 involvement. For inclusion in any match half, a players involvement had to be greater
154 than 20 min. This criteria was applied so that even if a player had two involvements
155 in a single half, only one data entry per half per player could be included in the final
156 analysis (Delaney et al., 2015). All natural match breaks (e.g. injury, try
157 scored/conceded) were included in the analysis.

158 To establish the duration-specific running intensities ($m \cdot min^{-1}$), a players
159 instantaneous speed ($m \cdot s^{-1}$), derived from the Doppler Shift method, was recorded
160 every 0.1s (i.e. 10Hz). A time-series file, detailing a record of instantaneous speed
161 every 0.1s was then exported from the proprietary software (Catapult Openfield
162 v1.14). Therefore, the first speed sample represents the 'start' of their match
163 involvement (i.e., half or interchange period), whilst the final speed sample represents
164 the 'end' of their involvement.

165 A custom-built algorithm using the zoo package (Zeileis & Grothendieck, 2005) in R
166 (v R-3.1.3, R Foundation for Statistical Computing, Vienna, Austria) was developed
167 to compute a moving-average of each player's instantaneous speed across different
168 durations. Moving-averages were calculated across five different durations (10 s, 30
169 s, 1 min, 5 min, 10 min) for each half. Like previous studies (Delaney et al., 2015),

170 these durations were arbitrarily chosen to represent shorter and prolonged durations
171 of activity due to their use in training prescription. For example, for a 10 min moving-
172 average, the algorithm computed a moving-average for every 6000 instantaneous
173 speed samples (i.e. 10 samples per second for 600 seconds [10 min]). This process
174 was repeated for each of the respective 'durations' in the study. For each player and
175 half, the respective computed moving-average values for each duration were then
176 concatenated into a data frame (with the columns representing the different moving
177 average durations [i.e. 10 s to 10 min] and the rows representing the moving average
178 instantaneous speed value). This was then exported to Microsoft Excel to determine
179 the maximum moving-average for each duration. This was multiplied by the moving-
180 average duration to determine a players maximal moving-average of relative distance
181 ($\text{m}\cdot\text{min}^{-1}$).

182 *Concurrent collision-, high-speed- and very-high-speed-distance within peak 10 min*
183 *relative distances*

184 The number of collisions, high-speed-distance (5 to $7 \text{ m}\cdot\text{s}^{-1}$) and very-high-speed-
185 distance ($>7 \text{ m}\cdot\text{s}^{-1}$) were selected to provide additional information of the concurrent
186 locomotor and collisions with the peak 10 min relative distances ($\text{m}\cdot\text{min}^{-1}$) identified
187 during ESL competition (Twist et al., 2014; McLellan, Lovell & Gass, 2011). The
188 minimum effort duration for high-speed and very-high-speed distance was set at 1
189 second (Varley, Jaspers, Helsen & Malone, 2017; Malone et al., 2017).

190 PlayerLoad™ was quantified as per previous methods which has demonstrated
191 acceptable reliability (Boyd, Ball & Aughey, 2011). The number of collisions were
192 quantified using the 'tackle' algorithm provided by the manufacturer which is derived
193 from the 100Hz tri-axial accelerometer and gyroscope also housed within the
194 microtechnology device as per previous methods. This has been reported to possess
195 acceptable validity to detect collision events, with specificity and sensitivity of $91.7 \pm$

2.5% and $93.9 \pm 2.4\%$ respectively, when short duration (< 1 second) and low-intensity (i.e. < 1 AU of PlayerLoad™) events were excluded (Hulin et al. 2017).

To export the number of collisions, high-speed- and very-high-speed-distance completed by each player, the GMT associated with the identified peak $10 \text{ m}\cdot\text{min}^{-1}$ moving-average for each half match file were coded within the proprietary software (Openfield v1.14, Catapult Innovations, Scoresby, Victoria, Australia) and exported into a customised spreadsheet.

203 **Statistical Analysis**

Linear mixed-effects models were used to estimate the differences between the positional groups and match half. For the continuous variables of 10 s, 30 s, 1-min, 5 min and 10 min peak $\text{m}\cdot\text{min}^{-1}$, 10 min high-speed-distance and very-high-speed-distance, estimations were made via PROC MIXED in SAS University Edition (SAS Institute, Cary, NC). For collision data, a generalised linear mixed-effects model was used, assuming a negative binomial distribution, via the *lme4* package (Bates, Maechler, Bolker & Walker 2015) in R (version 3.3.1). In both models the (fixed) effects of playing position and match-half were estimated. The interaction between these fixed effects was also explored, by including a multiplicative term in the models. The random effects in both models were match identity (differences between average match demands not accounted for by the fixed effects), athlete identity (differences between athletes' mean match demands) and the residual (within-athlete match-to-match variability). Magnitude-based inferences were used to provide an interpretation of the real-world relevance of the outcomes. For all peak relative distance durations, a difference of $10 \text{ m}\cdot\text{min}^{-1}$ was set as the smallest worthwhile effect threshold. This was chosen based on previous research (Delaney et al., 2015) as practitioners are unlikely to utilise between-position training prescription that is more specific than a 10 metre difference. For collisions, high-speed- and very-high-speed-distance comparisons, a value equivalent to a difference in means of 0.20 was set as the

223 smallest worthwhile effect threshold. For all comparisons, effects were classified as
224 *unclear* if the percentage likelihood that the true effect crossed both positive and
225 negative smallest worthwhile effect thresholds were both greater than 5%. Otherwise,
226 the effect was deemed clear, and was qualified with a probabilistic term using the
227 following scale: <0.5%, *most unlikely*; 0.5-5%, *very unlikely*; 5-25%, *unlikely*; 25-75%,
228 *possible*; 75-95%, *likely*; 95-99.5%, *very likely*; >99.5%, *almost certainly* (Hopkins,
229 2009).

230 **Results**

231 *Duration-specific peak relative distance ($m \cdot min^{-1}$)*

232 Table 1 details the mean \pm SD for peak relative distances from 10 s to 10 min by 1st
233 and 2nd half. Between halves of the match (1st vs 2nd) there were *likely* to *most likely*
234 trivial differences in these variables for all within-position comparisons.

235 Table 2 details the raw least square means positional differences and magnitude
236 based inferences for these variables. Although, full backs, outside backs and
237 adjustables covered substantially greater relative distances across 10 s periods, there
238 were *possibly* to *almost certainly* trivial differences between all positional groups in
239 peak 1, 5 and 10 min relative distances.

240 *Concurrent collisions, high-speed- and very-high-speed-distances within peak 10 min* 241 *relative distances ($m \cdot min^{-1}$)*

242 Table 1 displays the mean \pm SD for peak 10 min relative distance and concurrent
243 number of collisions, high-speed- and very-high-speed-distance for each positional
244 group.

245 Figure 1, 2 and 3 displays the standardised mean difference plus 90% confidence
246 intervals for positional differences in the concurrent number of collisions, high-speed-
247 and very-high-speed-distance completed during the peak 10 min relative distances.

248 Whilst there were *unclear* differences in the number of collisions between full backs,
249 adjustables and outside backs, edge and middle forwards completed a substantially
250 greater number of collisions compared to these three positional groups.

251 Between 1st and 2nd halves there were *possibly* reductions in high-speed distance for
252 full-backs (ES: 0.25 [-0.20 to 0.69]), *very likely* reductions for outside backs (ES: 0.54
253 [0.32 to 0.76]), *possibly* trivial reductions for adjustables (ES: 0.13 [-0.12 to 0.38]),
254 *possibly* reductions for middle forwards (ES: 0.29 [0.06 to 0.51]) and *likely* reductions
255 for edge forwards (ES: 0.40 [0.09 to 0.71]). For very-high-speed-distance, there were
256 *likely* reductions between 1st and 2nd halves for full backs (ES: 0.44 [-0.03 to 0.92])
257 and outside backs (ES: 0.40 [0.16 to 0.64]) and *possibly* trivial differences for
258 adjustables (ES: -0.10 [-0.36 to 0.17]) and middle forwards (ES: 0.12 [-0.13 to 0.36]).
259 *Unclear* differences were observed for wide-forwards (ES: -0.04 [-0.37 to 0.29]). For
260 collisions, there were *likely* trivial differences between 1st and 2nd halves for all
261 positional groups.

262 **Discussion**

263 The primary aim of the study was to establish the positional differences in peak
264 duration-specific relative distances and the number of collisions, high-speed-, and
265 very-high-speed-distances completed within the peak 10 min locomotor period of ESL
266 competition. A secondary aim was to determine whether these peak demands differed
267 between the 1st half and 2nd half of competition within positional groups.

268 The main findings were that whilst adjustables, outside- and full-backs completed
269 greater peak running 'intensities' during 10 s locomotor bouts, *likely to almost certainly*
270 trivial differences were observed between all the positional groups as the duration
271 increased (30 s to 10 min). Although, during the peak 10 min locomotor period,
272 adjustables outside- and full-backs covered greater high-speed and very-high-speed-
273 distances than middle- and edge-forwards, the latter positional groups completing a

274 substantially greater number of collisions. The difference in demands between 1st and
275 2nd halves were *likely* to *almost certainly* trivial across the majority of variables,
276 although there were small decreases in high-speed- and very-high-speed-distance
277 across all positional groups during the peak 10 min locomotor period of the 2nd half.
278 Collectively this suggests for prolonged periods of an ESL match (i.e. 2 x 10 min
279 periods), the positions demonstrate limited practical differences in overall relative
280 distance, although middle- and edge-forwards complete a greater number of
281 collisions, whereas fullbacks, outside backs and adjustables complete greater
282 distances at high-speed during this time. This study is the first to provide data of the
283 peak locomotor and concurrent collision activity of ESL rugby by halves of the match.
284 The findings suggest that it is important for coaches to prescribe periods of training
285 that provide positional groups with similar exposures to relative distance, while still
286 ensuring that the respective positions achieve this in a different manner (i.e. backs
287 more high-speed running) and that they are concurrently exposed to varying collision
288 activity (i.e. forwards more collisions).

289 Compared to previous literature (Delaney et al., 2015), the peak duration-specific
290 relative distances of ESL competition appear comparable to those reported within the
291 NRL. This suggests that the peak locomotor demands of the two competitions are
292 consistent. Consequently, there appears to be a growing body of evidence to suggest
293 that the peak duration-specific relative distances of professional rugby league
294 competition are consistent across teams and competitions and therefore, when
295 controlling for contextual influences, there appears to be a 'ceiling' requirement of
296 relative distance that professional rugby league players are required to complete.
297 Importantly, it must be considered that the data in the current study represents the
298 average of the maximal relative distances covered by players per half, per game.
299 Therefore, detailing the range of peak demands experienced by players, including the
300 maximal recorded exposure during competition can also provide useful information of

301 the highest recorded demands (Table 1). For example, whilst whole-game relative
302 distances are $\sim 94.7 \text{ m}\cdot\text{min}^{-1}$ (Johnston et al., 2014), at least 10% of the match (i.e. 2
303 x 5 min) is spent covering relative distances between 107 and $116 \text{ m}\cdot\text{min}^{-1}$. Depending
304 on position, this rose to between 134 to $165 \text{ m}\cdot\text{min}^{-1}$ during some matches (Table 1).
305 Practitioners should therefore aim to ensure players receive an appropriate exposure
306 to technical-tactical activities at these maximal competition 'intensities'.

307 Due to the importance of 'winning' the collision contest and its interplay with locomotor
308 activity, a novel aspect of the current investigation was the detail and positional
309 comparison of the frequency of collision bouts during the peak 10 min locomotor
310 periods of ESL competition. These appear similar to the whole-game collision
311 frequencies ($\text{number}\cdot\text{min}^{-1}$) reported in the NRL (Gabbett et al., 2012) which revealed
312 middle-forwards (mean [range]: $1.09 [0.96 \text{ to } 1.22]$) to exhibit the greatest frequency
313 of collisions, with differences also observed between wide-running-forwards (0.76
314 $[0.69 \text{ to } 0.84]$), adjustables ($0.58 [0.45 \text{ to } 0.71]$) and outside backs ($0.38 [0.32 \text{ to } 0.43]$). This suggests that during the peak locomotor passages of ESL competition,
316 the frequency of collision activity is maintained at whole-game 'intensities'. Therefore,
317 practitioners should consider the amalgamation of collision activity whilst aiming to
318 replicate the 'peak' relative distances reported in the current study. However, it is
319 important to note that the current study quantified the collisions embedded within the
320 peak locomotor demands and it is plausible that the peak frequency of collisions for
321 a given duration could be substantially greater than those reported. Future research
322 should therefore seek to establish the peak collision frequencies experienced by the
323 positional groups for a range of durations to further strengthen the understanding
324 between locomotor and collision activity during professional rugby league
325 competition.

326 In professional rugby league it is commonplace for forwards (particularly middle-
327 forwards) to complete reduced time on the pitch during matches (Waldron et al., 2011;

Twist et al., 2014; Johnston et al., 2014). This has previously been attributed to forwards possessing reduced prolonged intermittent running capacity (Scott et al., 2017), greater body mass (Jones et al., 2015; Darrall-Jones et al., 2016) and greater collision activity compared to backs (Gabbett et al., 2012). Our study suggests that this is likely because middle-forwards complete similar peak locomotor intensities to backs whilst concurrently completing substantially more collisions for prolonged periods of the match (i.e. 2 x 10 min). When locomotor bouts are controlled, the addition of collisions have been reported to increase a players rating of perceived exertion, blood lactate concentration and heart rate (Johnston & Gabbett, 2011; Mullen, Highton & Twist, 2015; Norris, Highton, Hughes & Twist, 2016), suggesting that the internal physiological cost of competition would be greater in the forwards position. In addition, the total number of contacts in the forwards position has previously been reported to relate to decrements in perceptual muscle soreness ($r = 0.62$), perceptual fatigue ($r = 0.69$) and countermovement jump flight time ($r = -0.55$) 24 hours post ESL competition (Twist, Waldron, Highton, Burt & Daniels, 2012). Despite this, such substantial relationships appear to be absent in the backs (soreness: $r = 0.20$; fatigue: $r = 0.11$; jump flight time: $r = -0.25$) (Twist et al., 2012). Collectively, this suggests that rugby league forwards are subjected to greater psycho-physiological and biomechanical loads (Soligard et al., 2016; Vanrenterghem et al., 2017) per min of competition than backs, leading to similar amounts of “fatigue” in the days following competition, despite forwards competing for a reduced amount of time (Twist et al., 2012; Johnston et al., 2014). Therefore, practitioners should ensure that training prescription and recovery periodisation reflect this, particularly when forwards complete substantially greater playing times than typically accustomed to.

Whilst this study is the first to detail the interplay between locomotor and collision activity during the peak passages of competition and how they differ between position

355 and halves of the match, the study is not without limitations. Firstly, the data were
356 collected from a single ESL club, which may not be representative of the differences
357 observed with other teams in the competition. Secondly, the collision, high-speed-
358 and very-high-speed-distance demands embedded within the peak 10 min duration
359 were extracted on the assumption that the measurement of instantaneous speed
360 ($\text{m}\cdot\text{s}^{-1}$) provides a valid representation of the peak locomotor demands of professional
361 rugby league competition (Delaney et al., 2015). Acceleration and deceleration events
362 are prevalent in professional rugby league, due to the spatial constraints imposed by
363 the 10-metre rule separating the opposing structures of the attacking and defending
364 teams. Therefore, determining the collisions, high-speed- and very-high-speed
365 distances completed within the peak acceleration demands could arguably provide a
366 more valid representation of the peak locomotor demands of competition.

367 Despite this, for the practitioner wishing to optimise training prescription, it is important
368 to find the balance between the validity of the measurement and practical/actionable
369 data. In particular, during the planning and prescription of training a fundamental
370 strategy adopted by practitioners is to control and manipulate the overall distance
371 covered per unit of time (i.e. relative distance). This warrants consideration, as
372 technical-tactical training is the most frequently prescribed modality in professional
373 rugby league, particularly during the in-season period which lasts the majority of the
374 calendar year (Gabbett et al., 2012; Lovell et al., 2013; Weaving et al., 2017).
375 Therefore, it would be preferable to appropriately expose players to these peak
376 demands (e.g. 10 min continuous bouts) within this mode of training to concurrently
377 satisfy both the physical and technical-tactical requirements of training. Achieving this
378 would allow practitioners to prescribe an appropriate range of training stimuli whilst
379 also ensuring players are contained within an appropriate overall accumulation of
380 training load (Gabbett, 2016). Furthermore, instantaneous speed can also be
381 monitored in real-time (Barrett et al., 2017; Weaving et al., 2017). This is unlike

382 acceleration data, which can only be monitored post-session. Consequently,
383 acceleration variables can be difficult for the practitioner to translate into the
384 actionable manipulation of training content. Regardless, previous work has reported
385 the peak duration-specific acceleration and relative distance demands to occur at
386 different periods within a match (Delaney et al., 2016). This highlights that the
387 retrospective analysis of acceleration demands during specific training drills is
388 warranted. In particular, within a specific duration of rugby league activity, it is likely
389 that the interplay between the magnitude of instantaneous speed and acceleration
390 plus collision activity would provide the best representation of the 'most demanding'
391 durations of professional rugby league competition.

392 It is therefore recommended that further research be undertaken in order to better
393 understand the interaction between these three components coupled with their own
394 individual peak demands (which may occur at different times to each other) during
395 the peak passages of competition. Ideally, future research should look to the link the
396 peak interactions between these three modes of activity and the associated technical-
397 tactical/skill activities that are completed within such periods.

398 **Conclusions**

399 Aside from very-short-duration bouts (i.e. 10 s), there are trivial differences in the peak
400 relative distances covered between positions during ESL competition. However,
401 adjustables, outside- and full-backs cover substantially greater high-speed- and very-
402 high-speed-distances during the peak 10 min relative distance period than middle-
403 and edge-forwards whilst the forwards positional groups complete a greater number
404 of collisions. There are likely trivial differences between these demands between
405 halves of competition, suggesting that players are likely to be exposed to similar peak
406 intensities for each given period in both halves of the match.

407 **Practical Applications**

- 408 • To simulate the peak running intensities of ESL competition, practitioners
409 should expose positional groups to similar peak relative distances and
410 durations during training.
- 411 • Given the similarities between match halves across durations, programming
412 multiple peak bouts within a training session could help to prepare players for
413 competition.
- 414 • How positions achieve this overall relative distance should differ, with
415 adjustables, outside- and full-backs completing greater high-speed- and very-
416 high-speed-distances.
- 417 • During the peak 10 min running 'intensity' of ESL, forwards complete a greater
418 frequency of collisions and should be exposed to these demands whilst
419 completing similar relative distances.

420 **References**

- 421 Barrett, S. (2017). Monitoring elite soccer players external loads using real-time data.
422 *International Journal of Sports Physiology and Performance*, 2, 1-9.
- 423 Bates, D., Sarkar, D., Bates, M. D., & Matrix, L. (2007). The lme4 package. *R package*
424 *version*, 2(1), 74.
- 425 Boyd, L.J., Ball, K. & Aughey, R.J. (2011). The reliability of MinimaxX accelerometers
426 for measuring physical activity in Australian football. *International Journal of Sports*
427 *Physiology and Performance*, 6, 311-321.
- 428 Cummins, C., & Orr, R. (2015). Analysis of physical collisions in elite National Rugby
429 League match play. *International Journal of Sports Physiology and Performance*, 10
430 (6), 732-739.
- 431 Darrall-Jones J., Roe G., Carney S., Clayton R., Phibbs P., Read D., Weakley J., Till
432 K., & Jones B. (2016). The effect of body mass on the 30-15 intermittent fitness test

433 in rugby union players. *International Journal of Sports Physiology and Performance*,
 434 11(3), 400-403.

435 Delaney, J.A., Duthie, G.M., Thornton, H.R., Scott, T.J., Gay, D., Dascombe, B.J.
 436 (2016). Acceleration-based running intensities of professional rugby league match
 437 play. *International Journal of Sports Physiology and Performance*, 11(6), 802-809.

438 Delaney J.A., Scott T.J., Thornton H.R., Bennett K.J., Gay, D., Duthie, G.M., &
 439 Dascombe, B.J. (2015). Establishing duration-specific running intensities from match-
 440 play analysis in rugby league. *International Journal of Sports Physiology and*
 441 *Performance*, 10(6), 725-731.

442 Gabbett T.J., Jenkins D.G., & Abernethy, B. (2011). Physical collisions and injury in
 443 professional rugby league match-play. *Journal of Science and Medicine in Sport*, 14,
 444 210–215.

445 Gabbett T.J., Jenkins D.G., & Abernethy, B. (2012). Physical demands of professional
 446 rugby league training and competition using microtechnology. *Journal of Science and*
 447 *Medicine in Sport*, 15, 80–86.

448 Gabbett, T.J. (2016). The training-injury prevention paradox: should athletes be
 449 training smarter and harder? *British Journal of Sports Medicine*, 50(5), 273-280.

450 Hopkins, W.G., Marshall, S.W., Batterham, A.M., Hanin, J. (2009). Progressive
 451 statistics for studies in sports medicine and exercise science. *Medicine and Science*
 452 *in Sports and Exercise*, 41, 3-13.

453 Hulin B.T., Gabbett T.J., Johnston R.D., Jenkins D.G. (2017). Wearable
 454 microtechnology can accurately identify collision events during professional rugby
 455 league match-play. *Journal of Science and Medicine in Sport*, 20(7), 638-642.

456 Impellizzeri, F.M., Rampinini, E., & Marcora, S.M. (2005). Physiological assessment
 457 of aerobic training in soccer. *Journal of Sports Sciences*, 23, 583-592.

458 Johnston, R.D., Gabbett, T.J., & Jenkins, D.G. (2014). Applied sport science of rugby
459 league. *Sports Medicine*, 44(8), 1087-1100.

460 Johnston, R.D., Gabbett, T.J., Jenkins, D.G., Hulin, B.T. (2015). Influence of physical
461 qualities on post-match fatigue in rugby league players. *Journal of Science and*
462 *Medicine in Sport*, 18(2), 209-213.

463 Jones B., Till, K., Barlow, M., Lees, M., O'Hara, J.P., & Hind, K. (2015). Anthropometric
464 and three-compartment body composition differences between Super League and
465 Championship Rugby League Players: Considerations for the 2015 season and
466 beyond. *PLoS One*, 10(7), e0133188.

467 Lovell, T.W.J., Sirotic, A.C., Impellizzeri, F.M., & Coutts, A.J. (2013). Factors affecting
468 perception of effort (session rating of perceived exertion) during Rugby League
469 training. *International Journal of Sports Physiology and Performance*, 8, 62-69.

470 Malone, J.J., Lovell, R., Varley, M.C., Coutts, A.J. Unpacking the black box:
471 applications and considerations for using GPS devices in sport. (2017). *International*
472 *Journal of Sports Physiology and Performance*, 12 (Suppl 2), S218-S226.

473 McLellan C.P., Lovell D.I., & Gass G.C. (2011). Performance analysis of elite rugby
474 league match play using global positioning systems. *Journal of Strength and*
475 *Conditioning Research*, 25(6), 1703-1710.

476 McNamara D.J., Gabbett, T.J., Chapman, P., Naughton, G., Farhart, P. (2015).
477 Variability of PlayerLoad, bowling velocity, and performance execution in fast bowlers
478 across repeated bowling spells. *International Journal of Sports Physiology and*
479 *Performance*, 10, 1009-1014.

480 Mullen, T., Highton, J., Twist, C. (2015). The internal and external responses to a
481 forward-specific rugby league simulation protocol performed with and without physical
482 contact. *International Journal of Sports Physiology and Performance*, 10(6), 746-753.

483 Norris, J.P., Highton, J., Hughes, S.F., Twist, C. (2016). The effects of physical contact
 484 type on the internal and external demands during a rugby league match simulation
 485 protocol. *Journal of Sports Sciences*, 34(19), 1859-66.

486 Robertson, S.J. & Joyce, D.G. (2015). Informing in-season tactical periodisation in
 487 team sport: development of a match difficulty index for Super Rugby. *Journal of Sports*
 488 *Sciences*, 33(1), 99-107.

489 Scott, M.T., Scott, T.J., Kelly, V.G. (2016). The validity and reliability of global
 490 positioning systems in team sport: a brief review. *Journal of Strength and Conditioning*
 491 *Research*, 30(5), 1470-1490.

492 Scott, T.J., Thornton, H.R., Scott, M.T.U., Dascombe, B.J., & Duthie, G.M. (2017).
 493 Differences between relative and absolute speed and metabolic thresholds in rugby
 494 league. *International Journal of Sports Physiology and Performance*, 28, 1-21.

495 Soligard, T., Schwellnus, M., Alonso, J.M., Bahr, R., Clarsen, B., Dijkstra, H.P.,
 496 Gabbett, T., ... Engebretsen, L. (2016). How much is too much? (Part 1) International
 497 Olympic Committee consensus statement on load in sport and risk of injury. *British*
 498 *Journal of Sports Medicine*, 50(17), 1030-1041.

499 Twist, C., Highton, J., Waldron, M., Edwards, E., Austin, D., & Gabbett, T.J. (2014).
 500 Movement demands of elite rugby league players during Australian National Rugby
 501 League and European Super League matches. *International Journal of Sports*
 502 *Physiology and Performance*, 9(6), 925-930.

503 Twist, C., Waldron, M., Highton, J., Burt, D., & Daniels, M. (2012). Neuromuscular,
 504 biochemical and perceptual post-match fatigue in professional rugby league forwards
 505 and backs. *Journal of Sports Sciences*, 30, 359-367.

506 Speranza, M.J., Gabbett, T.J., Johnston, R.D., Sheppard, J.M. (2016). Effect of
 507 strength and power training on tackling ability in semi-professional rugby league
 508 players. *Journal of Strength and Conditioning Research*, 30(2), 336-343.

509 Scott T.J., Thornton, H.R., Scott, M.T.U., Dascombe, B.J., & Duthie, G.M. (2017).
 510 Differences between relative and absolute speed and metabolic thresholds in rugby
 511 league. *International Journal of Sports Physiology and Performance*, 28, 1-21.

512 Vanrenterghem J., Nedergaard N.J., Robinson M.A., & Drust, B. (2017). Training load
 513 monitoring in team sports: a novel framework separating physiological and
 514 biomechanical load-adaptation pathways. *Sports Medicine*, 47(11), 2135-2142.

515 Varley M.C., Jaspers, A., Helsen, W.F., & Malone, J.J. Methodological considerations
 516 when quantifying high-intensity efforts in team sport using global positioning system
 517 technology. (2017). *International Journal of Sports Physiology and Performance*,
 518 12(8), 1059-1068.

519 Waldron, M., Twist, C., Highton, J., Worsfold, P., & Daniels, M. (2011). Movement and
 520 physiological match demands of elite rugby league using portable global positioning
 521 systems. *Journal of Sports Sciences*, 29(11), 1223-1230.

522 Waldron, M., Worsfold, P.R., Twist, C., & Lamb, K. (2014). The relationship between
 523 physical abilities, ball-carrying and tackling among elite youth rugby league players.
 524 *Journal of Sports Sciences*, 32(6), 542-49.

525 Weaving, D., Jones, B., Till, K., Marshall, P., Earle, K., Abt, G. (2017). Quantifying
 526 the external and internal loads of professional rugby league training modes:
 527 consideration for concurrent field-based training prescription. *Journal of Strength*
 528 *and Conditioning Research*. doi: 10.1519/JSC.0000000000002242.

529 Weaving, D., Jones, B., Marshall, P., Till, K., & Abt, G. (2017). Multiple measures are
530 needed to quantify training loads in professional rugby league. *International Journal*
531 *of Sports Medicine*, 38(10), 735-740.

532 Weaving D., Whitehead S., Till K., & Jones, B. (2017). Validity of real-time data
533 generated by a wearable microtechnology device. *Journal of Strength and*
534 *Conditioning Research*, 31(10), 2876-2879.

535 Zeileis, A., & Grothendieck, G. (2005). zoo: S3 Infrastructure for Regular and Irregular
536 Time Series. *Journal of Statistical Software*, 14(6), 1–27.

537

538

539

540

541





Figure 1. The standardised mean positional differences plus 90% confidence intervals for the number of collisions completed during the peak 10 min of European Super League rugby. Only substantial differences are detailed. Outside backs vs. fullbacks (-0.07 [-0.47 to 0.34]), adjustables vs. fullbacks (-0.15 [-0.61 to 0.32]), outside backs vs. adjustables (0.09 [-14.59 to 14.76]) and middle forwards vs. edge forwards (-0.06 [-3.76 to 3.64]) were all *unclear*.



555 **Figure 2. Standardised mean positional differences plus 90% confidence**
556 **intervals in high-speed-distance (5 to 7 m·s⁻¹) completed during the peak 10**
557 **min of European Super League rugby.**

558

559

560

561

562

563

564


565

566

567

568

569



570 **Figure 3. Standardised mean differences plus 90% confidence intervals for the**
 571 **positional differences in very-high-speed-distance (> 7 m·s⁻¹) completed**
 572 **during the peak 10 min of European Super League rugby.**

573

574

575

576

577

578

Table 1. The mean \pm standard deviation of duration-specific peak 1st and 2nd half relative distances of European Super League rugby and the concurrent number of collisions, and high-speed- and very-high-speed-distances covered during the peak 10-minute relative distances.

	10s (m·min ⁻¹)	30s (m·min ⁻¹)	1-min (m·min ⁻¹)	5-min (m·min ⁻¹)	10-min (m·min ⁻¹)	10-min HSD (m)	10-min VHSD (m)	Collisions (n)
Fullback								
1 st Half	320.8 \pm 10.6	209.7 \pm 5.2	169.7 \pm 3.8	118.7 \pm 2.8	102.9 \pm 2.3	89.2 \pm 8.4	9.3 \pm 2.1	5.2 \pm 3.0
2 nd Half	331.4 \pm 10.9	209.5 \pm 5.3	167.7 \pm 3.8	115.2 \pm 2.8	101.6 \pm 2.3	80.9 \pm 8.6	4.8 \pm 2.2	5.7 \pm 3.6
Outside back								
1 st Half	325.2 \pm 6.3	200.5 \pm 3.3	169.7 \pm 3.8	110.9 \pm 1.9	96.1 \pm 1.8	82.6 \pm 5.2	10.1 \pm 1.1	5.1 \pm 2.6
2 nd Half	325.2 \pm 6.3	203.1 \pm 3.3	167.7 \pm 3.8	106.4 \pm 1.9	92.3 \pm 1.8	64.5 \pm 5.2	5.9 \pm 1.1	4.9 \pm 2.5
Adjustable								
1 st Half	313.0 \pm 7.1	200.3 \pm 3.7	160.7 \pm 2.4	113.8 \pm 2.1	99.9 \pm 2.0	70.1 \pm 5.9	5.5 \pm 1.3	5.5 \pm 3.3
2 nd Half	322.5 \pm 6.9	205.8 \pm 3.6	159.5 \pm 2.4	111.6 \pm 2.1	96.9 \pm 1.9	65.7 \pm 5.8	6.5 \pm 1.2	5.2 \pm 3.2
Middle forward								
1 st Half	291.5 \pm 6.2	195.5 \pm 3.2	163.1 \pm 2.4	111.1 \pm 1.9	99.0 \pm 1.8	48.7 \pm 5.1	1.9 \pm 1.1	9.7 \pm 2.6
2 nd Half	281.7 \pm 6.3	195.8 \pm 3.2	160.8 \pm 2.4	106.0 \pm 1.9	94.2 \pm 1.8	39.2 \pm 5.1	0.7 \pm 1.2	9.1 \pm 2.8
Edge forward								
1 st Half	296.2 \pm 7.9	191.8 \pm 3.9	159.9 \pm 2.9	110.0 \pm 2.2	99.3 \pm 2.0	60.6 \pm 5.4	2.0 \pm 1.5	8.9 \pm 2.7
2 nd Half	298.3 \pm 7.7	197.8 \pm 3.8	160.4 \pm 2.8	105.9 \pm 2.1	95.1 \pm 1.9	53.9 \pm 6.1	2.4 \pm 1.5	8.2 \pm 2.7

HSD = high-speed-distance (5-7 m·s⁻¹); VHSD = very-high-speed-distance (>7 m·s⁻¹)

Table 2. Raw mean positional differences [90% confidence limits] and likelihoods in peak relative distances across durations

	10s (m·min⁻¹)	30s (m·min⁻¹)	1 min (m·min⁻¹)	5 min (m·min⁻¹)	10 min (m·min⁻¹)
FB vs. OB	0.96 [-14.2 to 16.1] <i>Unclear</i>	7.8 [0.4 to 15.2] <i>Possibly trivial</i>	8.6 [3.2 to 14.1] <i>Possibly trivial</i>	8.3 [4.4 to 12.1] <i>Likely trivial</i>	8.0 [5.2 to 10.9] <i>Likely trivial</i>
FB vs. ADJ	8.38 [-7.7 to 24.4] <i>Unclear</i>	6.5 [-1.5 to 14.6] <i>Likely trivial</i>	5.3 [-0.7 to 11.3] <i>Likely trivial</i>	4.2 [-0.1 to 8.5] <i>Very likely trivial</i>	3.8 [0.5 to 7.1] <i>Almost certainly trivial</i>
FB vs. MF	39.6 [24.2 to 55.0] <i>Almost Certainly ↑</i>	14.0 [6.3 to 21.7] <i>Likely ↑</i>	6.7 [1.0 to 12.5] <i>Likely trivial</i>	8.4 [4.3 to 12.6] <i>Possibly trivial</i>	5.7 [2.3 to 9.1] <i>Very likely trivial</i>
FB vs. EF	28.9 [12.3 to 45.4] <i>Very likely ↑</i>	14.8 [6.6 to 23.0] <i>Likely ↑</i>	8.5 [2.4 to 14.7] <i>Possibly trivial</i>	9.0 [4.6 to 13.4] <i>Possibly trivial</i>	5.0 [1.4 to 8.6] <i>Very likely trivial</i>
OB vs. ADJ	7.4 [-3.8 to 18.6] <i>Possibly Trivial</i>	-1.3 [-7.0 to 4.4] <i>Very likely trivial</i>	-3.3 [-7.6 to 1.0] <i>Almost certainly trivial</i>	-4.0 [-7.1 to -1.0] <i>Almost certainly trivial</i>	-4.2 [-6.6 to 1.8] <i>Almost certainly trivial</i>
OB vs. MF	38.6 [27.8 to 49.4] <i>Almost Certainly ↑</i>	6.2 [0.5 to 11.8] <i>Likely trivial</i>	-1.9 [-6.2 to 2.5] <i>Almost certainly trivial</i>	0.1 [-3.1 to 3.4] <i>Almost certainly trivial</i>	-2.4 [-5.3 to 0.6] <i>Almost certainly trivial</i>
OB vs. EF	27.9 [15.6 to 40.2] <i>Very likely ↑</i>	7.0 [0.7 to 13.3] <i>Likely trivial</i>	-0.1 [-4.9 to 4.7] <i>Almost certainly trivial</i>	0.7 [-2.8 to 4.2] <i>Almost certainly trivial</i>	-3.0 [-6.1 to 0.1] <i>Almost certainly trivial</i>
ADJ vs. MF	31.2 [19.6 to 42.8] <i>Almost certainly ↑</i>	7.4 [1.4 to 13.5] <i>Likely trivial</i>	1.4 [-3.2 to 6.1] <i>Almost certainly trivial</i>	4.2 [0.71 to 7.6] <i>Almost certainly trivial</i>	1.83 [-1.3 to 5.0] <i>Almost certainly trivial</i>
ADJ vs. EF	20.5 [7.5 to 33.5] <i>Likely ↑</i>	8.3 [1.6 to 15.0] <i>Possibly trivial</i>	3.2 [-1.9 to 8.3] <i>Very likely trivial</i>	4.7 [1.0 to 8.5] <i>Very likely trivial</i>	1.17 [-2.2 to 4.5] <i>Almost certainly trivial</i>

579	MF vs. EF	-10.7 [-22.0 to 0.7]	0.9 [-4.7 to 6.4]	1.8 [-2.3 to 5.9]	0.6 [-2.3 to 3.5]	-0.7 [-2.8 to 1.47]
580		<i>Possibly ↓</i>	<i>Almost certainly</i> <i>trivial</i>	<i>Almost certainly</i> <i>trivial</i>	<i>Almost certainly</i> <i>trivial</i>	<i>Almost certainly</i> <i>trivial</i>

581 FB = fullback; OB = outside back; ADJ = adjustables; MF = middle forward; EF = edge forward. The direction of
582 difference is in relation to the first named positional group.

583