



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Allen, H and Hull, J and Backhouse, SH and Price, OJ (2018) The Allergy Questionnaire for Athletes: A simple screening tool for the assessment of exercise-induced bronchoconstriction. In: British Association of Sport and Exercise Medicine 2018, 04 October 2018 - 05 October 2018, Leeds. (Unpublished)

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/5391/>

Document Version:

Conference or Workshop Item (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

**THE ALLERGY QUESTIONNAIRE FOR ATHLETES: A SIMPLE SCREENING TOOL FOR THE ASSESSMENT  
OF EXERCISE-INDUCED BRONCHOCONSTRICTION**

**H. Allen<sup>1</sup>**, JH Hull<sup>2</sup>, SH Backhouse<sup>1</sup>, OJ Price<sup>1</sup>

<sup>1</sup>Carnegie School of Sport, Leeds Beckett University, Leeds, United Kingdom (UK); <sup>2</sup>Department of Respiratory Medicine, Royal Brompton Hospital, London, UK

**Aim:** Evaluating respiratory symptoms in athletic individuals can be difficult and therefore robust clinical assessment tools are required (1). The Allergy Questionnaire for Athletes (AQUA) is validated to assess allergic +/- respiratory symptoms (2). The purpose of this study was therefore to determine the value of AQUA as a screening tool to confirm or refute exercise-induced bronchoconstriction (EIB) in athletes.

**Methods:** One-hundred and forty-seven athletes (male:  $n = 100$ ; female:  $n = 47$ ) (age:  $32 \pm 9$  years) completed AQUA and performed spirometry pre-and-post a eucapnic voluntary hyperpnoea challenge (EVH). A positive AQUA was determined as a score  $\geq 5$  (2). Sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were calculated and evaluated against objective evidence of EIB: (EVH threshold [-10%  $\Delta FEV_1$  at two consecutive time-points] and [-15%  $\Delta FEV_1$  at one time-point]) (3). Diagnostic accuracy was calculated using receiver operating characteristics area under the curve (ROC-AUC).

**Results:** All participants demonstrated normal baseline lung function ( $FEV_1$  % predicted  $>80\%$ ). The prevalence of EIB was 16% (-10%  $\Delta FEV_1$ ) and 11% (-15%  $\Delta FEV_1$ ) (mean  $\Delta FEV_1 = -7.4\% \pm 7.9$ ). Eighty-seven (59%) provided a positive AQUA score (range: 0-25). ROC-AUC for AQUA was 65% (-10%  $\Delta FEV_1$ ) and 69% (-15%  $\Delta FEV_1$ ). A negative AQUA score was highly predictive of negative EVH test outcome. Sensitivity, specificity, PPV and NPV are presented in Table 1.

**Conclusion:** AQUA is a simple screening tool that provides value in ruling out EIB, and should be considered during clinical assessment (i.e. inform referral for objective testing) or as a 'first-step' as part of screening interventions. It is important to acknowledge that a positive AQUA score should not be used to confirm EIB, in the absence of indirect bronchoprovocation. The development and validation of an athlete specific questionnaire to confirm EIB remains an avenue for future research.

1. Price OJ, Hull JH, Ansley L, Thomas M, Eyles C. Exercise-induced bronchoconstriction in athletes—a qualitative assessment of symptom perception. *Respiratory medicine*. 2016;120:36-43.
2. Bonini M, Braido F, Baiardini I, Del Giacco S, Gramiccioni C, Manara M, et al. AQUA: allergy questionnaire for athletes. Development and validation. *Med Sci Sports Exerc*. 2009;41(5):1034-41.
3. Price OJ, Ansley L, Levai IK, Molphy J, Cullinan P, Dickinson JW, et al. Eucapnic voluntary hyperpnea testing in asymptomatic athletes. *American journal of respiratory and critical care medicine*. 2016;193(10):1178-80.

**Table 1.** AQUA sensitivity, specificity, PPV and NPV for the detection of EIB.

<b>Population: <math>n = 147</math></b>						
Prevalence (-10% $\Delta$ FEV <sub>1</sub> ): 16%						
Prevalence (-15% $\Delta$ FEV <sub>1</sub> ): 11%						
<b>EVH</b>						
<b>-10% <math>\Delta</math>FEV<sub>1</sub></b>						
<b>-15% <math>\Delta</math>FEV<sub>1</sub></b>						
		<b>+</b>	<b>-</b>	<b>+</b>	<b>-</b>	
<b>AQUA score</b>	<b>+</b>	20	70	+	15	75
	<b>-</b>	3	54	-	1	56
<b>AQUA score</b>						
<b>-10% <math>\Delta</math>FEV<sub>1</sub></b>						
<b>-15% <math>\Delta</math>FEV<sub>1</sub></b>						
<b>Sensitivity (%)</b>		87			94	
<b>Specificity (%)</b>		44			43	
<b>PPV (%)</b>		22			17	
<b>NPV (%)</b>		95			98	

**Presenting author details:**

**Name:** Mr Hayden Allen BSc (Hons.) MRes

**E-mail.** [hayden.allen@leedsbeckett.ac.uk](mailto:hayden.allen@leedsbeckett.ac.uk)

**Tel.** 01138122042

**Mailing address:** Carnegie School of Sport, Leeds Beckett University, Cavendish Hall, Headingley Campus, Leeds, LS6 3QU.