



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Goff, L and Huang, P and Silva, MJ and Bordoli, C and Enayat, EZ and Molaodi, OR and Cassidy, A and Maynard, M and Harding, S (2019) Associations of dietary intake with cardiometabolic risk in a multi-ethnic cohort: a longitudinal analysis of the Determinants of Adolescence, now young Adults, Social well-being and Health (DASH) study. British Journal of Nutrition. ISSN 0007-1145 DOI: <https://doi.org/10.1017/S0007114519000291>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/5704/>

Document Version:

Article (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

# Accepted manuscript

**Associations of dietary intake with cardiometabolic risk in a multi-ethnic cohort: a longitudinal analysis of the Determinants of Adolescence, now young Adults, Social well-being and Health (DASH) study**

Louise M. Goff<sup>1,2</sup>, Peiyuan Huang<sup>1</sup>, Maria J. Silva<sup>1</sup>, Claire Bordoli<sup>1</sup>, Elli Z. Enayat<sup>1</sup>, Oarabile R. Molaodi<sup>3</sup>, Aidan Cassidy<sup>1</sup>, Maria Maynard<sup>4</sup>, Seeromanie Harding<sup>1</sup>

**Affiliations:** <sup>1</sup>Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, London SE1 9NH <sup>2</sup>Department of Diabetes, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, London SE1 1UL; <sup>3</sup>MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow; <sup>4</sup>School of Clinical & Applied Sciences, Leeds Beckett University, Leeds.

**Pubmed indexing:** Goff, Huang, Silva, Bordoli, Enayat, Molaodi, Cassidy, Maynard, Harding.

**Corresponding author:** Dr Louise M. Goff, Diabetes Research Group, Departments of Diabetes & Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, Waterloo Campus, Franklin-Wilkins Building, Room 3.87, London, SE1 9NH, United Kingdom. T: +44 (0)20 7848 4273; E: [louise.goff@kcl.ac.uk](mailto:louise.goff@kcl.ac.uk)

**Short running title:** Ethnicity, diet and cardiometabolic risk

**Keywords:** ethnicity, breakfast, diet, cardiometabolic risk, African, Caribbean, South Asian

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

10.1017/S0007114519000291

# Accepted manuscript

## Abbreviations:

BMI: Body Mass Index

CHD: Coronary Heart Disease

DASH: Determinants of Adolescent Social well-being and Health

T2D: type 2 diabetes

WHtR: waist to height ratio

## ABSTRACT

Unfavourable dietary habits, such as skipping breakfast are common among ethnic minority children and may contribute to inequalities in cardiometabolic disease. We conducted a longitudinal follow-up of a subsample of the UK multi-ethnic Determinants of Adolescent Social well-being and Health (DASH) cohort, which represents the main UK ethnic groups and is now aged 21-23 years. We aimed to describe longitudinal patterns of dietary intake and investigate their impact on cardiometabolic risk in young adulthood. Participants completed a dietary behaviours questionnaire and a 24-hr dietary intake recall; anthropometry, blood pressure, total and HDL-cholesterol and HbA1c were measured. The cohort consisted of 107 White British, 102 Black Caribbean, 132 Black African, 98 Indian, 111 Bangladeshi/Pakistani and 115 Other/Mixed ethnicity. Unhealthful dietary behaviours such as skipping breakfast and low intakes of fruit and vegetables were common (56%, 57% and 63%, respectively). Rates of skipping breakfast and low fruit and vegetable consumption were highest among Black African and Black Caribbean participants. BMI and cholesterol levels at 21-23 years were higher among those who regularly skipped breakfast at 11-13 years (BMI 1.41 (95% CI 0.57, 2.26)  $p=0.001$ ; cholesterol 0.15 (95% CI -0.01, 0.31)  $p=0.063$ ) and 21-23 years (BMI 1.05 (95% CI 0.22, 1.89)  $p=0.014$ ; cholesterol 0.22 (95% CI 0.06, 0.37)  $p=0.007$ ). Childhood breakfast skipping is more common in certain ethnic groups and is associated with cardiometabolic risk factors in young adulthood. Our findings highlight the importance of targeting interventions to improve dietary behaviours such as breakfast consumption at specific population groups.

# Accepted manuscript

## INTRODUCTION

Ethnic inequalities in a wide range of chronic diseases have been documented <sup>(1)</sup>, with risk factors evident early in life <sup>(2)</sup>. Children and youth from ethnic minority groups are more likely to be overweight or obese than White-European populations <sup>(3; 4)</sup>. The cardiometabolic risks associated with obesity are well known, therefore it is not surprising that some ethnic minority populations suffer disproportionately high rates of chronic diseases such as type 2 diabetes (T2D), hypertension and coronary heart disease (CHD) <sup>(1;5)</sup>.

Ethnic inequalities in obesity and chronic disease risk may be explained by differences in diet and lifestyle. Differences in the nutritional composition of the diet are apparent; a recent cohort study of children in the UK reported significantly higher energy, fat and sugar intakes among South Asians, while Black Africans demonstrated lower intakes of fat and saturated fat but higher intakes of carbohydrate, compared to White British <sup>(6)</sup>. Unfavourable dietary habits, such as regularly skipping breakfast and consumption of sugar-sweetened fizzy drinks and fast foods, are also more common among children and adolescents from some ethnic minority groups <sup>(7; 8; 9; 10)</sup>. The detrimental impact of skipping breakfast, and fast food and fizzy drink consumption on childhood and adolescent health has been documented through a number of cohort studies, which have shown that those who regularly skip breakfast have an increased risk of obesity and detrimental cardiometabolic risk profiles <sup>(10; 11; 12)</sup>. Additionally, studies have shown that children who skip breakfast regularly have poorer quality diets, particularly higher energy density, saturated fat and lower micronutrient intakes <sup>(13)</sup>. More recently longitudinal studies have investigated the prevalence of these behaviours in young adulthood and the detrimental impact of childhood behaviours on adult health <sup>(11; 14; 15)</sup>.

Young adulthood is recognised as an important transition period, with increasing responsibilities and the development of independent health behaviours. The Australian Childhood Determinants of Adult Health (CDAH) study found skipping breakfast in both childhood and young adulthood to be associated with larger waist circumferences, higher fasting insulin, and higher total- and LDL-cholesterol concentrations <sup>(11)</sup>, while in the US-based Project EAT socio-economic status was found to be a principal determinant of increases in breakfast skipping between childhood and adulthood <sup>(15)</sup>. To date there have been no longitudinal cohort studies to describe ethnic patterns in adolescent and young adult nutritional intake and dietary behaviours, and to determine their longitudinal effects on body weight and cardiometabolic risk factors. The 'DASH 10 years on' study is the first multi-ethnic cohort of young adults in the UK and provides longitudinal follow-up data for a subset of the Determinants of Adolescent Social well-being and Health (DASH) cohort, who are now young adults (21-23 years). We aimed to describe longitudinal patterns of dietary intake in the

# Accepted manuscript

multi-ethnic Determinants of Adolescent, now young Adults, Social well-being and Health cohort, and investigate the impact of dietary intake on cardiometabolic risk in young adulthood.

## **SUBJECTS AND METHODS**

### *Study Design*

Details of the DASH study can be found at <http://dash.sphsu.mrc.ac.uk/> and in a published cohort profile <sup>(16)</sup>. The original cohort of 6643 students, aged 11-13y, was recruited between 2002 and 2003, from 51 secondary schools in 10 London boroughs and sampled to represent the main ethnic groups of the UK population. A pilot follow-up study on 665 participants was completed in March 2014; dietary data was collected from 627 participants who took part in the follow-up.

The study was approved by the National Health Service South East Scotland Research Ethics Committee (Ref:11/SS/0051). Written informed consent was obtained from participants. The study is registered on clinicaltrials.gov ID: NCT03283332.

### *Participants*

The subsample consisted of 107 White British, 102 Black Caribbean, 132 Black African, 98 Indian, 111 Bangladeshi or Pakistani and 115 Other (mainly mixed) ethnicities, and was purposively sampled to be representative by gender and socio-economic circumstances (SEC) across the 10 London boroughs and 51 schools. Ethnicity was self-declared, measured by self-report using Census categories, checked against reported parental ethnicity and grandparents' country of birth. Bangladeshis and Pakistanis were approximately equally represented and combined because of relatively small sample sizes; participants declaring their ethnicity as White European, White Irish, Chinese, mixed ethnicity were classified as 'Other ethnicity'.

### *Procedures*

Measurement protocols and questionnaires can be found at <http://dash.sphsu.mrc.ac.uk/>. Nurses and research assistants were trained in the assessment methods by the research team for 1 week prior to the start of fieldwork, and were re-certified at 6 monthly intervals. All equipment was checked and calibrated regularly by the field supervisors. Participants were given a choice of locations (local general practitioner's surgeries, local community pharmacies, Clinical Research Centres and Kings College London) and assessments took about 2 hours.

### *Dietary behaviours and intake*

## Accepted manuscript

At age 11-13 and 21-23 years participants completed a short questionnaire regarding their consumption of breakfast, fruit and vegetables. For breakfast consumption, participants were given a choice of four responses (every day, 3-4 days a week, 1-2 days a week, never/hardly ever). Fruit and vegetable consumption were separate questions with a choice of seven responses for each (5 or more portions a day, 4 portions a day, 3 portions a day, 2 portions a day, 1 portions a day, I eat fruit/vegetables some days but not every day, I never eat fruit/vegetables). Examples of portions (e.g. handful of grapes, an apple, serving spoon of vegetables, bowl of salad) were given alongside the questions for more accurate estimation of portion size. In addition, at age 11-13 years participants answered an additional question regarding consumption of fizzy drinks, for which a choice of five answers were available (every day, most days, weekly, less than weekly, never).

At 21-23 years dietary intake was assessed using the standardised triple-pass 24-hour recall methodology <sup>(17)</sup>. The study investigator, trained in dietary assessment methods, interviewed the participant face-to-face to recall their food and beverage intake for the preceding 24 hours, extracting details regarding food types, brands, recipes and cooking methods. Detailed information was collected for pre-packaged foods including brands and portion sizes, and full recipes and cooking methods were recorded for home-prepared foods and dishes. Portion sizes were confirmed using a photographic food atlas <sup>(18)</sup> and household measures.

Dietary intake records were coded for analysis by a single coder (CB) and analysed for food and nutrient composition using WISP software, which is based on McCance and Widdowson's *The Composition of Foods*, 6<sup>th</sup> edition. Where food items were not available within the WISP database new entries were created, particularly for home-prepared recipes. Nutritional composition information was sought from food labels and manufacturer websites for branded items, and information for cultural foods were sought from our own database of African and Caribbean dishes <sup>(19)</sup>, the West African Food Table <sup>(20)</sup>, and recipe books. Food weights were assigned from the photo atlases that were used in the interviews. Where food weights were not indicated, average adult portion sizes from the National Diet and Nutrition Survey were used <sup>(21)</sup>.

The quality of reporting in the 24-hour recall records was assessed using a scoring system developed by the researchers. Seven components were identified as important to assess the quality of dietary information (portion sizes, brand information, food description, cooking methods, recipe information, beverage information, intake timings) and these were scored on a scale of 0-2, in which zero indicated no information provided, 1 indicated incomplete information provided, and 2

## Accepted manuscript

indicated complete information provided. The scores assigned to each category were totalled and a percentage calculated, indicating the quality of dietary information obtained. For dietary recalls which did not include categories such as composite or homemade dishes these categories were excluded when calculating the diet quality percentage. To test the sensitivity of the protocol to assess the quality of the dietary information obtained, two researchers trialled a sample of recalls deemed of poor (n=10) and high quality (n=10). Researchers compared quality scores in order to remain consistent. Overall the mean quality score of the dietary records was 69%, ranging from 21 – 100%, with 85% of the recalls scoring  $\geq 50\%$  for the quality of the data recording. In preliminary analysis we investigated the impact of reporting quality on the dietary data to determine whether recalls of poorer quality should be excluded from our final analyses; we found no statistical difference in reported energy intake when records scoring  $< 50\%$  were excluded, therefore, the final analyses included data from the whole cohort.

Macronutrient intakes were expressed as percentage of total energy intake. Dietary fibre (AOAC method) was expressed in grams per 1000 kcals and sodium intake in milligrams per 1000 kcals. Energy intakes were compared to estimated energy requirements to assess potential under-reporting; age and sex specific standard equations were used to estimate basal metabolic rate (BMR)<sup>(22)</sup>. These were multiplied by a lower cut-off physical activity level (PAL) of 1.2 to provide estimates of minimum total energy expenditure and recognise probable under-reporting<sup>(23)</sup>; 36% of participants reported an energy intake below this cut-off and were considered probable under-reporters (rates did not vary by gender, ethnicity or educational attainment (data not shown)). Data from these participants were not excluded from further analyses.

### ***Body size, biological, and social measures***

Height was measured without shoes, using portable stadiometers, to the nearest 0.1 cm. Weight was measured in light clothing using Salter electronic scales, to the nearest 0.1 kg. Body Mass Index (BMI) was derived as weight (kg)/height (m)<sup>2</sup>; BMI  $> 25$  kg/m<sup>2</sup> was defined as overweight/obese. Waist circumference (cm) was measured midway between the 10<sup>th</sup> rib and the top of the iliac crest, and 0.5 cm subtracted to correct for measurement over T-shirt or vest. The mean of two duplicate measures was derived, with waist to height ratio calculated. Systolic blood pressure (sBP) was measured using validated OMRON M5-1 instruments and appropriately sized cuffs, after sitting quietly for a timed 5 minutes, with  $> 1$  minute between 3 subsequent readings. The mean of the second and third readings was used in analysis. A 25 ml blood sample was taken by venepuncture for the assessment of total and HDL cholesterol and glycated haemoglobin (HbA1c); lipids were

# Accepted manuscript

analysed using colour photometric assays on an automated clinical chemistry analyser (Siemens Advia 2400 analyser), HbA1c was analysed using HPLC (Premier 9210 analyser). Physical activity was measured using ActivPal™ monitors (PAL Technologies, Glasgow UK). The monitor, validated for identifying sitting, standing and walking<sup>(24)</sup> was worn continuously on the front of the thigh for 5 days; the number of minutes spent sitting or in sedentary activity per day was derived and used as the measure of physical inactivity.

A self-complete questionnaire measured current smoking behaviour and SEC (i.e. educational attainment and employment status).

## *Statistical Analysis*

The dietary behaviours (skipping breakfast, fizzy drink, fruit, and vegetable consumption) were recoded into dichotomous variables as follows: skipping breakfast was defined as not consuming breakfast every day; fizzy drink consumption was defined as consuming fizzy drinks every day or most days; low daily fruit consumption was defined as consuming less than 2 portions of fruit per day; and low daily vegetable consumption was defined as consuming less than 3 portions of vegetables per day. Missing data in each categorical variable were recoded as 'not stated'. Missing data among continuous variables were 0.001% for gender, 0.15% for BMI, 4.5% for waist:height ratio, 25.0% for total cholesterol, 27.2% for HDL-cholesterol, 27.7% for HbA1c, 0.15% for systolic blood pressure, and 5.6% for all six nutritional composition variables.

Continuous variables were tested for normality using the Shapiro-Wilk test, and no transformation was necessary. Ethnic differences in anthropometry, cardiometabolic risks factors and dietary intake were performed using ANOVA with post-hoc analysis. Longitudinal patterns in dietary behaviours (skipping breakfast, fruit and vegetable intake) from 11-13 years to 21-23 years were examined with mixed-effects logistic regression, adjusted for age, gender and ethnicity. Multivariate linear regression was used to examine the association between dietary behaviours at 11-13 years and at 21-23 years and cardiometabolic risk factors at 21-23 years, adjusted for gender and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), smoking (current smoker/non-smoker), energy intake and saturated fat intake at 21-23 years. N=664, 635, 499, 484, 481, and 664, respectively, for linear regression models with BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c, and systolic blood pressure, respectively, as the outcome. Multivariate linear regression was used to examine the association between dietary behaviours at



# Accepted manuscript

11-13 years and at 21-23 years and nutritional composition of the diet at 21-23 years, adjusted for gender, and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), and smoking (current smoker/non-smoker) at 21-23 years. N=628 for linear regression models for all six nutritional composition variables as the outcome. All analyses were performed using STATA (version 13; StataCorp LP, College Station, Texas, USA) and statistical significance was considered at  $p<0.05$ .

## RESULTS

The cohort consisted of 329 men and 336 women. The representation by ethnicity was: 107 White British, 102 Black Caribbean, 132 Black African, 98 Indian, 111 Bangladeshi or Pakistani and 115 Other (mainly mixed ethnicity). The mean age of the cohort was 22.7 years, mean BMI 24.7 kg/m<sup>2</sup>, 39% were overweight or obese (BMI>25 kg/m<sup>2</sup>), 24% were smokers and 47% were educated to degree level. Rates of smoking were highest among White British (47%) and lowest among Black African (11%) participants. Attainment of a university degree was highest among Black African (58%) and lowest among Black Caribbean (34%) participants.

Dietary data, collected for 627 participants, are shown in Table 1 alongside anthropometry and cardiometabolic risk factors for the whole cohort and for each ethnic group at 21-23 years. At 21-23 years, overall, 56% of participants reported skipping breakfast on a regular basis, 57% reported consuming less than 2 portions of fruit a day, and 63% reported consuming less than 3 portions of vegetables a day. Ethnic differences in diet, anthropometry and cardiometabolic risk were evident. Black Caribbean and Black African participants were significantly heavier than the Indian and Pakistani/Bangladeshi participants, and had higher BMI compared with White British and Indian participants but there were no differences in waist circumference or waist to height ratio. Total cholesterol was lowest in the Black African and highest in the Pakistani/Bangladeshi participants, while HDL-cholesterol was highest in the Black Caribbean and Black African participants, and lowest in the Indian participants. The highest proportion of breakfast skipping occurred among the Black Caribbean (66%) and Black African (64%) groups and was lowest in the Indian group (46%), although there were no statistical differences. Saturated fat intakes were lowest among the Black African and Indian participants and highest among the Pakistani/Bangladeshi participants. Dietary fibre intake was highest in the White British and Indian groups and was significantly higher than the Black groups or the Pakistani/Bangladeshi participants. The Pakistani/Bangladeshi participants were significantly more likely to have a low daily consumption of vegetables but there were no ethnic differences in fruit consumption.

## Accepted manuscript

We examined longitudinal patterns in the diet behaviours; there were significant increases in the rates of skipping breakfast (37 vs 56%  $p=0.035$ ), low consumption of fruit (46 vs 57%  $p=0.024$ ) and of vegetables (57 vs 63%  $p=0.021$ ) from 11-13 years to 21-23 years among the whole cohort (Figure 1). Age was associated with risk of unhealthful diet behaviours, independent of gender and ethnicity. When they were 21-23 years old, individuals were more likely to skip breakfast (OR: 2.65 (95% CI 2.00, 3.50),  $p<0.001$ ), have low consumption of fruit (OR: 1.39 (95% CI 1.08, 1.79),  $p=0.011$ ), and vegetables (OR: 1.28 (95% CI 0.99, 1.66),  $p=0.059$ ) than when they were 11-13 years old, although the association for low vegetable consumption did not reach statistical significance. After adjusting for gender, ethnicity and educational attainment, skipping breakfast regularly at 11-13 years was a significant determinant of skipping breakfast in young adulthood (OR: 2.36 (95% CI 1.62, 3.44),  $p<0.001$ ). Low fruit consumption and low vegetable consumption at 21-23 years were also significant determinants of skipping breakfast at 21-23 years (Fruit: OR: 2.21 (95% CI 1.54, 3.18),  $p<0.001$ ; Vegetables: OR: 1.91 (95% CI 1.32, 2.76),  $p=0.001$ ) but low fruit consumption and low vegetable consumption at 11-13 years were not significant determinants of skipping breakfast in young adulthood (Fruit: OR: 1.29 (95% CI 0.88, 1.88),  $p=0.190$ ; Vegetables: OR: 1.32 (95% CI 0.91, 1.91),  $p=0.150$ ). Figure 1 shows the rates of (a) skipping breakfast, (b) low fruit and (c) low vegetable consumption at 11-13 years and 21-23 years by ethnic group. Small ethnic sample sizes precluded rigorous testing of ethnic specific effects, nonetheless the descriptive data indicates ethnic variations. Increases in the rates of skipping breakfast were evident in all ethnic groups, with the greatest increase occurring in the White British group. Prevalence of low fruit consumption increased in all ethnic groups, with the Black African group experiencing the greatest increase. While the rates of low vegetable consumption did not increase in either the White British or 'Other' ethnic groups but increased in all other ethnicities, with the greatest increase occurring among the Black African group.

### *Longitudinal associations between dietary behaviours, anthropometric and cardiometabolic risk factors*

The influence of the dietary behaviours during both adolescence and young adulthood on cardiometabolic risk factors during young adulthood were investigated in multivariate analysis, adjusting for gender, ethnicity, educational attainment, smoking, physical activity, energy and saturated fat intake (Table 2). Skipping breakfast at 11-13 years was a significant determinant of BMI at 21-23 years (1.41 (95% CI 0.57, 2.26),  $p<0.001$ ) as was skipping breakfast at 21-23 years, although the effect was slightly attenuated in this age group (1.05 (95% CI 0.22, 1.89),  $p=0.014$ ). Skipping breakfast at both 11-13 years and 21-23 years were also important determinants of total

## Accepted manuscript

cholesterol levels (11-13 years: 0.15 (95% CI -0.01, 0.31),  $p=0.063$ ; 21-23 years: 0.22 (95% CI 0.06, 0.37),  $p=0.007$ ). Low intake of fruit at 11-13 years was a significant determinant of waist to height ratio at 21-23 years (0.01 (95% CI 0.00, 0.03),  $p=0.034$ ), and regular fizzy drink consumption at 11-13 years was a significant determinant of systolic blood pressure at 21-23 years (1.77 (95% CI 0.12, 3.43),  $p=0.036$ ).

### *Longitudinal associations between dietary behaviours and nutritional composition of the diet*

The influence of the dietary behaviours during both adolescence and young adulthood on the nutritional composition of the diet during young adulthood were investigated in multivariate analysis, adjusting for gender, ethnicity, educational attainment, smoking and physical activity (Table 3). Skipping breakfast at both 11-13 years and 21-23 years were significant determinants of lower dietary fibre intakes (11-13 years: -0.81 (95% CI -1.49, -0.12),  $p=0.022$ ; 21-23 years: -1.33 (95% CI -2.01, -0.65),  $p<0.001$ ), as were regular fizzy drink consumption at 11-13 years (-0.72 (95% CI -1.42, -0.02),  $p=0.045$ ), low fruit consumption at 21-23 years (-1.22 (95% CI -1.90, -0.54),  $p<0.001$ ) and low vegetable consumption at 21-23 years (-1.30 (95% CI -2.04, -0.56),  $p=0.001$ ). Low fruit and vegetable consumption at 21-23 years were also important determinants of higher saturated fat intakes (fruit: 0.89 (95% CI -0.01, 1.78),  $p=0.052$ ; vegetables: 1.06 (95% CI 0.09, 2.02),  $p=0.032$ ).

# Accepted manuscript

## DISCUSSION

We have undertaken a longitudinal assessment of nutritional intake and dietary behaviours and their association with cardiometabolic risk factors among an ethnically diverse cohort of young adults in the UK. In this work, we report the detrimental impact of childhood eating behaviours, such as regularly skipping breakfast, on cardiometabolic risk factors in young adulthood. We have identified an increasing prevalence, from childhood to young adulthood, of less healthy dietary behaviours. Furthermore, we have described distinct ethnic patterns, such that skipping breakfast is most prevalent among Black African and Caribbean groups.

The DASH cohort was recruited primarily to investigate ethnic determinants and patterns of health, and through the collection of dietary data we have been able to explore potential predictors of cardiometabolic risk in our young adults. In this study, we found the Black African and Caribbean groups to have the highest body weights and BMIs, however waist circumference and waist to height ratio were comparable to the other groups. In the Indian ethnic group, body weight and BMI were significantly lower than the White British and both Black ethnic groups. These findings are in line with the existing literature and suggest that higher body weight/BMI amongst African ancestry populations is not related to abdominal fat accumulation <sup>(25)</sup>, although fat deposition in the abdominal region is usually more prevalent in South Asian populations <sup>(26)</sup>. The Black Africans exhibited low total cholesterol and high HDL-cholesterol concentrations; these cardiometabolic risk differences may be attributed to dietary differences, namely lower saturated fat, which was also apparent among the Black African participants. Among the Indian participants HDL-cholesterol was lower. Distinctions in nutritional intake were also evident for this group, namely higher intakes of fibre in the Indian group. Ethnic differences in cardiometabolic risk factors have been extensively reported and in general our data corroborate the literature. However, our categorisation of ethnicity, which is more detailed than that which is often employed, highlights some differences between our findings and the existing literature. Higher HbA1c levels amongst African-Americans relative to White-Americans have been previously reported <sup>(27; 28)</sup>; in our study the Black Caribbeans and Black Africans did not exhibit higher HbA1c levels. It is not clear the extent to which higher HbA1c among Black populations is a result of dietary intake versus biological factors. Higher sugar intakes among Black Caribbean compared to Black West African adults have previously been described <sup>(19)</sup>, however we did not find this in our participants, which is in agreement with Donin *et al.* who reported no difference in sugar intake between Black Caribbean and Black African children in the CHASE study <sup>(6)</sup>. Lower total cholesterol amongst Black Africans but not Black Caribbeans is in agreement with findings from the CHASE study <sup>(2)</sup>, and may relate to

## Accepted manuscript

lower saturated fat intake amongst Black Africans but not Black Caribbeans; a finding that is consistent with that of the CHASE study. Historically the African-Caribbean diet has been recognised for its low intake of saturated fat, which has been proposed to contribute to the cardioprotective lipid profile that these ethnicities traditionally exhibited <sup>(29; 30; 31)</sup>. However, dietary acculturation and ‘Westernisation’ of the diet results in increasing fat intake, a change that is more apparent amongst Black Caribbean than Black African communities <sup>(19)</sup>, which may explain why the total cholesterol concentration was only lower amongst the Black African group. Lower HDL-cholesterol amongst South Asian populations has been extensively reported in the literature <sup>(31; 32; 33)</sup>, and the nutritional profile that we identified is also consistent with previous reports <sup>(29; 31; 32; 34)</sup>.

Unhealthy dietary behaviours were highly prevalent among our young adults. In agreement with other longitudinal investigations we saw that adolescent breakfast skipping was predictive of the behaviour in adulthood <sup>(11; 14)</sup>. In our study breakfast skipping was particularly prevalent amongst certain ethnic groups. In agreement with other studies of children and young adults <sup>(10; 14; 35)</sup> our Black African and Black Caribbean participants had the highest rates of breakfast skipping, although the White British group saw the greatest increase in skipping breakfast from childhood to young adulthood. Our longitudinal analysis indicated that skipping breakfast in childhood was an important determinant of BMI and total cholesterol concentration in young adulthood. Whilst there have been a number of longitudinal studies to demonstrate the detrimental effects of skipping breakfast on body weight and the development of obesity, there has been very little demonstration of the long-term effects of this behaviour on cardiometabolic risk factors, such as blood lipids. Our study is the first to focus on an ethnically diverse cohort that represents the main ethnic groups of the UK population. Prior to this the long term evidence was limited to the Australian CDAH study, which studied a predominantly White Australian population, and found skipping breakfast in adolescence to have a unfavourable impact on waist circumference, fasting insulin and total- and LDL-cholesterol <sup>(11)</sup>. Furthermore, adolescent and young adult dietary behaviours were seen to be significant determinants of nutritional composition. Skipping breakfast and regular fizzy drink consumption during adolescence were associated with significantly lower intakes of dietary fibre in young adulthood. In young adulthood skipping breakfast and low consumption of fruit and vegetables were associated with significantly lower dietary fibre and significantly higher saturated fat intakes. These findings corroborate with the literature, demonstrating the detrimental impact of skipping breakfast and other dietary behaviours on the quality of the diet <sup>(13)</sup>; these results are particularly alarming given these unhealthful behaviours are most prevalent among some of our ethnic minority groups, who are most at risk of developing obesity and cardiometabolic diseases, such as T2D. Promoting high fibre intakes and restriction of saturated fat are two of the most

## Accepted manuscript

important public health nutrition campaigns in a bid to reduce chronic disease risk among the general population. Our findings provide a useful insight into dietary behaviours and specific ethnic groups that health promotion campaigns could target in aiming to improve the nutritional intake of the diets of young people.

The limitations of our study warrant consideration. The size of the cohort subsample is relatively small and we did not have statistical power to investigate longitudinal patterns by ethnic group however, it is the first cohort study of young adults to specifically focus on ethnic determinants of health and chronic disease risk. Furthermore, our study contributes to the very limited longitudinal data on the role of adolescent diet on cardiometabolic risk in young adulthood and is the first such study in the UK. We achieved a response rate of 97% in our follow-up, which is remarkably high; this was achieved mainly through the use of social media, which allowed us to locate participants who had moved from their childhood residence. Our data collection focused on a relatively small number of cardiometabolic risk factors, which prevented us from assessing all aspects of risk, for example, the prevalence of Metabolic Syndrome, and making comparisons with the literature in this regard. Although our study is the first longitudinal study of adolescents and young adults to collect nutritional intake data, we were limited in our dietary data collection to assessing a single 24-hour period and therefore we have not assessed micronutrient intakes; this has limited our ability to investigate the role of, for example vitamin D, in determining cardiometabolic risk. Under-reporting is a common issue in dietary data collection; we estimated under-reporting to affect 36% of our recalls, which is comparable to other studies <sup>(36)</sup>. Importantly, under-reporting was not associated with gender, ethnicity or educational attainment, suggesting there is no systematic bias in our dietary data. We did not exclude under-reporters from our analyses; our reporting period was relatively short, and we deemed low energy intakes to be plausible over a 24-hour period. In our dietary behaviours questionnaire, which asked about daily portions of fruit and vegetables, we provided portion size guidance to aid the participants in estimating their daily portions, however our guidance was not exhaustive and did not focus on composite dishes such as soup, which may have led to inaccuracies in reporting. We did not collect data on fizzy drink consumption at 21-23 years due to time restrictions for our data collection visits; sugar sweetened beverages are receiving significant attention at present due to their links with cardiometabolic risk; our data has not enabled us to look at longitudinal patterns of consumption. Our dietary analysis is also limited to focusing on single nutrients and food groups and does not consider interactions between these or reflect the complexity of dietary intake; dietary pattern analysis would enable a better understanding of dietary interactions.

# Accepted manuscript

In conclusion, findings from this cohort of young adults representing the major ethnic groups of the UK population demonstrate the detrimental effects of adolescent dietary behaviours, particularly skipping breakfast on cardiometabolic risk factors. The results also show these behaviours to be particularly prevalent amongst Black African and Caribbean ethnic groups. Our study highlights the importance of targeting health promotion interventions to improve dietary behaviours such as breakfast consumption at specific groups of young adults in the population.

## ACKNOWLEDGEMENTS

We acknowledge the invaluable support of participants and their parents, the Participant Advisory Group, schools, civic leaders, local GP surgeries and community pharmacies, the Clinical Research Centre at Queen Mary University of London, the Clinical Research Facility at University College Hospital, the survey assistants and nurses involved with data collection, Rachel Bonner, senior respiratory physiologist (UCL GOS ICH) for spirometry training and over-reading of data for quality control, the Primary Care Research Network, and Professor Sanders and Cruickshank at the Diabetes and Nutritional Sciences Division at Kings College London for hosting the feasibility study. Seeromanie Harding is the Principal Investigator of DASH. All authors contributed to study design, analyses and writing of the article. The study was funded by the MRC (MC\_U130015185/MC\_UU\_12017/13), Chief Scientist Office (SPHSU13), North Central London Research Consortium and the Primary Care Research Network.

**Data sharing:** The DASH data are available to researchers via a data request to the MRC Social and Public Health Science Unit. Applications and the data sharing policy for DASH can be found at [http://dash.sphsu.mrc.ac.uk/DASH\\_dsp\\_v1\\_November-2012\\_draft.pdf](http://dash.sphsu.mrc.ac.uk/DASH_dsp_v1_November-2012_draft.pdf). It reflects the MRC guidance on data sharing with the aim of making the data as widely and freely available as possible while safeguarding the privacy of participants, protecting confidential data, and maintaining the reputation of the study. All potential collaborators work with a link person, an experienced DASH researcher—to support their access to and analysis of the data. The variable-level metadata is available from the study team and also via the MRC Data Gateway.

**FINANCIAL SUPPORT:** MRC (MC\_U130015185/MC\_UU\_12017/13), Chief Scientist Office (SPHSU13), North Central London Research Consortium and the Primary Care Research Network.

**CONFLICTS OF INTEREST:** the authors declare no conflicts of interest.

# Accepted manuscript

**AUTHORS CONTRIBUTIONS:** SH, MM, LMG designed research; EZE, AC, CB conducted research; PH, CB, MJS, ORM analysed data and performed statistical analyses; LMG wrote manuscripts and had primary responsibility for final content.

## REFERENCES

1. Becker E, Boreham R, Chaudhury M *et al.* (2006) *Health Survey for England 2004. The health of minority ethnic groups*. London: Joint Health Surveys Unit, National Centre for Social Research, Department of Epidemiology and Public Health at the Royal Free and University College Medical School.
2. Donin AS, Nightingale CM, Owen CG *et al.* (2010) Ethnic differences in blood lipids and dietary intake between UK children of black African, black Caribbean, South Asian, and white European origin: the Child Heart and Health Study in England (CHASE). *Am J Clin Nutr* **92**, 776-783.
3. Ogden CL, Flegal KM, Carroll MD *et al.* (2002) Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA* **288**, 1728-1732.
4. Nelson TF, Gortmaker SL, Subramanian SV *et al.* (2007) Disparities in overweight and obesity among US college students. *Am J Health Behav* **31**, 363-373.
5. Zhang Q, Wang Y, Huang ES (2009) Changes in racial/ethnic disparities in the prevalence of Type 2 diabetes by obesity level among US adults. *Ethn Health* **14**, 439-457.
6. Donin AS, Nightingale CM, Owen CG *et al.* (2010) Nutritional composition of the diets of South Asian, black African-Caribbean and white European children in the United Kingdom: the Child Heart and Health Study in England (CHASE). *Br J Nutr* **104**, 276-285.
7. Harding S, Teyhan A, Maynard MJ *et al.* (2008) Ethnic differences in overweight and obesity in early adolescence in the MRC DASH study: the role of adolescent and parental lifestyle. *Int J Epidemiol* **37**, 162-172.
8. Nicklas TA, Myers L, Reger C *et al.* (1998) Impact of breakfast consumption on nutritional adequacy of the diets of young adults in Bogalusa, Louisiana: ethnic and gender contrasts. *J Am Diet Assoc* **98**, 1432-1438.
9. Ritterman Weintraub ML, Fernald LC, Adler N *et al.* (2015) Perceptions of social mobility: development of a new psychosocial indicator associated with adolescent risk behaviors. *Front Public Health* **3**, 62.
10. Donin AS, Nightingale CM, Owen CG *et al.* (2014) Regular breakfast consumption and type 2 diabetes risk markers in 9- to 10-year-old children in the child heart and health study in England (CHASE): a cross-sectional analysis. *PLoS Med* **11**, e1001703.
11. Smith KJ, Gall SL, McNaughton SA *et al.* (2010) Skipping breakfast: longitudinal associations with cardiometabolic risk factors in the Childhood Determinants of Adult Health Study. *Am J Clin Nutr* **92**, 1316-1325.
12. Berkey CS, Rockett HR, Gillman MW *et al.* (2003) Longitudinal study of skipping breakfast and weight change in adolescents. *Int J Obes Relat Metab Disord* **27**, 1258-1266.
13. Szajewska H, Rusczyński M (2010) Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Crit Rev Food Sci Nutr* **50**, 113-119.
14. Merten MJ, Williams AL, Shriver LH (2009) Breakfast consumption in adolescence and young adulthood: parental presence, community context, and obesity. *J Am Diet Assoc* **109**, 1384-1391.
15. Watts AW, Mason SM, Loth K *et al.* (2016) Socioeconomic differences in overweight and weight-related behaviors across adolescence and young adulthood: 10-year longitudinal findings from Project EAT. *Prev Med* **87**, 194-199.
16. Harding S, Whitrow M, Maynard MJ *et al.* (2007) Cohort profile: The DASH (Determinants of Adolescent Social well-being and Health) Study, an ethnically diverse cohort. *Int J Epidemiol* **36**, 512-517.
17. Nelson M, Holmes B, Roberts C *et al.* (2013) *Dietary assessment - Recalls. Diet and physical activity measurement toolkit*. <http://dapa-toolkit.mrc.ac.uk/dietary-assessment/methods/recalls/index.html>: MRC.
18. Nelson, Atkinson, Meyer J (1997) *A photographic atlas of food portion sizes*: MAFF Publications, London, UK.



# Accepted manuscript

19. Goff LM, Timbers L, Style H *et al.* (2014) Dietary intake in Black British adults; an observational assessment of nutritional composition and the role of traditional foods in UK Caribbean and West African diets. *Public Health Nutr*, 1-11.
20. Stadlmayr B, Charrondiere RVE, Bayli R *et al.* (2012) *West African Food Composition Table*. Rome.
21. Department of Health (2011) National Diet and Nutrition Survey: Headline results from Years 1 and 2 (combined) of the rolling programme 2008-9 - 2009-10. England: Department of Health.
22. Henry CJ (2005) Basal metabolic rate studies in humans: measurement and development of new equations. *Public Health Nutr* **8**, 1133-1152.
23. Rennie KL, Coward A, Jebb SA (2007) Estimating under-reporting of energy intake in dietary surveys using an individualised method. *Br J Nutr* **97**, 1169-1176.
24. Dowd KP, Harrington DM, Donnelly AE (2012) Criterion and concurrent validity of the activPAL professional physical activity monitor in adolescent females. *PLoS One* **7**, e47633.
25. Gower BA, Nagy TR, Goran MI (1999) Visceral fat, insulin sensitivity, and lipids in prepubertal children. *Diabetes* **48**, 1515-1521.
26. Ramachandran A, Snehalatha C, Viswanathan V *et al.* (1997) Risk of noninsulin dependent diabetes mellitus conferred by obesity and central adiposity in different ethnic groups: a comparative analysis between Asian Indians, Mexican Americans and Whites. *Diabetes Res Clin Pract* **36**, 121-125.
27. Herman WH, Ma Y, Uwaifo G *et al.* (2007) Differences in A1C by race and ethnicity among patients with impaired glucose tolerance in the Diabetes Prevention Program. *Diabetes Care* **30**, 2453-2457.
28. Ziemer DC, Kolm P, Weintraub WS *et al.* (2010) Glucose-independent, black-white differences in hemoglobin A1c levels: a cross-sectional analysis of 2 studies. *Ann Intern Med* **152**, 770-777.
29. Vyas A, Greenhalgh A, Cade J *et al.* (2003) Nutrient intakes of an adult Pakistani, European and African-Caribbean community in inner city Britain. *J Hum Nutr Diet* **16**, 327-337.
30. Goff LM, Griffin BA, Lovegrove JA *et al.* (2013) Ethnic differences in beta-cell function, dietary intake and expression of the metabolic syndrome among UK adults of South Asian, black African-Caribbean and white-European origin at high risk of metabolic syndrome. *Diab Vasc Dis Res* **10**, 315-323.
31. Miller GJ, Kotecha S, Wilkinson WH *et al.* (1988) Dietary and other characteristics relevant for coronary heart disease in men of Indian, West Indian and European descent in London. *Atherosclerosis* **70**, 63-72.
32. McKeigue PM, Marmot MG, Adelstein AM *et al.* (1985) Diet and risk factors for coronary heart disease in Asians in northwest London. *Lancet* **2**, 1086-1090.
33. Knight TM, Smith Z, Whittles A *et al.* (1992) Insulin resistance, diabetes, and risk markers for ischaemic heart disease in Asian men and non-Asian in Bradford. *Br Heart J* **67**, 343-350.
34. Sevak L, McKeigue PM, Marmot MG (1994) Relationship of hyperinsulinemia to dietary intake in south Asian and European men. *AmJ Clin Nutr* **59**, 1069-1074.
35. Deshmukh-Taskar P, Nicklas TA, Radcliffe JD *et al.* (2013) The relationship of breakfast skipping and type of breakfast consumed with overweight/obesity, abdominal obesity, other cardiometabolic risk factors and the metabolic syndrome in young adults. The National Health and Nutrition Examination Survey (NHANES): 1999-2006. *Public Health Nutr* **16**, 2073-2082.
36. Poslusna K, Ruprich J, de Vries JH *et al.* (2009) Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *Br J Nutr* **101 Suppl 2**, S73-85.

## Accepted manuscript

**Table 1. Dietary intake, anthropometry and cardiometabolic risk factors of participants, at 21-23y, in the Determinants of Adolescent, now young Adults, Social well-being and Health study**

|                                   | Whole cohort<br>(n=665) | White British<br>(n=107)            | Black Caribbean<br>(n=102)          | Black African<br>(n=132)            | Indian<br>(n=98)                    | Pakistani Bangladeshi<br>(n=111)    | Other<br>(n=115)     | <i>p</i> |
|-----------------------------------|-------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|----------------------|----------|
|                                   | Mean<br>(95% CI)        | Mean<br>(95% CI)                    | Mean<br>(95% CI)                    | Mean<br>(95% CI)                    | Mean<br>(95% CI)                    | Mean<br>(95% CI)                    | Mean<br>(95% CI)     |          |
| Male/Female (n)                   | 329/336                 | 52/55                               | 47/55                               | 58/74                               | 52/46                               | 62/49                               | 57/57                | 0.842    |
| Age (y)                           | 22.7<br>(22.7, 22.8)    | 22.7<br>(22.5, 22.8)                | 22.7<br>(22.5, 22.9)                | 22.6<br>(22.4, 22.9)                | 22.9<br>(22.7, 23.1)                | 22.6<br>(22.7, 23.1)                | 22.8<br>(22.6, 23.0) | 0.129    |
| Weight (kg)                       | 71.6<br>(70.0, 73.3)    | 72.5<br>(68.8, 76.2)                | 76.6 <sup>a</sup><br>(71.7, 81.5)   | 77.1 <sup>b</sup><br>(73.0, 81.2)   | 66.3 <sup>a,b</sup><br>(62.7, 69.9) | 68.0 <sup>a,b</sup><br>(64.1, 71.9) | 69.9<br>(66.5, 73.3) | <0.001   |
| Height (cm)                       | 170<br>(169, 171)       | 172 <sup>a</sup><br>(170, 175)      | 170<br>(168, 173)                   | 172 <sup>b</sup><br>(170, 174)      | 168 <sup>a,b</sup><br>(166, 171)    | 168 <sup>a,b</sup><br>(165, 170)    | 169<br>(167, 171)    | <0.001   |
| BMI (kg/m <sup>2</sup> )          | 24.7<br>(24.2, 25.2)    | 24.2 <sup>a,c</sup><br>(23.1, 25.4) | 26.2 <sup>a,b</sup><br>(24.8, 27.7) | 26.1 <sup>c,d</sup><br>(24.8, 27.4) | 23.4 <sup>b,d</sup><br>(22.2, 24.5) | 24.0<br>(22.9, 25.2)                | 24.4<br>(23.4, 25.5) | <0.001   |
| Overweight/obese (%) <sup>a</sup> | 39                      | 35                                  | 50                                  | 44                                  | 36                                  | 36                                  | 35                   | 0.063    |
| Waist (cm)                        | 82.2<br>(80.9, 83.5)    | 83.0<br>(80.2, 85.8)                | 83.6<br>(80.0, 87.3)                | 83.0<br>(79.9, 86.1)                | 80.8<br>(77.6, 84.1)                | 81.7<br>(79.0, 84.4)                | 81.2<br>(78.4, 84.1) | 0.878    |
| Waist:height ratio                | 0.48<br>(0.48, 0.49)    | 0.48<br>(0.46, 0.50)                | 0.49<br>(0.47, 0.51)                | 0.48<br>(0.46, 0.50)                | 0.48<br>(0.46, 0.50)                | 0.49<br>(0.47, 0.50)                | 0.48<br>(0.46, 0.50) | 0.429    |
| Total cholesterol (mmol/l)        | 4.35<br>(4.26, 4.43)    | 4.34<br>(4.17, 4.52)                | 4.34<br>(4.14, 4.54)                | 4.07 <sup>a</sup><br>(3.93, 4.22)   | 4.36<br>(4.21, 4.51)                | 4.53 <sup>a</sup><br>(4.35, 4.72)   | 4.33<br>(4.14, 4.52) | 0.007    |
| HDL cholesterol (mmol/l)          | 1.54<br>(1.50, 1.57)    | 1.56<br>(1.49, 1.64)                | 1.61 <sup>a</sup><br>(1.53, 1.70)   | 1.60 <sup>b</sup><br>(1.52, 1.67)   | 1.41 <sup>a,b</sup><br>(1.34, 1.48) | 1.47<br>(1.40, 1.54)                | 1.55<br>(1.46, 1.64) | 0.003    |
| HbA1c (mmol/mol)                  | 34<br>(34, 35)          | 34<br>(33, 35)                      | 35<br>(34, 36)                      | 35<br>(34, 36)                      | 35<br>(34, 36)                      | 34<br>(33, 35)                      | 34<br>(33, 35)       | 0.164    |
| Systolic blood pressure (mmHg)    | 114<br>(113, 115)       | 117<br>(114, 120)                   | 116<br>(113, 119)                   | 116<br>(113, 119)                   | 113<br>(110, 116)                   | 112<br>(109, 115)                   | 113<br>(110, 115)    | 0.090    |
| Diastolic blood pressure (mmHg)   | 72<br>(72, 73)          | 73<br>(71, 75)                      | 74<br>(72, 76)                      | 74<br>(72, 76)                      | 72<br>(70, 74)                      | 72<br>(70, 74)                      | 70<br>(68, 72)       | 0.073    |
| Energy intake                     | 2192                    | 2008                                | 2242                                | 1976                                | 2423                                | 2260                                | 2238                 | 0.255    |

## Accepted manuscript

|                                      |              |                  |                    |                    |                    |                   |              |        |
|--------------------------------------|--------------|------------------|--------------------|--------------------|--------------------|-------------------|--------------|--------|
| <b>(kcal/day)</b>                    | (2097, 2288) | (1811, 2205)     | (2018, 2466)       | (1766, 2186)       | (2164, 2682)       | (2001, 2520)      | (1982, 2494) |        |
| <b>Fat intake</b>                    | 38.0         | 37.9             | 38.5               | 36.5               | 35.6               | 41.1              | 38.6         | 0.168  |
| <b>(% of energy)</b>                 | (37.0, 39.0) | (35.5, 40.2)     | (36.4, 40.5)       | (33.7, 39.3)       | (33.0, 38.3)       | (38.9, 43.4)      | (36.1, 41.1) |        |
| <b>Saturated fat intake</b>          | 14.4         | 15.5             | 13.9               | 13.0 <sup>a</sup>  | 13.1 <sup>a</sup>  | 16.4 <sup>a</sup> | 14.5         | 0.031  |
| <b>(% of energy)</b>                 | (13.9, 15.0) | (14.2, 16.9)     | (12.7, 15.2)       | (11.7, 14.3)       | (11.8, 14.4)       | (14.9, 17.9)      | (13.3, 15.6) |        |
| <b>Carbohydrate intake</b>           | 47.4         | 46.1             | 48.1               | 47.9               | 50.4               | 45.6              | 46.5         | 0.170  |
| <b>(% of energy)</b>                 | (46.3, 48.5) | (43.8, 48.4)     | (45.4, 50.7)       | (44.8, 51.0)       | (47.8, 53.0)       | (42.9, 48.3)      | (43.8, 49.2) |        |
| <b>Protein intake</b>                | 16.4         | 16.0             | 15.9               | 17.1               | 15.7               | 16.6              | 16.8         | 0.953  |
| <b>(% of energy)</b>                 | (15.7, 17.1) | (14.4, 17.6)     | (14.2, 17.6)       | (14.8, 19.5)       | (13.9, 17.4)       | (15.1, 18.1)      | (15.6, 18.2) |        |
| <b>Sugar intake</b>                  | 17.1         | 16.6             | 19.5               | 18.8               | 16.2               | 14.5              | 17.1         | 0.151  |
| <b>(% of energy)</b>                 | (16.2, 18.0) | (14.5, 18.8)     | (17.0, 22.0)       | (16.1, 21.6)       | (14.2, 18.2)       | (12.7, 16.2)      | (15.0, 19.3) |        |
| <b>Fibre intake</b>                  | 8.1          | 9.3 <sup>a</sup> | 7.3 <sup>a,c</sup> | 7.7 <sup>b,c</sup> | 8.8 <sup>b,c</sup> | 7.3 <sup>c</sup>  | 8.0          | <0.001 |
| <b>(g/1000kcal)</b>                  | (7.7, 8.5)   | (8.1, 10.5)      | (6.3, 8.2)         | (6.8, 8.7)         | (7.6, 9.9)         | (6.3, 8.3)        | (6.9, 9.0)   |        |
| <b>Sodium intake</b>                 | 1285         | 1407             | 1320               | 1122               | 1192               | 1314              | 1345         | 0.082  |
| <b>(mg/1000kcal)</b>                 | (1231, 1340) | (1270, 1544)     | (1157, 1483)       | (1010, 1235)       | (1054, 1330)       | (1191, 1436)      | (1218, 1471) |        |
| <b>Sedentary time</b>                | 738          | 798              | 670                | 833                | 772                | 720               | 692          | 0.234  |
| <b>(mins/day)</b>                    | (683, 792)   | (650, 947)       | (573, 767)         | (6565, 1000)       | (654, 889)         | (560, 880)        | (575, 807)   |        |
| <b>Skipping breakfast (%)</b>        | 56           | 53               | 66                 | 64                 | 46                 | 51                | 54           | 0.118  |
| <b>Low fruit consumption (%)</b>     | 57           | 52               | 59                 | 65                 | 55                 | 56                | 52           | 0.373  |
| <b>Low vegetable consumption (%)</b> | 63           | 54               | 75                 | 73                 | 55                 | 73*               | 51*          | <0.001 |
| <b>Smoker (%)</b>                    | 24           | 47*              | 21                 | 11*                | 24                 | 24                | 24           | 0.001  |
| <b>Educated to degree level (%)</b>  | 47           | 40               | 34*                | 58*                | 57                 | 49                | 44           | 0.001  |

BMI, body mass index. Overweight/obese defined as BMI>25kg/m<sup>2</sup>. Low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; skipping breakfast defined as not eating breakfast every day of the week. Continuous variables are shown as mean (95% CI), categorical variables shown as n or %, as indicated in the variable label. Differences between ethnic groups in continuous variables were tested with one-way ANOVA with post-hoc analysis, the same superscript letters within a row denote significant differences between those ethnic groups; ethnic differences in frequency variables were tested with chi-square test with post-hoc analysis, differences are denoted with \*.

## Accepted manuscript

**Table 2. Influence of dietary behaviours at 11-13 years and at 21-23years on cardiometabolic risk factors at 21-23 years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.**

|   | BMI                    |          | Waist:height ratio      |          | Total cholesterol       |          | HDL-cholesterol         |          | HbA1c                  |          | Systolic blood pressure |          |
|---|------------------------|----------|-------------------------|----------|-------------------------|----------|-------------------------|----------|------------------------|----------|-------------------------|----------|
|   | Coef.<br>(95% CI)      | <i>p</i> | Coef.<br>(95% CI)       | <i>p</i> | Coef.<br>(95% CI)       | <i>p</i> | Coef.<br>(95% CI)       | <i>p</i> | Coef.<br>(95% CI)      | <i>p</i> | Coef.<br>(95% CI)       | <i>p</i> |
| <b>11-13y</b>                           |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| Skipping breakfast (No - Ref)           |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| Yes                                     | 1.41<br>(0.57, 2.26)   | <0.001   | 0.02<br>(0.01, 0.03)    | <0.001   | 0.15<br>(-0.01, 0.31)   | 0.063    | 0.01<br>(-0.06, 0.08)   | 0.725    | 0.43<br>(-0.53, 1.39)  | 0.379    | 0.74<br>(-0.89, 2.38)   | 0.370    |
| Not stated                              | 0.78<br>(-0.70, 2.26)  | 0.302    | 0.01<br>(-0.01, 0.03)   | 0.353    | 0.16<br>(-0.12, 0.45)   | 0.266    | 0.02<br>(-0.12, 0.15)   | 0.820    | 0.05<br>(-1.66, 1.75)  | 0.958    | 0.33<br>(-2.54, 3.19)   | 0.823    |
| Fruit ( $\geq 2$ serves/day - Ref)      |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| <2 serves/day                           | 0.68<br>(-0.17, 1.53)  | 0.115    | 0.01<br>(0.00, 0.03)    | 0.034    | -0.004<br>(-0.16, 0.16) | 0.958    | -0.05<br>(-0.12, 0.02)  | 0.197    | -0.16<br>(-1.11, 0.79) | 0.742    | 0.66<br>(-0.97, 2.30)   | 0.425    |
| Not stated                              | 0.65<br>(-0.58, 1.90)  | 0.299    | 0.01<br>(-0.01, 0.03)   | 0.225    | 0.03<br>(-0.20, 0.27)   | 0.779    | -0.003<br>(-0.11, 0.10) | 0.951    | -0.70<br>(-2.10, 0.70) | 0.325    | 0.21<br>(-2.17, 2.59)   | 0.865    |
| Vegetables ( $\geq 3$ serves/day - Ref) |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| <3 serves/day                           | -0.03<br>(-0.89, 0.82) | 0.941    | 0.002<br>(-0.01, 0.01)  | 0.723    | -0.17<br>(-0.33, -0.01) | 0.033    | -0.007<br>(-0.06, 0.08) | 0.838    | 0.18<br>(-0.75, 1.12)  | 0.702    | -0.003<br>(-1.64, 1.63) | 0.997    |
| Not stated                              | -1.70<br>(-1.00, 1.29) | 0.788    | -0.004<br>(-0.03, 0.02) | 0.709    | -0.11<br>(-0.40, 0.17)  | 0.427    | -0.007<br>(-0.12, 0.14) | 0.905    | -0.38<br>(-2.07, 1.30) | 0.656    | -0.79<br>(-3.66, 2.09)  | 0.591    |
| Fizzy drinks (No - Ref)                 |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| Yes                                     | 0.03<br>(-0.89, 0.84)  | 0.947    | 0.001<br>(-0.01, 0.01)  | 0.888    | 0.05<br>(-0.11, 0.21)   | 0.542    | -0.04<br>(-0.11, 0.03)  | 0.240    | -0.34<br>(-1.30, 0.62) | 0.488    | 1.77<br>(0.12, 3.43)    | 0.036    |
| Not stated                              | 0.06<br>(-1.44, 1.56)  | 0.938    | -0.002<br>(-0.02, 0.02) | 0.858    | 0.05<br>(-0.24, 0.34)   | 0.742    | -0.007<br>(-0.14, 0.12) | 0.920    | -0.53<br>(-2.25, 1.19) | 0.546    | 1.22<br>(-1.64, 4.09)   | 0.403    |
| <b>21-23y</b>                           |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| Skipping breakfast (No - Ref)           |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| Yes                                     | 1.05<br>(0.22, 1.89)   | 0.014    | 0.02<br>(0.01, 0.03)    | 0.002    | 0.22<br>(0.06, 0.37)    | 0.007    | 0.0002<br>(-0.07, 0.07) | 0.995    | 0.73<br>(-0.22, 1.67)  | 0.130    | 0.44<br>(-1.18, 2.05)   | 0.597    |
| Not stated                              | -1.32<br>(-4.44, 1.80) | 0.406    | -0.01<br>(-0.05, 0.04)  | 0.731    | 0.30<br>(-0.26, 0.85)   | 0.292    | 0.07<br>(-0.17, 0.32)   | 0.556    | 0.12<br>(-2.93, 3.18)  | 0.937    | -3.56<br>(-9.56, 2.44)  | 0.244    |
| Fruit ( $\geq 2$ serves/day - Ref)      |                        |          |                         |          |                         |          |                         |          |                        |          |                         |          |
| <2 serves/day                           | -0.26<br>(-1.11, 0.59) | 0.552    | 0.001<br>(-0.01, 0.01)  | 0.912    | -0.12<br>(-0.28, 0.04)  | 0.147    | -0.04<br>(-0.11, 0.03)  | 0.270    | -0.50<br>(-1.46, 0.46) | 0.305    | 0.26<br>(-1.37, 1.88)   | 0.758    |
| Not stated                              | -1.65<br>(-4.32, 1.02) | 0.225    | -0.008<br>(-0.05, 0.03) | 0.676    | 0.25<br>(-0.23, 0.74)   | 0.301    | 0.03<br>(-0.19, 0.24)   | 0.809    | -0.11<br>(-2.95, 2.74) | 0.940    | -0.39<br>(-5.51, 4.73)  | 0.881    |

## Accepted manuscript

| Vegetables ( $\geq 3$ serves/day - Ref) |                        |       |                        |       |                        |       |                        |       |                        |       |                        |       |
|---|------------------------|-------|------------------------|-------|------------------------|-------|------------------------|-------|------------------------|-------|------------------------|-------|
| <3 serves/day                           | -0.07<br>(-0.99, 0.84) | 0.873 | 0.001<br>(-0.01, 0.01) | 0.859 | -0.06<br>(-0.24, 0.11) | 0.456 | -0.02<br>(-0.09, 0.06) | 0.630 | 0.29<br>(-0.72, 1.30)  | 0.571 | -0.72<br>(-2.49, 1.04) | 0.420 |
| Not stated                              | -1.40<br>(-3.66, 0.85) | 0.221 | -0.01<br>(-0.04, 0.02) | 0.574 | -0.12<br>(-0.51, 0.28) | 0.559 | 0.08<br>(-0.10, 0.25)  | 0.385 | -0.17<br>(-2.49, 2.15) | 0.884 | -3.27<br>(-7.58, 1.05) | 0.137 |

BMI, body mass index. Skipping breakfast defined as eating breakfast less than 5 days per week; low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; fizzy drink consumption defined as consuming fizzy drinks every day or most days. Multivariate linear regression models for BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c and systolic blood pressure (separate models); coefficients were estimated with linear regression, adjusted for gender and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), smoking (current smoker/non-smoker), energy intake and saturated fat intake at 21-23 years. N=664, 635, 499, 484, 481, and 664, respectively, for linear regression models with BMI, waist:height ratio, total cholesterol, HDL-cholesterol, HbA1c, and systolic blood pressure, respectively, as the outcome.

## Accepted manuscript

**Table 3. Influence of dietary behaviours at 11-13 years and at 21-23years on nutritional composition of the diet at 21-23years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.**

|   | Energy intake (kcal)         |          | Fat (% of energy)      |          | Saturated fat (% of energy) |          | Sugar (% of energy)    |          | Fibre (g/1000kcal)      |          | Sodium (mg/1000kcal)         |          |
|---|------------------------------|----------|------------------------|----------|-----------------------------|----------|------------------------|----------|-------------------------|----------|------------------------------|----------|
|   | Coef. (95% CI)               | <i>p</i> | Coef. (95% CI)         | <i>p</i> | Coef. (95% CI)              | <i>p</i> | Coef. (95% CI)         | <i>p</i> | Coef. (95% CI)          | <i>p</i> | Coef. (95% CI)               | <i>p</i> |
| <b>11-13y</b>                           |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| Skipping breakfast (No - Ref)           |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| Yes                                     | -116.25<br>(-263.65, 31.15)  | 0.122    | -0.19<br>(-1.86, 1.48) | 0.826    | 0.42<br>(-0.48, 1.31)       | 0.361    | 0.11<br>(-1.43, 1.65)  | 0.889    | -0.81<br>(-1.49, -0.12) | 0.022    | -58.83<br>(-184.66, 67.00)   | 0.359    |
| Not stated                              | -176.37<br>(-435.51, 82.77)  | 0.182    | 0.14<br>(-2.80, 3.08)  | 0.927    | 0.66<br>(-0.92, 2.24)       | 0.411    | 2.13<br>(-0.58, 4.85)  | 0.123    | -0.74<br>(-1.95, 0.47)  | 0.230    | 50.38<br>(-170.84, 271.61)   | 0.655    |
| Fruit ( $\geq 2$ serves/day - Ref)      |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| <2 serves/day                           | 142.89<br>(-4.17, 289.95)    | 0.057    | -0.57<br>(-2.24, 1.10) | 0.503    | -0.44<br>(-1.33, 0.46)      | 0.339    | -0.14<br>(-1.68, 1.40) | 0.859    | -0.36<br>(-1.04, 0.33)  | 0.312    | -5.49<br>(-131.17, 120.19)   | 0.932    |
| Not stated                              | 23.48<br>(-191.86, 238.83)   | 0.830    | -0.84<br>(-3.29, 1.60) | 0.498    | -0.65<br>(-1.96, 0.66)      | 0.330    | 1.34<br>(-0.92, 3.59)  | 0.244    | 0.16<br>(-0.85, 1.17)   | 0.752    | 68.67<br>(-115.37, 252.70)   | 0.464    |
| Vegetables ( $\geq 3$ serves/day - Ref) |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| <3 serves/day                           | -48.49<br>(-197.09, 100.11)  | 0.522    | -1.15<br>(-2.83, 0.53) | 0.178    | -0.60<br>(-1.50, 0.30)      | 0.193    | 1.32<br>(-0.23, 2.86)  | 0.095    | -0.31<br>(-1.00, 0.39)  | 0.383    | -27.77<br>(-154.45, 98.91)   | 0.667    |
| Not stated                              | -108.10<br>(-368.68, 152.47) | 0.416    | -1.23<br>(-4.17, 1.72) | 0.414    | -0.07<br>(-1.65, 1.51)      | 0.930    | 3.56<br>(0.85, 6.27)   | 0.010    | -0.10<br>(-1.32, 1.12)  | 0.869    | 47.31<br>(-174.83, 269.45)   | 0.676    |
| Fizzy drinks (No - Ref)                 |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| Yes                                     | 34.82<br>(-115.70, 185.35)   | 0.650    | 0.04<br>(-1.66, 1.74)  | 0.964    | 0.34<br>(-0.58, 1.25)       | 0.469    | -0.08<br>(-1.65, 1.48) | 0.915    | -0.72<br>(-1.42, -0.02) | 0.045    | -51.30<br>(-179.49, 76.89)   | 0.432    |
| Not stated                              | -47.59<br>(-308.79, 213.60)  | 0.721    | -0.27<br>(-3.23, 2.69) | 0.857    | 0.66<br>(-0.93, 2.25)       | 0.414    | 3.15<br>(0.44, 5.87)   | 0.023    | -0.66<br>(-1.88, 0.56)  | 0.287    | 70.21<br>(-152.23, 292.64)   | 0.536    |
| <b>21-23y</b>                           |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| Skipping breakfast (No - Ref)           |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| Yes                                     | -68.16<br>(-214.82, 78.49)   | 0.362    | 1.10<br>(-0.56, 2.75)  | 0.195    | 0.26<br>(-0.63, 1.15)       | 0.565    | -0.49<br>(-2.02, 1.04) | 0.530    | -1.33<br>(-2.01, -0.65) | <0.001   | -35.13<br>(-160.14, 89.88)   | 0.581    |
| Not stated                              | -144.67<br>(-689.91, 400.57) | 0.602    | 1.70<br>(-4.46, 7.87)  | 0.587    | -0.99<br>(-4.30, 2.33)      | 0.558    | -2.17<br>(-7.87, 3.54) | 0.456    | 0.00<br>(-2.52, 2.52)   | 0.998    | -199.83<br>(-664.60, 264.93) | 0.399    |
| Fruit ( $\geq 2$ serves/day - Ref)      |                              |          |                        |          |                             |          |                        |          |                         |          |                              |          |
| <2 serves/day                           | 27.47<br>(-119.98, 174.91)   | 0.715    | 0.70<br>(-0.97, 2.37)  | 0.411    | 0.89<br>(-0.01, 1.78)       | 0.052    | -0.47<br>(-2.01, 1.07) | 0.547    | -1.22<br>(-1.90, -0.54) | <0.001   | -29.32<br>(-154.98, 96.35)   | 0.647    |
| Not stated                              | -7.71<br>(-473.24, 457.83)   | 0.974    | 1.19<br>(-4.08, 6.45)  | 0.658    | 0.14<br>(-2.68, 2.96)       | 0.921    | -3.14<br>(-8.00, 1.72) | 0.205    | 0.67<br>(-1.48, 2.82)   | 0.542    | -33.96<br>(-430.73, 362.82)  | 0.867    |

## Accepted manuscript

|   |                             |       |                       |       |                       |       |                        |       |                         |       |                             |       |
|---|-----------------------------|-------|-----------------------|-------|-----------------------|-------|------------------------|-------|-------------------------|-------|-----------------------------|-------|
| Vegetables ( $\geq 3$ serves/day - Ref) |                             |       |                       |       |                       |       |                        |       |                         |       |                             |       |
| <3 serves/day                           | -27.94<br>(-187.19, 131.32) | 0.731 | 1.50<br>(-0.29, 3.30) | 0.100 | 1.06<br>(0.09, 2.02)  | 0.032 | 0.11<br>(-1.55, 1.78)  | 0.896 | -1.30<br>(-2.04, -0.56) | 0.001 | -101.31<br>(-236.80, 34.18) | 0.143 |
| Not stated                              | 26.10<br>(-366.17, 418.37)  | 0.896 | 4.01<br>(-0.42, 8.43) | 0.076 | 1.27<br>(-1.10, 3.65) | 0.294 | -1.83<br>(-5.93, 2.27) | 0.380 | -0.76<br>(-2.58, 1.05)  | 0.410 | 18.78<br>(-314.96, 352.51)  | 0.912 |

Skipping breakfast defined as eating breakfast less than 5 days per week; low fruit consumption defined as less than 2 portions per day; low vegetable consumption defined as less than 3 portions per day; fizzy drink consumption defined as consuming fizzy drinks every day or most days. Multivariate linear regression models for energy intake, % energy from fat, % energy from saturated fat, % energy from sugar, fibre intake, sodium intake (separate models); coefficients were estimated with linear regression, adjusted for gender, and ethnicity at 11-13 years, and additionally adjusted for educational attainment (attainment/non-attainment of higher education degree), physical activity (based on tertiles of sitting/sedentary time), and smoking (current smoker/non-smoker) at 21-23 years. N=628 for linear regression models for all six nutritional composition variables as the outcome.



# Accepted manuscript

## FIGURE LEGENDS

**Figure 1.** Rates of (a) regularly skipping breakfast, (b) low daily fruit and (c) low daily vegetable consumption according to ethnicity at 11-13 years and at 21-23 years. The Determinants of Adolescent, now young Adults, Social well-being and Health study.



## Accepted manuscript



## Accepted manuscript

