

---

Citation:

Sheehy, K and Noureen, A and Khaliq, A and Dhingra, K and Husain, N and Pontin, E and Cawley, R and Taylor, P (2019) An examination of the relationship between shame, guilt and self-harm: A systematic review and meta-analysis. *Clinical Psychology Review*, 73. ISSN 0272-7358 DOI: <https://doi.org/10.1016/j.cpr.2019.101779>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/6166/>

Document Version:

Article (Accepted Version)

---

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

## RUNNING HEAD: SHAME, GUILT, AND SELF-HARM

### **An examination of the relationship between shame, guilt and self-harm:**

#### **A systematic review and meta-analysis**

Kate Sheehy<sup>1</sup> (Kate.Sheehy@merseycare.nhs.uk)

Amna Noureen<sup>2</sup> (Amna.Naureen@pill.org.pk)

Ayesha Khaliq<sup>2</sup> (Ayesha.Khaliq@pill.org.pk)

Katie Dhingra<sup>3</sup> (K.Dhingra@outlook.com)

Nusrat Husain<sup>4</sup> (Nusrat.Husain@manchester.ac.uk)

Eleanor E. Pontin<sup>1</sup> (Ellie.Pontin@nhs.net)

Rosanne Cawley<sup>1</sup> (Rosanne.Cawley@mft.nhs.uk)

Peter J. Taylor<sup>4\*</sup> (Peter.Taylor-2@manchester.ac.uk)

<sup>1</sup> Institute of Psychology, Health and Society, University of Liverpool

<sup>2</sup> Pakistan Institute of Living & Learning, Karachi, Pakistan

<sup>3</sup> School of Social Sciences, Leeds Beckett University, England, LS1 3HE

<sup>4</sup> Division of Psychology & Mental Health, School of Health Sciences, University of Manchester, Manchester Academic Health Sciences Centre, Manchester, United Kingdom

\*Corresponding author

Correspondence Address: 2nd floor, Zochonis building, Brunswick Park, University of Manchester, Manchester, M13 9PL, England.

Tel: +44(0)1613060425

Email: [Peter.Taylor-2@Manchester.ac.uk](mailto:Peter.Taylor-2@Manchester.ac.uk)

### **Role of Funding Source**

Funding supporting this study was provided by *Medical Research Council* (MR/N006062/1).

The MRC had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

### **Contributors**

Dr Sheehy was involved in conceptualisation, design and planning of the review (including protocol development), literature searches and screening, data extraction, synthesis and evaluation, and write-up. Dr Dhingra, Ms Noureen and Ms Khaliq were involved in screening, data extraction, quality checking, evaluation and write-up. Dr Pontin and Prof. Husain were involved in supervision, planning, and write-up. Dr Cawley was involved in conceptualisation, planning and evaluation. Dr Taylor was involved in project management, supervision, conceptualisation, design and planning of the review, literature searches and screening, data extraction, synthesis and evaluation, and write-up.

### **Conflicts of Interest**

None

### **Acknowledgments**

We would like to acknowledge those authors who supported this review by providing additional information and data. These include Craig Bryan, David Kealy, Hilary Weingarden, Nicolas Rüesch, and Maria Wiklander.

## Abstract

Self-harm is a major public health concern associated with suicide risk and significant psychological distress. Theories suggest that aversive emotional states are an important process that drives self-harm. Shame and guilt may, in particular, be important emotions in self-harm. This review therefore sought to provide a systematic review and meta-analysis of the relationship between shame, guilt, and self-harm. A systematic search of electronic databases (PsycINFO; Medline; CINAHL Plus; Web of Science and ProQuest) was undertaken to identify studies measuring shame, guilt and self-harm (including suicidal and non-suicidal behaviour). Meta-analysis was undertaken where papers focused on the same subtype of shame or guilt and shared a common outcome. Thirty studies were identified for inclusion. Most forms of shame were associated with non-suicidal self-injury (NSSI), but research was sparse concerning suicidal behaviour. Fewer studies examined guilt and findings were more varied. Methodological issues included a paucity of longitudinal designs and lack of justification for sample sizes. Results of this review support the link between shame and self-harm, particularly NSSI. The direction of this relationship is yet to be established. Clinically, consideration should be given to the role of shame amongst individuals who present with NSSI. This review was pre-registered on PROSPERO (CRD42017056165).

**Keywords:** shame, guilt, self-harm, non-suicidal self-injury (NSSI), suicide, systematic review

### **Highlights**

- Shame has a positive association with self-harm
- Guilt proneness does not appear to be related to self-harm
- However, exposure to state guilt is greater in those with a history of self-harm
- Shame should be considered in psychological assessments with those who self-harm
- Longitudinal studies in this area are lacking

## Introduction

Self-harm poses a significant public health concern worldwide, because of its high prevalence (Muehlenkamp, Claes, Havertape, & Plener, 2012; Swannell, Martin, Page, Hasking, & St John, 2014), and association with subsequent suicide risk (Hawton et al., 2015; Ribeiro et al., 2016). It is also often indicative of psychological distress and additional need (Goldman-Mellor et al., 2014), and reduced life expectancy from any cause (approximately 26 years of life lost; Bergen et al., 2012). Self-harm refers to the deliberate destruction or damage to one's own body tissue, irrespective of suicidal intent, and can be applied to a range of behaviours including overdose, cutting, burning, and self-battery (National Collaborating Centre for Mental Health, 2004; Royal College of Psychiatrists, 2010). Self-harm therefore encompasses suicidal behaviour (i.e. suicide attempts) and non-suicidal self-injury (NSSI; Klonsky, 2011), as well as behaviours where the level of suicidal intent is unclear or ambiguous. Recent reviews suggest that one of the most commonly reported reasons for self-harm is around coping with or regulating difficult emotional states (Edmondson, Brennan, & House, 2016; Taylor et al., 2018). For example, this function is endorsed by 71% (95% CI: 63-78%) of those who engage in NSSI (Taylor et al., 2018). Thus, mechanisms involving exposure to and regulation of emotional states appear key to understanding self-harm (Nock, 2009). A better understanding of these mechanisms can help inform the development and adaptation of interventions for those who struggle with self-harm (Muehlenkamp, 2006). Certain emotions appear especially important in understanding self-harm (Klonsky, 2009). The current review focuses on two such emotions, shame and guilt.

Shame and guilt have been described as self-conscious, 'moral' emotions, which arise in response to an evaluation of the self (Tangney & Dearing, 2002; Tangney, Stuewig, & Mashek, 2007). Although routinely considered in tandem, shame and guilt are thought to represent distinct, yet overlapping, emotional experiences (Tangney et al., 2007). Current

thinking regarding this distinction points to a differential focus on the self, versus one's behaviour. At its core, shame can be seen as a cognitive affective construct, comprising negative judgements of the self (Chou et al., 2018). These judgements are global, undesirable, and characterised by an evaluation of the self as inherently flawed, inadequate or bad (Blythin et al., 2018; Carden, Saini, Seddon, Watkins, & James Taylor, 2018; Gilbert & Procter, 2006). By contrast, guilt is concerned with one's behaviour, and the negative evaluation of this (Tangney & Dearing, 2002; Tangney et al., 2007). Hence, the object of focus is something done by the individual that is perceived as bad or wrong, rather than the individual themselves. As a result, the phenomenological experiences of guilt and shame are said to diverge significantly (Lewis, 1971).

Traditionally, the conceptualisation of shame has centred upon the individual's perception of themselves. However, some researchers have distinguished between this and an individual's representation of how they are perceived by others (specifically the individual's perception of being negatively judged by others), referred to as 'external' shame (Gilbert, 1997, 1998). In addition, shame may be thought to arise in relation to different aspects of the self, such as one's character, behaviour, or body (Andrews, Qian, & Valentine, 2002). As a result, a range of psychometric measures have been developed and used to assess these various components of shame. No such distinctions have been made in relation to guilt as far as we are aware. Furthermore, whilst it is acknowledged that shame and guilt may occur in relation to specific incidents or events, it is now also recognised that some individuals have a greater tendency, or proneness, to experience feelings of shame or guilt across a range of situations (Tangney, 1990).

Both shame and guilt may be experienced as unwanted or aversive emotional states. However, literature suggests that shame may be particularly pernicious due to its close ties with an individual's sense of self (Lewis, 1971). Indeed, shame is closely linked with various



psychological difficulties including depression, psychosis, Post-Traumatic Stress Disorder (PTSD), and eating disorders (Blythin et al., 2018; Carden et al., 2018; Kim, Thibodeau, & Jorgensen, 2011; Pugh, Taylor, & Berry, 2015). Across the available research, there is evidence that shame is more robustly associated with psychological difficulties and that when adjusting for overlapping shame, guilt at times is no longer associated with mental health difficulties (Blythin et al., 2018; Kim et al., 2011; Pugh et al., 2015). Nonetheless, guilt too may be experienced as painful, and may give rise to feelings of regret or remorse (Pugh et al., 2015; Tangney, Miller, Flicker, & Barlow, 1996). Whilst guilt may lead an individual to engage in reparative action to address perceived problematic behaviour (Tangney & Dearing, 2002), responses to shame are typically less adaptive and include rumination (Cheung, Gilbert, & Iron, 2004), submission (Gilbert, Pehl, & Allan, 1994), avoidance (Schoenleber & Berenbaum, 2012), and attempts to conceal oneself or one's perceived faults (Tangney et al., 1996). In light of the available literature, it is hypothesised that shame will show a stronger relationship with self-harm than guilt.

Theoretical models have typically focused either on suicidal behaviour or NSSI, rather than the broader construct of self-harm. Theoretical models of NSSI explicitly suggest that NSSI is maintained by negative reinforcement, characterised by escape from unpleasant internal states, including emotions like shame and guilt (Chapman, Gratz, & Brown, 2006; Hasking, Whitlock, Voon, & Rose, 2017; Nock, 2009). Some theorists have developed these ideas further by implying a specific role of shame or guilt in the aetiology and maintenance of NSSI. For example, some NSSI may arise out of beliefs about the self as deserving of punishment (Glassman, Weierich, Hooley, Deliberto, & Nock, 2007; Nock, 2009), which could be a consequence of strong feelings of shame. Schoenleber and Berenbaum (2012), for example, propose that individuals may engage in NSSI as a means of managing feelings of shame. Theories of suicidal behaviour have also posited that strong negative emotions may

drive suicidal behaviour (Baumeister, 1990; Williams, 1997). The Interpersonal Theory of suicide (Joiner, 2005; Van Orden et al., 2010) suggests feelings of burdensomeness are key to the desire for suicide, and that feelings of self-hate are a facet of this construct (Van Orden, Cukrowicz, Witte, & Joiner, 2012; Van Orden et al., 2010). Since shame is the emotion perhaps most synonymous with self-hate it may therefore be relevant in driving suicidal urges. Shame may be part of the mechanism explaining the increased risk of self-harm in some marginalized groups, such as those who are lesbian, gay or bisexual (Taylor, Dhingra, Dickson, & McDermott, 2018), or those belonging to alternative subcultures (Hughes, Knowles, Dhingra, Nicholson, & Taylor, 2018). For example, experiences of rejection associated with belonging to a marginalized group (e.g. being lesbian, gay, bisexual or transgender) are associated with self-harm risk (Cawley, Pontin, Touhey, Sheehy, & Taylor, 2019).

The current research aims to provide a systematic review and meta-analysis of the available literature pertaining to self-harm and its relationships with shame and guilt. We will appraise the weight of evidence concerning the relationship between these constructs, and via the meta-analysis quantify the size of these associations. It has been noted that shame and guilt overlap with one another, and also with depressive symptoms. In addition to focussing on bivariate associations, we also review associations whilst adjusting for guilt (when the effect involves shame) or shame (when the effect involves guilt) and depression.

## **Method**

### **Protocol Registration**

A systematic review protocol was developed and pre-registered online with PROSPERO (CRD42017056165). This review followed the PRISMA reporting guidelines (Moher, Liberati, Tetzlaff, Altman, & The, 2009). Departures from protocol include the addition of meta-analyses, which were included to provide a further summary of associations,

the exclusion of ETHOS as a database (as it is limited to UK dissertations), the expansion of the review team, and the secondary outcome concerning help-seeking being dropped in light of recent existing reviews in this area (Rowe et al., 2014).

### **Search Strategy**

First, scoping searches were undertaken to aid the identification of relevant search terms. Following this, four databases were searched (PsycINFO; Medline; CINAHL Plus; Web of Science) to identify relevant published studies (from earliest records until March 2017). These searches were later updated to December 2018. The following search terms were used related to: (a) self-harm: *NSSI* OR *suicid\** OR *self-harm* OR *self-injur\** OR *self-mutilation* OR *overdose* OR *DSH* OR *parasuicid\**; and (b) shame or guilt: *ashamed* OR *shame\** OR *guilt\** OR *self-blame* OR *self-disgust*. Search terms for the two groups were combined using the Boolean operator “AND”. The thesis and dissertation database ProQuest was also searched to identify relevant studies in the grey literature. First, titles and abstracts were screened independently by a single researcher. Following this, the full texts of the remaining articles were read to determine eligibility for inclusion. This was carried out independently by two researchers, with discrepancies addressed through discussion with a third author. Following the identification of included studies, reference lists of these papers were hand searched and corresponding authors were also emailed to identify any further potentially eligible studies.

### **Inclusion and Exclusion Criteria**

Studies included in the review were required to meet the following inclusion criteria: i) quantitative research studies, ii) comprising original research, iii) written in English, iv) measuring shame and/or guilt, v) measuring self-harm history or frequency (including NSSI and suicidal behaviour), vi) providing adequate information to estimate associations between variables. Studies using measures that conflated the constructs of shame and guilt were

excluded. For example, the Positive and Negative Affect scale has a guilt subscale that has items referring to feeling “ashamed” (Watson & Clark, 1994). Similarly, we excluded measures of shame focused on specific behaviours due to the overlap with guilt. Studies that assessed shame or guilt using a single item measure were also not included in the review, including studies in which guilt was assessed solely as part of a depression or mania measure, such as the Beck Depression Inventory (Beck, Steer, & Brown, 1996), as such measures may not provide a valid and accurate measure of the key constructs. Measures of shame or guilt related to experiences of trauma or grief or consisting of psychotic symptoms (i.e. delusional guilt) were also excluded, since these arguably have a distinct phenomenology relative to shame and guilt more generally.

### **Risk of Bias Assessment**

Studies included in the review were evaluated for risk of bias using an adapted version of the Agency for Healthcare Research and Quality (AHRQ; Williams, Plassman, Burke, Holsinger, & Benjamin, 2010) risk of bias tool. The AHRQ has previously been adapted for use in systematic reviews of self-injurious behaviour and associated constructs (Hughes et al., 2018; Taylor, Hutton, & Wood, 2015). The tool assesses risk of bias over eleven domains, including the validity of measures used, unbiased selection of participants, and appropriateness of analytic methods. Ratings were made independently by two research team members for each study, and then compared, with discrepancies resolved through discussion. These ratings were then used to identify common risks of bias across the literature as well as areas of strength.

### **Data Extraction**

Data extraction was undertaken independently by two research team members for each study, and then compared, with discrepancies resolved through discussion. Extracted data included study details (author, date, study location), study design information (type of design,

number of groups, recruitment method), participant characteristics (target sample, age, gender), measures used, and results of analyses.

### **Meta-Analytic Calculations**

Bivariate associations between shame, guilt and self-harm were grouped according to the emotion type (shame, guilt) and subtype (e.g. bodily shame, external shame, shame proneness) and outcome (NSSI, suicide attempt or self-harm not otherwise specified). Where two or more effects were grouped together these were aggregated via meta-analysis. This approach allowed the effects for different subtypes of shame or guilt to be compared, but it also meant that meta-analyses often had few included studies. We therefore also grouped studies by emotion type (shame or guilt), irrespective of subtype, and outcome. This allowed for effects from a larger set of studies to be combined by including studies investigating different subtypes of shame or guilt together in the same meta-analysis.

A random-effects model was adopted for all meta-analyses to accommodate the expected heterogeneity between studies in terms of sample, design, and measurement. The Dersimonian and Laird (DL; 1986) inverse variance estimator, within STATA 14 (StataCorp, 2015) was initially used. Whilst the DL estimator is commonly used, it has been noted that Restricted Maximum Likelihood (REML) estimator may better estimate between-study variance within random effects meta-analysis of continuous outcomes (Veroniki et al., 2014). We therefore repeated all meta-analyses using the REML estimator as well using the METAAN package (Kontopantelis & Reeves, 2010).

Within samples associations were captured with the correlation coefficient,  $r$ , whilst group differences were captured using cohen's  $d$ . Where necessary, effects were converted between effect size metrics following the steps outlined by Borenstein and colleagues (2009). For the meta-analyses undertaken on higher-order groupings of studies, a single effect size was taken from each study. Where studies included multiple different scales of the same emotion

(e.g. shame) these effects were first combined (following Borenstein et al., 2008). For one study this was not possible due to lack of information about the scales (Rusch et al., 2007) and so the effect size associated with the most commonly used scale (the TOSCA-3) was used. The  $I^2$  statistic was used to determine the impact of heterogeneity between studies (Higgins & Thompson, 2002).

## Results

### Study Characteristics

Thirty eligible papers were identified for inclusion in the review. A flow diagram of the screening process from identification through to inclusion is presented in Figure 1. Study characteristics are summarised in Table 1. All but two studies provided cross-sectional data on the relationship between shame, guilt and self-harm. All studies came from Western countries, most commonly the US ( $k = 18$ ) and UK ( $k = 6$ ). A large number of studies came from MSc or doctoral dissertations ( $k = 14$ ), though the majority derived from peer-reviewed journals ( $k = 16$ ). A single study was unpublished, with data made available by the author.

A broad variety of measures capturing distinct subtypes of shame and guilt were assessed across the included studies. The most common was shame or guilt proneness, typically assessed using the TOSCA (Tangney, 1990) or later versions of this measure ( $k = 11$ ). Body-related shame was the second most commonly assessed subtype of shame, usually measured using the Experiences of Shame Scale ( $k = 4$ ; ESS; Andrews et al., 2002). A number of studies also assessed exposure to general feelings of shame or guilt (not linked to a particular aspect of the self) using the Personal Feelings Questionnaire ( $k = 6$ ; PFQ2; Harder & Zalma, 1990). Within this review we refer to this as ‘state shame’ and ‘state guilt’, to distinguish it from other forms. This broad range of emotion subtypes meant that few studies were identified focusing on any one subtype, and thus limited the number of studies contributing to any one meta-analysis.

FIGURE 1 ABOUT HERE

TABLE 1 ABOUT HERE

### **Risk of bias**

The assessment of risk of bias is presented in Table 1. Overall, risk of bias was relatively low with regards to the data that were the focus of this review. Notably, for unpublished data sets the information was not always available to ascertain the risk of bias associated with these data. The most common methodological problems related to justification of sample size, the use of heavily self-selecting samples (e.g. participants responding to flyers or online advertisements), blinding of researchers, and measurement of self-harm. Only four studies justified their sample size in terms of power calculations. This may mean that analyses were underpowered in some cases, leading to inflated Type II error rates. Attempts at blinding researchers or interviewers to participants' status were rarely undertaken, which may have introduced rater bias and expectancy effects. Although most studies still employed widely used and validated tools to assess self-harm status, around a third used single-item (sometimes unvalidated) self-report measures, which may have led to misclassification, thus increasing the likelihood of statistical decision errors. However, there is support for the predictive and convergent validity of the Beck Scale for Suicide Ideation (BSS; Beck & Steer, 1991) suicidal behaviour item, which was often used.

In terms of methodological strengths, five of the six studies involving group comparisons attempted to match groups on key socio-demographic variables (e.g. age, gender, ethnicity, socio-economic status). Hence confounding variables are unlikely to have biased group comparisons. Furthermore, all but one study used a validated method for ascertaining the clinical status or participant group, and most studies used valid and reliable measures to rate shame and/or guilt. Missing data also appeared minimal (i.e. < 20%) for a

large proportion of studies, and in cases where missing data was apparent, appropriate details were provided in terms of how this was managed (e.g. use of imputation strategies to minimize bias). Finally, the analytic techniques adopted were appropriate in the large majority of studies.

## TABLE 2 ABOUT HERE

### **Association between Shame, Guilt and NSSI**

Table 3 reports the bivariate association between shame, guilt, and self-harm, grouped in terms of emotion subtype and outcome. Where data on two or more comparable associations were identified, a random-effects meta-analysis was undertaken to produce aggregate effect sizes. Bivariate effects for individual studies are reported in Supplementary Table 1. When meta-analyses were repeated using REML rather than DL estimation results were very similar (see Supplementary Table 2).

Individuals with a history of NSSI reported greater levels of shame proneness, characterological shame, and external shame, with moderate to large effect sizes. No significant difference was apparent for body-related shame ( $k = 5$  studies). This lack of difference was largely informed by the study by Duggan and colleagues in school students (2015), whose data suggests little overall difference in body shame at time 2 between those with and without a history of NSSI. However, Duggan and colleagues (2015) did report significant longitudinal effects, discussed below. There was also an unusually large effect size for a single study looking at body shame, linked to unusually small standard deviations (Nelson & Muehlenkamp, 2012). One study focused on external shame in a high secure inpatient sample noted that self-reported external shame was lower than is typically seen in the general population, though it is unclear whether this reflects a response bias or a characteristic of the population (Mallindine, 2002). When all studies investigating the relationship between shame (irrespective of subtype) and NSSI history were included



together in a meta-analysis ( $k = 10$ ), a large association was identified,  $d = 1.09$  (0.17, 2.01),  $I^2 = 98\%$ , but this reduced to a more moderate effect when the one study with unusually small standard deviations (Nelson & Muehlenkamp, 2012) was excluded,  $d = 0.47$  (0.17, 0.78),  $I^2 = 82\%$ , and with REML estimation,  $r = .48$  (.20, .76).

There were also small to moderate positive associations between NSSI frequency and shame proneness, external shame and characterological shame. However, again, there was no significant relationship with body shame based on a meta-analysis of two studies (Etzel, 2004; Pritchard, 2009). When all studies investigating the relationship between shame (irrespective of subtype) and NSSI frequency were included together in a meta-analysis ( $k = 7$ ), a moderate association was identified,  $r = .24$  (.06, .40),  $I^2 = 88\%$ , and with REML estimation,  $r = .24$  (.07, .39). In summary, there is evidence that shame (with the possible exception of body shame) is elevated in those with a history of NSSI, and to a lesser extent, associated with the frequency of NSSI. This was apparent across a range of populations including university and high school students, survivors of sexual abuse and domestic violence, inpatients, adults with experiences of NSSI, and people in prison. However, most studies relied on small samples ( $k = 7$  with  $n < 100$ ) and many shame subtypes were investigated by only one or two studies, making these findings preliminary.

In contrast to shame, only two studies of undergraduate students examined the relationship between guilt and NSSI. Guilt proneness was not related to either NSSI history or frequency (Schoenleber, 2013; VanDerhei, Rojahn, Stuewig, & McKnight, 2014).

#### TABLE 3 ABOUT HERE

### **Association between Shame, Guilt and Suicidal Behaviour**

Shame proneness was elevated in participants with a history of suicide attempts compared to those without, but not state shame (see Table 3). These two meta-analyses demonstrated high inconsistency, suggesting that effect sizes are moderated by other factors.

The number of studies contributing to these meta-analyses were too low to warrant statistical testing of moderating variables (Higgins & Green, 2009), but it is notable that smaller effects were apparent for one study focused on women with a borderline personality disorder (BPD) diagnosis (Rusch et al., 2007). It is possible that because experiences of shame are already elevated in those diagnosed with BPD (Rizvi, Brown, Bohus, & Linehan, 2011) the subsequent association with suicidal behaviour is attenuated. Two studies also found associations between shame and suicide attempt frequency, reporting small but significant positive correlations with shame related to character, body, performance, appearance and relationships (McLeod, 2002; Rutherford, 2015). When all studies investigating the relationship between shame (irrespective of subtype) and suicidal behaviour were included together in a meta-analysis ( $k = 8$ ), a moderate association was identified,  $d = 0.49$  (0.27, 0.70),  $I^2 = 70\%$ , and with REML estimation,  $d = 0.48$  (0.28, 0.69). Overall, there is evidence of a small relationship between some forms of shame and suicidal behaviour, but this evidence is limited, and further confirmation is required.

State guilt, but not guilt proneness, was also elevated in those with a history of suicide attempts compared to those without (See Table 3). When all studies investigating the relationship between guilt (irrespective of subtype) and suicidal behaviour were included together in a meta-analysis ( $k = 8$ ), a small association was identified,  $d = 0.29$  (0.06, 0.51),  $I^2 = 67\%$ , and with REML estimation,  $d = .32$  (.06, .60).

### **Association between Shame, Guilt and Self-Harm**

A subset of three studies measured self-harm, rather than NSSI or suicidal behaviour more specifically (Kealy, 2019; Lamb, 2004; Milligan & Andrews, 2005). Two out of three studies reported significantly greater levels of shame in those with a history of self-harm than those without (see Table 3). Shame was not significantly correlated with self-harm frequency in one study of psychiatric outpatients (Lamb, 2004), and the direction of the trend was

actually negative (greater guilt related to less frequent self-harm), though the sample was very small ( $n = 20$ ) increasing the risk of unusual and unrepresentative results. Unpublished data also indicated elevated state guilt in psychiatric outpatients with a history of self-harm (Kealy, 2019).

### **Adjusted Associations**

The association between shame and NSSI (frequency or history) remained significant even after adjusting for guilt in most analyses (5 out of 6). In a sample of undergraduate students, appearance-related shame emerged as the only significant correlate of suicide attempt history (OR = 1.04) when adjusting for shame and guilt proneness, relationships, and performance related shame (McLeod, 2002). However, another study of individuals with a history of suicidal thoughts found no significant relationship between shame proneness and suicidal behaviour when adjusting for guilt proneness ( $r = .20$ ; Izadi, 2014). It is unclear if the form of correlation used in this study is suitable for a binary outcome like suicide attempt history (i.e. point-biserial or tetrachoric). Across two studies, shame remained significantly positively associated with NSSI whilst adjusting for negative affect (alongside guilt-proneness; OR = 2.12; Schoenleber, 2013) or internalizing symptoms (alongside guilt proneness and demographic information; OR = 1.37, 95% CI: 1.05-1.78; VanDerhei et al., 2014). A single study found that both guilt and shame were not correlated with suicidal behaviour whilst adjusting for depressive symptoms (Lester, 1998). As before, it was not clear if appropriate correlation coefficients were used given the binary outcome.

Guilt proneness did not have a significant bivariate association with NSSI history. This association remained non-significant when adjusting for shame proneness in one study (Schoenleber, 2013), and a negative association with NSSI history emerged in another study (OR = 0.76, 95% CI: 0.62-0.94), whereby greater guilt proneness was related to a lower risk of NSSI when adjusting for shame proneness (VanDerhei et al., 2014). Guilt proneness was

also not associated with NSSI when adjusting for negative affect (alongside shame proneness;  $OR = 0.69, p = .17$ ; Schoenleber, 2013).

### **Longitudinal Associations**

Only two studies adopted longitudinal methodologies. Duggan and colleagues (2015) investigated body shame over one year in high school students. They found that shame did not distinguish between those with an NSSI history who had not maintained this behaviour compared to controls with no history of NSSI. In contrast, shame remained elevated in those who continued to engage in NSSI relative to the control participants. Brown and colleagues (Brown, Linehan, Comtois, Murray, & Chapman, 2009) followed up a small sample of women diagnosed with BPD. Higher state shame, assessed using items from the PFQ-2, was associated with almost twice the risk of subsequent NSSI (relative risk ratio = 1.88, 95% CI: 1.04-3.38) within a survival analysis. This association did not remain significant when adjusting for feelings of fear. Non-verbal indicators of shame (rated based on video recorded interviews) were associated with NSSI occurrence in non-adjusted analyses (relative risk ratio = 1.99, 95% CI: 1.07-3.69), and remained significant when fear and sadness were also adjusted for (relative risk ratio = 1.86,  $p < .05$ ).

### **Discussion**

The aim of the current study was to provide a systematic review and meta-analysis of the available literature regarding self-harm and its relationships with shame and guilt. Thirty papers were identified for inclusion. Individuals with a history of NSSI typically reported greater shame across a range of different populations and shame sub-types, compared to those without a history of NSSI. Body shame was an exception, where evidence of an association was less clear and varied more dramatically between studies. Shame was also positively correlated with frequency of NSSI engagement. Effect sizes were typically small to moderate

according to Cohen's rules of thumb (Cohen, 1988). These associations between shame and NSSI typically held whilst adjusting for co-occurring feelings of guilt and mood-related symptoms. There was also evidence that shame had a positive association with both suicidal behaviour and self-harm (where measured as a general construct), but studies were fewer and results more varied. In contrast, results were mixed regarding the association between guilt and self-harm (including NSSI and suicidal behaviour). Guilt proneness did not appear to be associated with NSSI or suicidal behaviour, but state guilt was associated with suicide attempt history across four studies. Whilst shame appears linked to self-harm, the lack of longitudinal studies limits conclusions about the direction or temporal characteristics of these associations. The two longitudinal studies identified suggested that feelings of shame may contribute to the risk of NSSI over time.

The results are largely consistent with wider research, where shame has been found to be positively associated with a wide range of mental health difficulties, whilst results regarding guilt have been more ambiguous (Blythin et al., 2018; Carden et al., 2018; Kim et al., 2011; Pugh et al., 2015). This review did exclude some more specific forms of guilt, however, including trauma-related guilt, which may have more of a pronounced relationship with mental health difficulties (Pugh et al., 2015). The decision to exclude these experiences was taken in light of their distinct phenomenology, and the potential confounding effects of event-specific factors. For example, there is evidence the type of trauma can moderate the relationship between guilt and suicidal thinking (Bryan, Ray-Sannerud, Morrow, & Etienne, 2013). It was therefore anticipated that the inclusion of such studies would blur the relationships between key variables and impact upon the generalisability of findings. Overall, guilt has received less research attention than shame, and so we would recommend researchers study shame and guilt in tandem where possible to further establish any differential associations with mental health difficulties.

The results indicate that elevated experiences of shame are associated with self-harm behaviour. As the data are observational and correlational it is not possible to conclude that feelings of shame actively drive or maintain self-harm. These data are consistent with emotion-regulation orientated models of self-harm, which view self-harm as a potential response to aversive affective states like shame (Chapman et al., 2006; Hasking et al., 2017; Nock, 2009), and with people's self-reported reasons for self-harm, that most commonly concern managing negative internal states including shame (Breen, Lewis & Sutherland, 2013; Curtis, 2016; Taylor et al., 2018). Nonetheless, experiences of shame could also be a consequence of self-harm (e.g. self-injury scar-related shame; Bachtelle & Pepper, 2015), or an epiphenomenon related to other processes that drive self-harm. Qualitative research highlights how shame may both be an antecedent and consequence of self-harm (Curtis, 2016). It is also important to recognise that the functions of self-harm vary widely (Taylor et al., 2018) and that different emotions may play a greater or lesser role for different people. Theory suggests that shame may be particularly relevant because it is inherently aversive and closely tied to how individuals perceive and relate to themselves. Self-harm may emerge as a means of regulating these self-directed feelings. There is evidence that greater endorsement of shame-regulation reasons for NSSI (i.e. reducing shame) is associated with greater NSSI frequency (Schoenleber & Berenbaum, 2012; Schoenleber, 2013). These results are also consistent with evidence that a more hostile or critical style of relating to oneself is a risk factor for some forms of self-harm (Forrester, Slater, Jomar, Mitzman, & Taylor, 2017).

In summary, this review highlights that shame and self-harm are linked, but caution should be taken in making further conclusions at this stage. Future research may benefit from moving beyond cross-sectional designs to better understand this relationship. Whilst causality cannot be ascertained within observational data, further evidence of temporality (i.e. that shifts in shame or guilt preceded subsequent changes in self-harm behaviour over time)

would support a stronger case for a potentially causative relationship. Research methodologies that enable a more fine-grained investigation of how individuals respond to experiences of shame in the moment, such as experience sampling methodologies, could also be beneficial (Pratt & Taylor, 2019).

The review identified a number of methodological issues that were apparent in the extant literature. These include the variety of measures used to assess key constructs, namely shame. It is unclear the extent to which many of these measures tap meaningfully distinct constructs. Due to this uncertainty we did not plan to combine different types of guilt or shame into common meta-analyses. This heterogeneity of measures limits the extent to which comparisons between studies can be made and is problematic for summarising effects. It is suggested that researchers within this field would benefit from adopting a common set of measures across studies. This would facilitate comparisons regarding the severity and impact of guilt and shame across studies and population groups.

A further issue relates to the cultural sensitivity of the shame measures utilised within the reviewed studies. Cultural differences in both the precipitants to and manifestations of shame have been highlighted, underscoring the potential role of cultural expectations in the experience of shame (Abu-Kaf & Priel, 2008; Brown, 2006). However, most measures used within reviewed literature were both developed for and tested within predominantly western, individualist cultures. As a result, the extent to which these measures can be considered both sensitive and generalizable to a range of cultures is limited.

The use of meta-analysis provided a means of summarising the size and strength of the association between shame or guilt and self-harm. Whilst this approach has advantages over simple “vote-counting” of significant effects (taking into account the size of studies and the degree of heterogeneity), limitations should also be noted. The meta-analyses, especially those focused on emotion subtypes (i.e. body-related shame, external shame, etc.) included

small numbers of studies, and as such lack precision. Moreover, in many cases high inconsistency was present, but there were not enough studies available to examine possible moderators that may explain this. We also conducted larger meta-analyses by grouping emotion subtypes together, but it should be noted that these may also obscure important differences between subtypes (e.g. body-related shame had a weaker association with NSSI). Overall, these results should therefore be interpreted with caution. Whilst they arguably provide a starting estimate of the size of the bivariate association between these constructs, additional studies would help further confirm these associations. Larger-scale replications of earlier studies adopting the same measures and populations would be particularly beneficial given the current diversity in both populations and measures used.

This review is also limited by the exclusion of non-English language research. The current review also focussed solely on quantitative research. A review of the qualitative research concerning shame and self-harm may shed further light on the potential mechanisms underlying this relationship. Many studies used correlation coefficients to capture the relationship with self-harm frequency. For this reason, the correlation coefficient was typically used as the metric within our meta-analyses. However, in many cases self-harm frequency may be better represented as a count variable, with a Poisson or negative binominal distribution. In such cases correlation coefficients may not capture associations as well, introducing more error. Whilst it was beyond the scope of this review to focus on suicidal ideation as well as behaviour, there may be benefits to also reviewing this literature, particularly where studies examine the transition from thoughts to behaviour.

If shame does contribute to the onset and maintenance of self-harm, then approaches to self-harm intervention and prevention that focus on feelings of shame may be effective. In terms of direct intervention, compassion-focused therapy (CFT) has been developed specifically for individuals who struggle with shame and self-criticism (Gilbert, 2009). There



is preliminary evidence that individuals undergoing CFT experience reductions in shame (Judge, Cleghorn, McEwan, & Gilbert, 2012; Leaviss & Uttley, 2015). Other therapeutic approaches that focus on tolerance of difficult emotional states, such as Dialectical Behaviour Therapy (Rizvi et al., 2011), or focus on self-directed feelings and relating, such as Cognitive Analytic Therapy (Sheard et al., 2000), may also be helpful. Given the heterogeneity in the triggers, functions and forms of self-harm (Klonsky & Muehlenkamp, 2007; Taylor et al., 2018). However, a personalised approach, adopting shame-focused interventions where this appears to be part of the mechanism underlying that individual's self-harm, is likely to be better than a one-size-fits-all approach. At a societal level, recognition of and support for groups where experiences of shame and self-harm are elevated, such as those in the Lesbian, Gay, Bisexual or Transgender (LGBT) community may be beneficial (McDermott, Roen, & Scourfield, 2008; Taylor, Dhingra, Dickson, & McDermott, 2018a). Campaigns and programmes designed to reduce the stigma and shame may help in these instances.

This review summarises the extant literature concerning shame, guilt and self-harm. We provide a preliminary indication of the direction and magnitude of the association between these emotions (and their subtypes) and self-harm. We also highlight key gaps in the literature and future directions, including the need for longitudinal designs and larger-scale replications of earlier studies adopting the same measures and populations.

## References

- Abu-Kaf, S., & Priel, B. (2008). Dependent and self-critical vulnerabilities to depression in two different cultural contexts. *Personality and Individual Differences*, 44, 689-700. doi:10.1016/j.paid.2007.10.002
- Andrews, B., Qian, M., & Valentine, J. D. (2002). Predicting depressive symptoms with a new measure of shame: The Experience of Shame Scale. *The British Journal of Clinical Psychology*, 41, 29-42. doi:10.1348/014466502163778
- Bachtelle, S. E., & Pepper, C. M. (2015). The physical results of nonsuicidal self-injury: The meaning behind the scars. *Journal of Nervous and Mental Disease*, 203, 927-933. doi:http://dx.doi.org/10.1097/NMD.0000000000000398
- Baumeister, R. F. (1990). Suicide as escape from self. *Psychological Review*, 97, 90-113. doi:10.1037/0033-295X.97.1.90
- Beck, A. T., & Steer, R. A. (1991). *Manual for the beck scale for suicide ideation*. San Antonio, TX: Psychological Corporation.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory* (2nd ed.). San Antonio, Texas: Psychological Corporation.
- Bergen, H., Hawton, K., Waters, K., Ness, J., Cooper, J., Steeg, S., & Kapur, N. (2012). Premature death after self-harm: A multicentre cohort study. *The Lancet*, 380, 1568-1574. doi:10.1016/S0140-6736(12)61141-6
- Blythin, S. P. M., Nicholson, H. L., Macintyre, V. G., Dickson, J. M., Fox, J. R. E., & Taylor, P. J. (2018). Experiences of shame and guilt in anorexia and bulimia nervosa: A systematic review. *Psychology & Psychotherapy: Theory, Research & Practice*. doi:10.1111/papt.12198
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to meta-analysis*. Chichester, UK: John Wiley & Sons.

- Brown, B. (2006). Shame resilience theory: A grounded theory study on women and shame. *Families in Society*, 87, 43-52. doi:10.1606/1044-3894.3483
- Breen, A. V., Lewis, S. P., & Sutherland, O. (2013). Brief report: Non-suicidal self-injury in the context of self and identity development. *Journal of Adult Development*, 20, 57-62. doi: 10.1007/s10804-013-9156-8
- Brown, M. Z., Linehan, M. M., Comtois, K. A., Murray, A., & Chapman, A. L. (2009). Shame as a prospective predictor of self-inflicted injury in borderline personality disorder: A multi-modal analysis. *Behaviour Research and Therapy*, 47, 815-822. doi:http://dx.doi.org/10.1016/j.brat.2009.06.008
- Bryan, C. J., Morrow, C. E., Etienne, N., & Ray-Sannerud, B. (2013). Guilt, shame, and suicidal ideation in a military outpatient clinical sample. *Depression & Anxiety*, 30, 55-60. doi:10.1002/da.22002
- Bryan, C. J., Ray-Sannerud, B., Morrow, C. E., & Etienne, N. (2013). Guilt is more strongly associated with suicidal ideation among military personnel with direct combat exposure. *Journal of Affective Disorders*, 148, 37-41. doi:10.1016/j.jad.2012.11.044
- Carden, L., J., Saini, P., Seddon, C., Watkins, M., & James Taylor, P. (2018). Shame and the psychosis continuum: A systematic review of the literature. *Psychology & Psychotherapy: Theory, Research & Practice*. doi:10.1111/papt.12204
- Cawley, R., Pontin, E. E., Touhey, J., Sheehy, K., & Taylor, P. J. (2019). What is the relationship between rejection and self-harm or suicidality in adulthood? *Journal of Affective Disorders*, 242, 123-134. doi:https://doi.org/10.1016/j.jad.2018.08.082
- Chapman, A. L., Gratz, K. L., & Brown, M. Z. (2006). Solving the puzzle of deliberate self-harm: The experiential avoidance model. *Behaviour Research and Therapy*, 44, 371–394. doi:10.1016/j.brat.2005.03.005

- Cheung, M., Gilbert, P., & Iron, C. (2004). An exploration of shame, social rank and rumination in relation to depression. *Personality and Individual Differences*, 36, 1143-1153. doi:10.1016/S0191-8869(03)00206-X
- Chou, C. Y., Tsoh, J., Vigil, O., Bain, D., Uhm, S. Y., Howell, G., . . . Mathews, C. A. (2018). Contributions of self-criticism and shame to hoarding. *Psychiatry Research*, 262, 488-493. doi:10.1016/j.psychres.2017.09.030
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale: Erlbaum Associates.
- Cook, D. R. (1994). *Internalized Shame Scale: Professional Manual*. Menomonie, WI: Channel Press.
- Costa, P. T., & McCrae, R. R. (1985). *The NEO personality inventory manual*. Odessa, FL: Psychological Assessment Resources.
- Curtis, C. (2016). Young women's experiences of self-harm: Commonalities, distinctions and complexities. *Young*, 24, 17-35. doi: 10.1177/1103308815613680
- DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7, 177-188. doi:10.1016/0197-2456(86)90046-2
- Donhauser, S. (2007). *Examining the role of shame as a mediator of childhood sexual abuse and self-injury*. (PhD), St. Mary's University, Texas.
- Duggan, J., Heath, N., & Hu, T. (2015). Non-suicidal self-injury maintenance and cessation among adolescents: A one-year longitudinal investigation of the role of objectified body consciousness, depression and emotion dysregulation. *Child and Adolescent Psychiatry and Mental Health*, 9, 21. doi:10.1186/s13034-015-0052-9
- Edmondson, A. J., Brennan, C. A., & House, A. O. (2016). Non-suicidal reasons for self-harm: A systematic review of self-reported accounts. *Journal of Affective Disorders*, 191, 109-117. doi:http://dx.doi.org/10.1016/j.jad.2015.11.043

- Erchull, M. J., Liss, M., & Lichiello, S. (2013). Extending the negative consequences of media internalization and self-objectification to dissociation and self-harm. *Sex Roles*, 69, 583-593. doi:10.1007/s11199-013-0326-8
- Etzel, J. C. (2004). *A diagnostic exemplar of experiential avoidance: Examining shame and self-harm in battered women with PTSD*. (PhD), School of Saint Louis University.
- Fieldman, J. P. (1989). *Personality characteristics of adolescent suicide attempters*. (PhD), Hahnemann Medical College and Hospital.
- Forrester, R. L., Slater, H., Jomar, K., Mitzman, S., & Taylor, P. J. (2017). Self-esteem and non-suicidal self-injury in adulthood: A systematic review. *Journal of Affective Disorders*, 221, 172-183. doi:10.1016/j.jad.2017.06.027
- Gandy, S. (2013). *Undertsanding the mechanisms of non-suicidal self-inflicted injury (NSSII) among adult survivors of child sexual abuse*. (PhD), Alliant International University.
- Gilbert, P. (1997). The evolution of social attractiveness and its role in shame, humiliation, guilt and therapy. *British Journal of Medical Psychology*, 70, 113–147. doi:10.1111/j.20448341.1997.tb01893.x
- Gilbert, P. (1998). What is shame? Some core issues and controversies. In P. Gilbert & B. Andrews (Eds.), *Shame: Interpersonal behavior, psychopathology and culture* (pp. 3-36). New York: Oxford University Press.
- Gilbert, P. (2009). Introducing compassion-focused therapy. *Advances in Psychiatric Treatment*, 15, 199-208. doi:10.1192/apt.bp.107.005264
- Gilbert, P., Pehl, J., & Allan, S. (1994). The phenomenology of shame and guilt: An empirical investigation. *Psychology and Psychotherapy: Theory, Research and Practice*, 67, 23-36. doi:10.1111/j.2044-8341.1994.tb01768.x

- Gilbert, P., & Procter, S. (2006). Compassionate mind training for people with high shame and self-criticism: overview and pilot study of a group therapy approach. *Clinical Psychology & Psychotherapy*, *13*, 353-379. doi:10.1002/cpp.507
- Glassman, L. H., Weierich, M. R., Hooley, J. M., Deliberto, T. L., & Nock, M. K. (2007). Child maltreatment, non-suicidal self-injury, and the mediating role of self-criticism. *Behaviour Research and Therapy*, *45*, 2483-2490.  
doi:http://dx.doi.org/10.1016/j.brat.2007.04.002
- Goldman-Mellor, S. J., Caspi, A., Harrington, H., Hogan, S., Nada-Raja, S., Poulton, R., & Moffitt, T. E. (2014). Suicide attempt in young people: A signal for long-term health care and social needs. *JAMA Psychiatry*, *71*, 119-127.  
doi:10.1001/jamapsychiatry.2013.2803
- Goss, K., Gilbert, P., & Allan, S. (1994). An exploration of shame measures—I: The other as Shamer scale. *Personality and Individual Differences*, *17*, 713-717.  
doi:https://doi.org/10.1016/0191-8869(94)90149-X
- Gratz, K. L. (2001). Measurement of deliberate self-harm: Preliminary data on the Deliberate Self-Harm Inventory. *Journal of Psychopathology and Behavioral Assessment*, *23*, 253-263. doi:10.1023/A:1012779403943
- Hanson, R. K., & Tangney, J. P. (1996). *The Test of Self-Conscious Affect—Socially Deviant populations (TOSCA-SD)*. Ottawa, Ontario, Canada: Department of the Solicitor General of Canada.
- Harder, D. H., & Zalma, A. (1990). Two promising shame and guilt scales: A construct validity comparison. *Journal of Personality Assessment*, *55*, 729-745.  
doi:10.1080/00223891.1990.9674108

- Hasking, P., Whitlock, J., Voon, D., & Rose, A. (2017). A cognitive-emotional model of NSSI: Using emotion regulation and cognitive processes to explain why people self-injure. *Cognition and Emotion*, *31*, 1543-1556. doi:10.1080/02699931.2016.1241219
- Hawton, K., Bergen, H., Cooper, J., Turnbull, P., Waters, K., Ness, J., & Kapur, N. (2015). Suicide following self-harm: Findings from the Multicentre Study of self-harm in England, 2000-2012. *Journal of Affective Disorders*, *175*, 147-151. doi:10.1016/j.jad.2014.12.062
- Higgins, J. P. T., & Green, S. (2009). *Cochrane handbook for systematic reviews of interventions*: The Cochrane Collaboration.
- Higgins, J. P. T., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*, *21*, 1539-1558. doi:10.1002/sim.1186
- Hughes, M. A., Knowles, S. F., Dhingra, K., Nicholson, H. L., & Taylor, P. J. (2018). This corrosion: A systematic review of the association between alternative subcultures and the risk of self-harm and suicide. *British Journal of Clinical Psychology*. doi:10.1111/bjc.12179
- Izadi, M. (2014). *Perfectionism, shame, and non suicidal self injury*. (PhD), Alliant International University.
- Johns, D., & Holden, R. R. (1997). Differentiating suicidal motivations and manifestations in a nonclinical population. *Canadian Journal of Behavioural Science*, *29*, 266-274. doi:10.1037/0008-400X.29.4.266
- Joiner, T. E. (2005). *Why people die by suicide*. Cambridge, MA: Harvard University Press.
- Judge, L., Cleghorn, A., McEwan, K., & Gilbert, P. (2012). An exploration of group-based compassion focused therapy for a heterogeneous range of clients presenting to a community mental health team. *International Journal of Cognitive Therapy*, *5*, 420-429. doi:10.1521/ijct.2012.5.4.420

- Kealy, D. (2019). *Unpublished research data*.
- Kealy, D., Spidel, A., & Ogrodniczuk, J. S. (2017). Self-conscious emotions and suicidal ideation among women with and without history of childhood sexual abuse. *Counselling and Psychotherapy Research, 17*, 269–275. doi:10.1002/capr.12140
- Kim, S., Thibodeau, R., & Jorgensen, R. S. (2011). Shame, guilt, and depressive symptoms: A meta-analytic review. *Psychological Bulletin, 137*, 68-96. doi:10.1037/a0021466
- Klonsky, E. D. (2009). The functions of self-injury in young adults who cut themselves: Clarifying the evidence for affect-regulation. *Psychiatry Research, 166*, 260-268. doi:10.1016/j.psychres.2008.02.008
- Klonsky, E. D. (2011). Non-suicidal self-injury in United States adults: Prevalence, sociodemographics, topography and functions. *Psychological Medicine, 41*, 1981-1986. doi:doi:10.1017/S0033291710002497
- Klonsky, E. D., & Glenn, C. R. (2009). Assessing the functions of non-suicidal self-injury: psychometric properties of the Inventory of Statements About Self-injury (ISAS). *Journal of Psychopathology & Behavioral Assessment, 31*, 215-219. doi:10.1007/s10862-008-9107-z
- Klonsky, E. D., & Muehlenkamp, J. J. (2007). Self-injury: A research review for the practitioner. *Journal of Clinical Psychology, 63*, 1045-1056. doi:10.1002/jclp.20412
- Kontopantelis, E., & Reeves, D. (2010). Metaan: Random-effects meta-analysis. *The STATA Journal*. doi: 10.1177/1536867X1001000307
- Lamb, C. S. (2004). *Shame and self harm in axis II disorders*. (ClinPsyD), University of Edinburgh.
- Leaviss, J., & Uttley, L. (2015). Psychotherapeutic benefits of compassion-focused therapy: An early systematic review. *Psychological Medicine, 45*, 927-945. doi:10.1017/s0033291714002141



- Lester, D. (1998). The association of shame and guilt with suicidality. *The Journal of Social Psychology, 138*, 535-536. doi:10.1080/00224549809600407
- Lewis, H. B. (1971). Shame and guilt in neurosis. *Psychoanalytic Review, 58*, 419-438.
- Lindberg, S. M., Hyde, J. S., & McKinley, N. M. (2006). A measure of objectified body consciousness for preadolescent and adolescent youth. *Psychology of Women Quarterly, 30*, 65-76. doi:10.1111/j.1471-6402.2006.00263.x
- Linehan, M. M., Comtois, K. A., Brown, M. Z., Heard, H. L., & Wagner, A. (2006). Suicide Attempt Self-Injury Interview (SASII): Development, reliability, and validity of a scale to assess suicide attempts and intentional self-injury. *Psychological Assessment, 18*, 303-312. doi:10.1037/1040-3590.18.3.303
- Mallindine, C. (2002). *To what degree does social rank theory help us to understand selfinjurious behaviour amongst patients in a high security setting?* (ClinPsyD), University College London.
- Matos, M., Pinto-Gouveia, J., Gilbert, P., Duarte, C., & Figueiredo, C. (2015). The Other As Shamer Scale – 2: Development and validation of a short version of a measure of external shame. *Personality and Individual Differences, 74*, 6-11.  
doi:https://doi.org/10.1016/j.paid.2014.09.037
- McDermott, E., Roen, K., & Scourfield, J. (2008). Avoiding shame: young LGBT people, homophobia and self-destructive behaviours. *Culture, Health & Sexuality, 10*, 815-829. doi:10.1080/13691050802380974
- McKinley, N. M., & Hyde, J. S. (1996). The Objectified Body Consciousness Scale: Development and validation. *Psychology of Women Quarterly, 20*, 181-215.  
doi:10.1111/j.1471-6402.1996.tb00467.x
- McLeod, L. D. (2002). *The development and initial validation of the domains of shame questionnaire.* (PhD), Queen's University.

- Milligan, R.-J., & Andrews, B. (2005). Suicidal and other self-harming behaviour in offender women: The role of shame, anger and childhood abuse. *Legal and Criminological Psychology, 10*, 13-25. doi:10.1348/135532504X15439
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & The, P. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Medicine, 6*, e1000097. doi:10.1371/journal.pmed.1000097
- Morey, L. C. (1991). *Personality assessment inventory professional manual*. Odessa, FL: Psychological Assessment Resources.
- Muehlenkamp, J. J. (2006). Empirically supported treatments and general therapy guidelines for non-suicidal self-injury. *Journal of Mental Health Counseling, 28*, 166-185. doi:10.17744/mehc.28.2.6w61cut2lxjdg3m7
- Muehlenkamp, J. J., Claes, L., Havertape, L., & Plener, P. L. (2012). International prevalence of adolescent non-suicidal self-injury and deliberate self-harm. *Child and Adolescent Psychiatry and Mental Health, 6*, 10-18. doi:10.1186/1753-2000-6-10
- National Collaborating Centre for Mental Health. (2004). *Self-harm: The short-term physical and psychological management and secondary prevention of self-harm in primary and secondary care*: National Institute for Clinical Excellence.
- Nelson, A., & Muehlenkamp, J. J. (2012). Body attitudes and objectification in non-suicidal self-injury: Comparing males and females. *Archives of Suicide Research, 16*, 1-12. doi:10.1080/13811118.2012.640578
- Nock, M. K. (2009). Why do people hurt themselves? New insights into the nature and functions of self-injury. *Current Directions in Psychological Science, 18*, 78-83. doi:10.1111/j.1467-8721.2009.01613.x
- Nock, M. K., Holmberg, E. B., Photos, V. I., & Michel, B. D. (2007). Self-Injurious Thoughts and Behaviors Interview: Development, reliability, and validity in an

- adolescent sample. *Psychological Assessment*, 19, 309-317. doi:10.1037/1040-3590.19.3.309
- Paulsen, K. L. (2013). *Consequences of anger and shame reactions in adults sexually abused as children*. Alliant International University.
- Pratt, D., & Taylor, P. J. (2019). Experience sampling in the study of self-harm. In J. Palmier-Claus, G. Haddock & F. Varese (Eds.) *Experience sampling in mental health research* (pp. 97-110). Oxon: Routledge
- Pritchard, T. I. (2009). *Body attitudes, self-injury and eating disorders*. (MA), University of British Columbia.
- Pugh, L. R., Taylor, P. J., & Berry, K. (2015). The role of guilt in the development of post-traumatic stress disorder: A systematic review. *Journal of Affective Disorders*, 182, 138-150. doi:10.1016/j.jad.2015.04.026
- Resnick, H. S., & Weaver, T. (1994). *Habit Questionnaire*. Charleston: Medical University of South Carolina.
- Reynolds, W. M. (1987). *Suicidal Ideation Questionnaire (SIQ): Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Ribeiro, J. D., Franklin, J. C., Fox, K. R., Bentley, K. H., Kleiman, E. M., Chang, B. P., & Nock, M. K. (2016). Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: A meta-analysis of longitudinal studies. *Psychological Medicine*, 46, 225-236. doi:10.1017/S0033291715001804
- Rizvi, S. L., Brown, M. Z., Bohus, M., & Linehan, M. M. (2011). The role of shame in the development and treatment of borderline personality disorder *Shame in the therapy hour* (pp. 237-260). Washington, DC, US: American Psychological Association.

- Ross, S., & Heath, N. L. (2002). A study of the frequency of self-mutilation in a community sample of adolescents. *Journal of Youth and Adolescents*, 31, 66-77.  
doi:10.1023/A:1014089117419
- Rossotto, E., Yager, J., & Rorty, M. (1994). *Impulsive and self-harm behaviours among women with bulimia nervosa*. Paper presented at the The 6th International Conference on Eating Disorders, New York.
- Rowe, S. L., French, R. S., Henderson, C., Ougrin, D., Slade, M., & Moran, P. (2014). Help-seeking behaviour and adolescent self-harm: A systematic review. *Australian and New Zealand Journal of Psychiatry*, 48, 1083-1095. doi:10.1177/0004867414555718
- Royal College of Psychiatrists. (2010). *Self-harm, suicide and risk: Helping people who self-harm*. London: Royal College of Psychiatrists.
- Rusch, N., Lieb, K., Gottler, I., Hermann, C., Schramm, E., Richter, H., . . . Bohus, M. (2007). Shame and implicit self-concept in women with borderline personality disorder. *The American Journal of Psychiatry*, 164, 500-508.  
doi:10.1176/ajp.2007.164.3.500
- Rutherford, R. M. (2015). *Understanding the role of shame in women with depression*. (PhD), Pacifica Graduate Institute.
- Sansone, R. A., & Sansone, L. A. (2010). Measuring self-harm behavior with the self-harm inventory. *Psychiatry*, 7, 16-20. Retrieved from  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2877617/>
- Schaefer, K. E. (2014). *Suicidal ideation in a U.S. jail: Demographic and psychiatric correlates and a test of baumeister's escape theory*. (PhD), George Mason University.
- Schoenleber, M., & Berenbaum, H. (2012). Shame regulation in personality pathology. *Journal of Abnormal Psychology*, 121, 433-446. doi:10.1037/a0025281

- Schoenleber, M. L. (2013). *Testing alternative motivational models for self-injurious behaviour*. (PhD), University of Illinois.
- Seidlitz, L., Conwell, Y., Duberstein, P., Cox, C., & Denning, D. (2001). Emotion traits in older suicide attempters and non-attempters. *Journal of Affective Disorders*, 66, 123-131. doi: 10.1016/S0165-0327(00)00300-1
- Sheard, T., Evans, J., Cash, D., Hicks, J., King, A., Morgan, N., . . . Sunder, K. (2000). A CAT-derived one to three session intervention for repeated deliberate self-harm: a description of the model and initial experience of trainee psychiatrists in using it. *The British Journal of Medical Psychology*, 73, 179-196. doi:10.1348/000711200160417
- StataCorp. (2015). College Station, TX: StataCorp LP.
- Svanborg, P., & Asberg, M. (2001). A comparison between the Beck Depression Inventory (BDI) and the self-rating version of the Montgomery Asberg Depression Rating Scale (MADRS). *Journal of Affective Disorders*, 64, 203-216.
- Swannell, S. V., Martin, G. E., Page, A., Hasking, P., & St John, N. J. (2014). Prevalence of nonsuicidal self-injury in nonclinical samples: Systematic review, meta-analysis and meta-regression. *Suicide and Life-threatening Behavior*, 44, 273-303. doi: 10.1111/sltb.12070
- Tangney, J. P. (1990). Assessing individual differences in proneness to shame and guilt: Development of the self-conscious affect and attribution inventory. *Personality and Social Psychology*, 59, 102-111. doi:10.1037/0022-3514.59.1.102
- Tangney, J. P., & Dearing, R. L. (2002). *Shame and guilt*. New York: The Guilford Press.
- Tangney, J. P., Dearing, R. L., Wagner, P. E., & Gramzow, R. (2000). *The test of self-conscious affect-3 (TOSCA-3)*. Fairfax, VA: George Mason University.

- Tangney, J. P., Miller, R. S., Flicker, L., & Barlow, D. H. (1996). Are shame, guilt, and embarrassment distinct emotions? *Journal of Personality and Social Psychology*, 70, 1256-1269. doi: 10.1037/0022-3514.70.6.1256
- Tangney, J. P., Stuewig, J., Krishnan, S., Youman, K., Appel, M., Roop, L., & Durbin, K. M. (2008). *The Test of Self-Conscious Affect—Version 4 (TOSCA-4)*. Fairfax, VA: George Mason University.
- Tangney, J. P., Stuewig, J., & Mashek, D. J. (2007). Moral emotions and moral behavior. *Annual Review of Psychology*, 58, 345-372.  
doi:10.1146/annurev.psych.56.091103.070145
- Taylor, P. J., Dhingra, K., Dickson, J. M., & McDermott, E. (2018a). Psychological correlates of self-harm within gay, lesbian and bisexual UK university students. *Archives of Suicide Research*, 1-16. doi:10.1080/13811118.2018.1515136
- Taylor, P. J., Dhingra, K., Dickson, J. M., & McDermott, E. (2018). Psychological correlates of self-harm within gay, lesbian and bisexual UK university students. *Archives of Suicide Research*. doi: 10.1080/13811118.2018.1515136
- Taylor, P. J., Hutton, P., & Wood, L. (2015). Are people at risk of psychosis also at risk of suicide and self-harm? A systematic review and meta-analysis. *Psychological Medicine*, 45, 911-926. doi:10.1017/s0033291714002074
- Taylor, P. J., Jomar, K., Dhingra, K., Forrester, R., Shahmalak, U., & Dickson, J. M. (2018). A meta-analysis of the prevalence of different functions of non-suicidal self-injury. *Journal of Affective Disorders*, 227, 759-769. doi:10.1016/j.jad.2017.11.073
- Taylor, P. J., McDonald, J., Smith, M., Nicholson, H., & Forrester, R. (2019). Distinguishing those with current, past, and no history of Non-Suicidal Self-Injury: Shame, social comparison and self-concept integration *Journal of Affective Disorders*, 246, 182-188. doi: 10.1016/j.jad.2018.12.033

- Todd, L. (2002). *Intentional self-injury in male prisoners: The role of social status, shame and anger*. University College London.
- Van Orden, K. A., Cukrowicz, K. C., Witte, T. K., & Joiner, T. E. (2012). Thwarted belongingness and perceived burdensomeness: Construct validity and psychometric properties of the interpersonal needs questionnaire. *Psychological Assessment, 24*, 197-215. doi:10.1037/a0025358
- Van Orden, K. A., Witte, T. K., Cukrowicz, K. C., Braithwaite, S. R., Selby, E. A., & Joiner, T. E., Jr. (2010). The interpersonal theory of suicide. *Psychological Review, 117*, 575-600. doi:10.1037/a0018697
- VanDerhei, S., Rojahn, J., Stuewig, J., & McKnight, P. E. (2014). The effect of shame-proneness, guilt-proneness, and internalizing tendencies on nonsuicidal self-injury. *Suicide & Life-Threatening Behavior, 44*, 317-330. doi:10.1111/sltb.12069
- Veroniki, A. A., Jackson, D., Viechtbauer, W., Bender, R., Bowden, J.,...Salanti, G. (2014). Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Research Synthesis Methods, 7*, 55-79. doi: 10.1002/jrsm.1164
- Watson, D., & Clark, L. A. (1994). *The PANAS-X: Manual for the Positive and Negative Affect Schedule-expanded form*. Ames: The University of Iowa.
- Weingarden, H., Renshaw, K. D., Wilhelm, S., Tangney, J. P., & DiMauro, J. (2016). Anxiety and shame as risk factors for depression, suicidality, and functional impairment in body dysmorphic disorder and obsessive compulsive disorder. *Journal of Nervous and Mental Disease, 204*, 832-839. doi:10.1097/nmd.0000000000000498
- Wiklander, M., Samuelsson, M., Jokinen, J., Nilsson, A., Wilczek, A., Rylander, G., & Asberg, M. (2012). Shame-proneness in attempted suicide patients. *BMC psychiatry, 12*, 50-50. doi:10.1186/1471-244X-12-50

Williams, J. M. G. (1997). *Cry of pain: Understanding suicide and self-harm*. Oxford: Penguin Books.

Williams, J. W., Plassman, B. L., Burke, J., Holsinger, T., & Benjamin, S. (2010). *Preventing Alzheimer's disease and cognitive decline. Evidence report/technology assessment No. 193. (Prepared by the duke evidence-based practice center under contract No. HHSA 290-2007-10066-I.)*. Rockville, MD: Agency for Healthcare Research and Quality.

Xavier, A., Cunha, M., Pinto-Gouveia, J., & Paiva, C. (2013). *Exploratory study of the Portuguese version of the risk-taking and self-harm inventory for adolescents*. Paper presented at the I World Congress of Children and Youth Health Behaviors/IV National Congress on Health Education.

Xavier, A., Pinto Gouveia, J., & Cunha, M. (2016). Non-suicidal self-injury in adolescence: The role of shame, self-criticism and fear of self-compassion. *Child & Youth Care Forum*, 45, 571-586. doi:10.1007/s10566-016-9346-1

Zanarini, M. C., Vujanovic, A. A., Parachini, E. A., Boulanger, J. L., Frankenburg, F. R., & Hennen, J. (2003). A screening measure for BPD: The McLean Screening Instrument for Borderline Personality Disorder (MSI-BPD). *Journal of Personality Disorders*, 17, 568-573. Retrieved from <https://guilfordjournals.com/doi/pdf/10.1521/pedi.17.6.568.25355>



Table 1

## Summary of Study Characteristics

<b>Authors, years &amp; country</b>	<b>Design</b>	<b>Participant characteristics</b>	<b>Comparison group characteristics (if applicable)</b>	<b>NSSI measures</b>	<b>Shame/ guilt measures</b>
Brown et al. (2009), US	Longitudinal	N = 77 women seeking psychotherapy for BPD and self-harm; Mean age = 30.0 years, SD = 7.3	-	Suicide Attempt Self- Injury Interview (SASII)	Shame items adapted from the Harder Personal Feelings Questionnaire (PFQ2)
Bryan et al. (2013), US	Cross- sectional	N = 151 outpatients at military mental health clinics; Mean age = 34.12 years, SD = 8.41; Female = 36%	-	Self-Injurious Thoughts and Behavior Interview (SITBI)	The Harder Personal Feelings Questionnaire (PFQ2)
Donhauser (2007), US	Cross- sectional	N = 51 adult childhood sexual abuse survivors; 18-65 years; Female = 84%	-	Deliberate Self-Harm Inventory (DSHI)	Internalised Shame Scale (ISS)
Duggan et al. (2015), Canada	Longitudinal	N = 120 high school students; Mean age = 12.34 years, SD = 0.48; Female = 56%	-	How I Deal with Stress Questionnaire (HIDS)	Objectified Body Consciousness Scale – Youth (OBCS-Y)

Erchull et al. (2013); US	Cross-sectional	N = 160 Adult females; Mean age = 23.12, SD = 3.69	-	Deliberate Self-Harm Inventory (DSHI)	Objectified Body Consciousness Scale (OBCS)
Etzel (2004), US	Cross-sectional	N = 65 female victims of intimate partner violence; Mean age = 34.2 years, SD = 8.4	-	Habit Questionnaire for self-injurious behavior	Experience of Shame Scale (ESS)
Fieldman (1989), US	Cross-sectional	N = 30 acute adolescent psychiatric inpatients; Mean age = 15.8 years; Female = 66.7%	-	The Reynolds Suicide Ideation Questionnaire (RSIQ)	Guilt Scale
Gandy (2013), US	Cross-sectional	N = 103 adult survivor of childhood sexual abuse; Mean age = 39.88 years, SD = 13.82; Female = 71.8%	-	modified Suicidal Behavior Questionnaire, (SBQ)	Test of Self-Conscious Affect - 3 (TOSCA-3)
Izadi (2014), US	Cross-sectional	N = 50 individuals with a history of NSSI; Mean age = 27.0 years, SD = 7.1; Female = 74%	-	Deliberate Self-Harm Inventory (DSHI); modified Suicidal Behavior Questionnaire, (SBQ)	Test of Self-Conscious Affect - 3 (TOSCA-3); Shame Variability Questionnaire (SVQ; unpublished)
Kealy et al (2017), Canada	Cross-sectional	N = 68 female psychiatric outpatients; Mean age = 36.6 years, SD = 12.0	-	Suicidal Behavior Questionnaire - Revised (SBQ-R)	The Harder Personal Feelings Questionnaire (PFQ2)

Kealy (unpublished, 2019), Canada	Cross - sectional	N = 137 psychiatric outpatients; Mean age = 33.39, SD = 11.98; Female = 69%	-	Item taken from the McLean screening instrument for Borderline Personality Disorders (MSI-BPD)	The Harder Personal Feelings Questionnaire (PFQ2)
Lamb (2004), UK	Cross- sectional	N = 30 psychiatric inpatients and outpatients; Mean age = 37.67 years, SD = 12.4; Female = 86.7%	-	Self-harm Inventory (SHI)	Internalised Shame Scale (ISS)
Lester (1998), US	Cross- sectional	N = 116 university students; Mean age = 21.9 years, SD = 4.6; Female = 67.2%	-	Beck Depression Inventory (BDI)	Shame and Guilt scale
Mallindine (2002), UK	Cross- sectional	N = 21 male inpatients in high secure settings with a NSSI history; Mean age = 31.1 years; SD = 6.99  N = 13 female inpatients in a high secure setting with a NSSI history; Mean age = 28.1; SD = 5.33	N = 15 male inpatients in high secure settings without a history of NSSI; Mean age = 39.8 years; SD = 8.58	Information from clinical records	Other As Shamer scale (OAS)
McLeod (2002), Canada	Cross- sectional	N = 236 psychology university students; Mean	-	Suicidal Ideation and Behavior Questionnaire (SIBQ)	Test of Self-Conscious Affect (TOSCA); Harder The Harder Personal

		age = 19.2 years, SD = 1.0; Female = 72.4%			Feelings Questionnaire (PFQ2); The Domains of Shame Questionnaire (developed within this research)
Milligan & Andrews (2005), UK	Cross-sectional	N = 89 female prisoners; Mean age = 31.8 years, SD = 9.37	-	Impulsive Behavior scale (IBS)	Experience of Shame Scale (ESS)
Nelson & Muehlenkamp (2012), US	Cross-sectional	N = 341 university students; Mean age = 20.2 years, SD = 2.0; Female = 82.4%	-	The Deliberate Self-Harm Inventory (DSHI)	Objectified Body Consciousness Scale (OBCS)
Paulson (2013), US	Cross-sectional	N = 56 adults survivors of childhood sexual abuse; Mean age = 41.7 years; SD = 15.0; Female = 34%	-	Non Suicidal Self Injury Interview (NSSII)	Test of Self-Conscious Affect - 3 (TOSCA-3)
Pritchard (2009), Online (Canada, UK, US, Australia, New Zealand)	Cross-sectional	N = 54 adults reporting self-harm; Mean age = 21.85 years, SD = 3.58; Female = 66.7%	-	Deliberate Self-Harm Inventory (DSHI)	Objectified Body Consciousness Scale (OBCS)
		N = 18 adults reporting eating disorders; Mean age = 24.28 years, SD = 5.46; Female = 94.4%			
		N = 106 adults reporting eating disorder and self-			

		harm; Mean age = 23.0 years, SD = 6.65; Female = 90.56%			
Rusch et al. (2007), Germany	Cross-sectional	N = 60 female inpatients diagnosed with BPD; Mean age = 27.8 years, SD = 6.9  N = 30 female inpatients with Social Phobia; Mean age = 35.1, SD = 11.9	N = 60 female healthy comparison group; Mean age = 26.6, SD = 7.4	Psychiatric interview	Test of Self-Conscious Affect - 3 (TOSCA-3); The Harder Personal Feelings Questionnaire (PFQ2)
Rutherford (2015), US	Cross-sectional	N = 126 women from mental health clinics and hospitals; Mean age = 34 years, SD = 8.4	-	Study specific measure of suicidal behaviour	Experience of Shame Scale (ESS)
Schaefer (2014), US	Cross-sectional	N = 511 prison inmates; Mean age = 32.19 years, SD = 10.05; Female = 32%	-	Study specific measure of suicidal behaviour	Test of Self-Conscious Affect for Socially Deviant populations (TOSCA-SD)
Schoenleber (2013), US	Cross-sectional	N = 61 female university students; Mean age = 18.9 years, SD = 0.8  N = 54 women living in the community; Mean	-	Inventory of Statements About Self-Injury (ISAS)	Test of Self-Conscious Affect - 3 (TOSCA-3)

		age = 24.8 years, SD = 6.6			
Seidlitz et al. (2001), US	Cross-sectional	N = 47 inpatients aged over 50 years with depression and a history of suicide attempts; Mean age = 59.5 years, SD = 8.3; Female = 60%	N = 38 inpatients aged over 50 years with depression and no history of suicide attempts; Mean age = 62.9 years, SD = 10.3; Female = 55%	Study specific measure of suicidal behaviour	Emotional traits assessed via NEO Personality Inventory - Revised (NEO-PI-R)
Taylor et al. (2018), UK	Cross-sectional	N = 51 adults who report having engaged in NSSI in the past year; 96% aged under 30 years; Female = 80%	N = 110 university students with no history of NSSI; 100% aged under 30 years; Female = 86%	Self-Injurious Thoughts and Behavior Interview (SITBI)	Experience of Shame Scale (ESS)
		N = 44 adults with a history of NSSI, but no NSSI in the past year; 94% aged under 30 years; Female = 98%			
Todd (2002), UK	Cross-sectional	N = 73, male prisoners; Mean age; 30.88 years, SD = 7.08	-	Study specific measure of NSSI	Test of Self-Conscious Affect for Socially Deviant populations (TOSCA-SD); Other As Shamer scale (OAS)
VanDerhei et al. (2014), US	Cross-sectional	N = 378 university students; Mean age = 20.84 years, SD = 4.7; Female = 71%	-	Inventory of Statements About Self-Injury (ISAS)	Test of Self-Conscious Affect - 3 (TOSCA-3)

Weingarden et al. (2016), US	Cross-sectional	N = 114 adults with Body Dysmorphic Disorder; Mean age = 30.22 years, SD = 10.86; Female = 92%	N = 133 health adult controls; Mean age = 36.44 years, SD = 13.28; Female = 76%	Suicide Behaviors Questionnaire - Revised (SBQ-R)	Test of Self-Conscious Affect - 4 (TOSCA-4)
		N = 114 adults with Obsessive Compulsive Disorder; Mean age = 30.60 years, SD = 10.66; Female = 86%			
Wiklander et al. (2012), Sweden	Cross-sectional	N = 498 psychiatric outpatients	N = 161 healthy controls; Mean age = 44.6 years, SD = 7.7; Female = 63%	Montgomery Åsberg Depression Rating Scale (MADRS-S)	Test of Self-Conscious Affect (TOSCA)
		N = 108 patients with a BPD diagnosis and history of attempted suicide; Mean age = 29.9 years, SD = 8.0; Female = 97%			
		N = 67 patients without a BPD diagnosis and history of attempted suicide; Mean age = 34.8 years, SD = 12.4; Female = 67%			

		N = 162 patients without a history of attempted suicide; Mean age = 45.3 years, SD = 8.9; Female = 72%			
Xavier et al. (2016), Portugal	Cross-sectional	N = 782 Middle and secondary school children; Mean age = 14.9 years; SD = 1.8 Female = 52.8%	-	Risk Taking and Self Harm Inventory for Adolescents - Portuguese Version (RTSHIA)	Other As Shamer scale – 2, Portuguese version (OAS-2)

---

BDI (Beck et al., 1961) = Beck Depression Inventory; DSHI (Gratz, 2001) = Deliberate Self-Harm Inventory; HIDS (Ross & Heath, 2002) = How I Deal with Stress Questionnaire; HQ (Resnick & Weaver, 1994) = Habit Questionnaire for self-injurious behaviour; IBS (Rossotto et al., 1994) = Impulsive Behavior Scale; ISAS (Klonsky & Glenn, 2009) = Inventory of Statements About Self-injury; MADRS-S (Svanborg & Åsberg, 2001) = Montgomery-Asberg Depression Rating Scale; MSI-BPD (Zanarini et al., 2003) = Mclean Screening Instrument for BPD; PAI (Morey, 1991) = Personality Assessment Inventory; RTSHIA (Portuguese version: Xavier et al. 2013) = The Risk-Taking and Self-Harm Inventory for Adolescents Portuguese Version; SASII (Linehan et al., 2006) = Suicide Attempt Self-Injury Interview; SBQ-R (Osman et al., 2002) = Suicidal Behavior Questionnaire Revised; SHI (Sansone & Sansone, 1995) = Self-harm Inventory; SIBQ (Johns & Holden, 1997) = Self-Injurious Behaviour Questionnaire; SIQ (Reynolds, 1987) = Suicide Ideation Questionnaire; SITB (Nock et al., 2007) = Self-Injurious Thoughts and Behavior Interview. ESS (Andrews et al., 2002) = Experiences of Shame Scale; ISS (Cook, 1994) = Internalised Shame Scale; NEO-PI-R (Costa & McCrae, 1985) = the Revised NEO Personality Inventory; OAS (Goss et al., 1995) = The Other as Shamer Scale; OAS-2 (Matos et al., 2015) = The Other as Shamer Scale-2 Portuguese Version; OBCS (McKinley & Hyde, 1996) = Objectified Body Consciousness Scale; OBCS-Y (Lindberg et al., 2006) = Objectified Body Consciousness Scale – Youth; PFQ-2 (Harder & Zalma, 1990) = the Harder Personal Feelings Questionnaire; SVQ (Brown et al., unpublished) = Shame Variability Questionnaire; TOSCA (Tangney et al., 1989) = Test of Self-Conscious Affect; TOSCA-3 (Tangney et al., 2000) = Test of Self-Conscious Affect 3; TOSCA-SD (Hanson & Tangney, 1995) = Test of Self-Conscious Affect Socially Deviant; TOSCA-4 (Tangney et al., 2008) = Test of Self-Conscious Affect 4.



Table 2

## Risk of Bias Assessment

<b>Authors</b>	<b>Unbiased cohort selection</b>	<b>Selection minimizes baseline differences in demographic factors</b>	<b>Sample size calculated</b>	<b>Validated method for ascertaining clinical status or participant group</b>	<b>Validated methods for assessing shame/guilt</b>	<b>Validated methods for assessing self-harm</b>	<b>Blind outcome assessment</b>	<b>Adequate follow-up period</b>	<b>Adequate handling of missing data</b>	<b>Appropriate analytic methods</b>
Brown et al (2009)	Yes	Yes	No	Yes	Yes	Yes	No	Yes	No	Yes
Bryan et al (2013)	Yes	N/A	No	Yes	Yes	Yes	Yes	N/A	Yes	Yes
Donhauser (2008)	Partial	N/A	Yes	N/A	Yes	Yes	N/A	N/A	Yes	Yes
Duggan et al (2015)	Yes	Yes	No	Yes	Yes	Partial	No	Yes	Yes	Yes
Erchull et al (2013)	No	N/A	No	N/A	Yes	Yes	Yes	N/A	Yes	No
Fieldman (1989)	Yes	Yes	No	Yes	Partial	Yes	No	N/A	Unclear	Yes

---

Gandy										
(2014)	Partial	N/A	Yes	Yes	Yes	Yes	Yes	N/A	Yes	Yes
Etzel										
(2004)	Partial	N/A	No	No	Yes	Yes	No	N/A	Unclear	Yes
Izadi										
(2014)	Partial	N/A	No	Yes	Yes	Yes	No	N/A	Yes	Yes
Kealy et al										
(unpub.)	Yes	N/A	No	Yes	Yes	No	No	N/A	Yes	Yes
Kealey et al.										
(2017)	No	N/A	No	Yes	Yes	Yes	No	N/A	Yes	Yes
Lamb										
(2004)	No	N/A	Yes	Yes	Yes	Yes	No	N/A	Yes	Yes
Lester										
(1998)	No	N/A	No	N/A	Yes	Yes	No	N/A	Unclear	Unclear
Mallindine										
(2002)	Unclear	Yes	No	Yes	Yes	No	No	N/A	Yes	Yes
McLeod										
(2002)	No	N/A	No	N/A	Yes	Yes	Partial	N/A	Yes	Yes
Milligan &										
Andrews										
(2005)	No	N/A	No	Yes	Yes	Yes	No	N/A	Yes	Yes

---

---

Nelson & Muehlenkam p (2012)	No	N/A	No	N/A	Yes	Yes	Yes	N/A	Yes	Yes
Paulson (2014)	Partial	N/A	No	Yes	Yes	Partial	Yes	N/A	Yes	Yes
Pritchard (2009)	No	N/A	No	Yes	Yes	Yes	Yes	N/A	Yes	Yes
Rusch et al (2007)	Unclear	N/A	No	Yes	Yes	No	No	N/A	Yes	Yes
Rutherford (2016)	Partial	N/A	Yes	N/A	Yes	No	Yes	N/A	Yes	Yes
Schaefer (2014)	Yes	N/A	No	Yes	Yes	Yes	Partial	N/A	Unclear	Yes
Schoenleber (2013)	Partial	N/A	No	Yes	Yes	Yes	No	N/A	Yes	Yes
Seidlitz et al (2001)	Yes	N/A	No	Yes	Yes	No	No	N/A	Unclear	Yes
Taylor et al (2018)	No	N/A	No	Yes	Yes	Yes	No	N/A	Yes	Yes
Todd	Partial	Yes	No	Yes	Yes	No	No	N/A	Yes	Yes

---

---

(2002)										
VanDerhei										
et al (2014)	No	N/A	No	N/A	Yes	Yes	No	N/A	Yes	Yes
Weingarden										
et al (2016)	No	N/A	No	Yes	Yes	No	Yes	N/A	Yes	Yes
Wiklander et										
al (2012)	No	No	No	Yes	Yes	No	No	NA	Yes	Yes
Xavier et al										
(2016)	Unclear	N/A	No	N/A	Yes	Partial	Yes	N/A	Yes	Partial

---

N/A = Not Applicable

Table 3

Summary of Bivariate Associations between Shame or Guilt Variables and Self-Harm

Shame or guilt variable	Outcome	N/K	Association	$I^2$
Shame proneness	NSSI frequency	488/4	$r = .25$ (95% CI: .08, .40)	48%
	NSSI history (binary)	493/3	$d = 0.42$ (95% CI: 0.24, 0.60)	0%
	Suicide attempt history (binary)	1306/4	$d = 0.36$ (95% CI: 0.05, 0.66)	80%
Body shame	NSSI frequency	239/2	$r = .07$ (95% CI: -.30, .42)	85%
	NSSI history (binary)*	826/5	$d = 1.61$ (95% CI: -.32, 3.55)	99%
	Suicide attempt frequency	119/1	$r = .27$ (95% CI: .10, .43)	NA
	Self-harm history (binary)	89/1	$d = 1.24$ (95% CI: 0.78, 1.70)	NA
External shame	NSSI frequency	782/1	$r = .39$ (95% CI: .33, .45)	NA
	NSSI history (binary)	105/2	$d = 0.51$ (95% CI: 0.12, 0.90)	0%
Characterological or internal shame	NSSI frequency	62/1	$r = .33$ (95% CI: .09, .54)	NA
	NSSI history (binary)	205/1	$d = 1.71$ (95% CI: 1.39, 2.03)	NA
	Suicide attempt frequency	119/1	$r = .27$ (95% CI: .10, .43)	NA
	Self-harm history (binary)	119/2	$d = 0.39$ (95% CI: 0.02, 0.77)	0%
	Self-harm frequency	20/1	$r = -.38$ (95% CI: -.70, .08)	NA

State shame	Suicide attempt history (binary)	278/3	$d = 0.58$ (95% CI: -0.12, 1.27)	74%
	Self-harm history (binary)	137/1	<b><math>d = 0.37</math> (95% CI: 0.03, 0.70)</b>	NA
Performance, appearance and relationship related shame	Suicide attempt frequency	236/1	<b><math>r = .20 - .26</math> (95% CI: .07, .38)</b>	NA
Guilt proneness	NSSI frequency	386/2	$r = -.01$ (95% CI: -.11, .09)	0%
	NSSI history (binary)	360/1	$r = -.07$ (95% CI: -.17, .03)	NA
	Suicide attempt history (binary)	1335/5	$d = 0.12$ (95% CI: -0.08, 0.32)	51%
State guilt	Suicide attempt history (binary)	363/4	<b><math>d = 0.59</math> (95% CI: 0.25, 0.93)</b>	33%
	Self-harm history (binary)	137/1	<b><math>d = 0.47</math> (95% CI: 0.13, 0.81)</b>	NA

---

Note: NSSI = non-suicidal self-injury; *K* refers to independent samples rather than studies; Meta-analysis undertaken where two or more studies available.

Effects in bold are significant at  $p < .05$ ; \* Included one study with unusually large effect size. Exclusion of this study result in  $d = 0.35$  (-0.10, 0.79),  $I^2 = 81\%$




Figure 1: Flow diagram of included studies

Supplementary Table 1

## Summary of Bivariate Associations between Shame or Guilt Variables and Self-Harm by Individual Study

Study	Shame or guilt variable (measure)	Outcome	Standardized association	Non-standardized association
Bryan et al (2013)	State shame (PFQ2) State Guilt (PFQ2)	Suicide attempt history	$d = 1.25^*$ $d = 1.15^*$	Mean difference = 8.90 Mean difference = 5.93
Duggan et al (2015) <sup>a</sup>	Body shame (OBCS-Y)	NSSI history	Males: $d = -0.03$ Females: $d = 0.01$	Males: Mean difference = -0.04 Females: Mean difference = 0.01
Erchull et al (2013)	Body shame (OBCS)	NSSI history	$r = .18^*$	
Fieldman (1989)	Guilt proneness (Guilt Scale)	Suicide attempt history	$d = 1.13^*$	Mean difference = 18.3
Gandy (2014/2013)	Shame proneness (TOSCA-3)	NSSI frequency NSSI history	$r = .26^*$ $r = .25^*$	
Etzel (2004)	Characterological shame (ESS) Body shame (ESS)	NSSI frequency	$r = .33^*$ $r = .27^*$	
Kealy et al (unpub.)	State shame (PFQ2) State guilt (PFQ2)	Self-harm history Self-harm history	$d = 0.37^*$ $d = 0.47^*$	Mean difference = 2.71 Mean difference = 2.19
Kealey et al (2017)	State shame (PFQ2) State guilt (PFQ2)	Suicide attempt history Suicide attempt history	$d = 0.51$ $d = 0.52$	Mean difference = 3.20 Mean difference = 2.29
Lamb (2004)	Internal shame (ISS)	Self-harm frequency Self-harm history	$r = -.38$ $d = 0.09$	- Mean difference = 1.03
Mallindine (2002)	External shame (OAS)	NSSI history	$d = 0.53$	Mean difference = 8.40
McLeod (2003)	Performance, appearance and relationship related shame	Suicide attempt frequency	$r = .20-.26^*$	
Milligan & Andrews	Characterological shame (ESS) Body shame (ESS)	Self-harm history Self-harm history	$d = 0.49^*$ $d = 1.24^*$	Mean difference = 3.94 Mean difference = 4.35



(2005)					
Nelson & Muehlenkamp (2012) <sup>b</sup>	Body shame (OBCS)	NSSI history	$d = 6.86^*$	Mean difference = 0.59	
Paulson (2013)	Shame proneness (TOSCA-3)	NSSI frequency	$r = .15$		
Pritchard (2014)	Body shame (OBCS)	NSSI frequency	$r = -.11$		
Rusch et al (2007)	Shame proneness (TOSCA-3)	Suicide attempt history	$d = -0.26$	Mean difference = -1.76	
	Guilt proneness (TOSCA-3)	Suicide attempt history	$d = -0.24$	Mean difference = -1.07	
	State shame (PFQ2)	Suicide attempt history	$d = -0.04$	Mean difference = -0.24	
	State guilt (PFQ2)	Suicide attempt history	$d = 0.52$	Mean difference = 1.93	
Rutherford (2016)	Characterological shame (ESS)	Suicide attempt frequency	$r = .27^*$		
	Body shame (ESS)	Suicide attempt frequency	$r = .27^*$		
Schaefer (2014)	Shame proneness (TOSCA-SD)	Suicide attempt frequency	$r = .09$		
	Guilt proneness (TOSCA-SD)	Suicide attempt frequency	$r = .02$		
Schoenleber (2013)	Shame proneness (TOSCA-3)	NSSI frequency	$r = .58^*$	-	
	Guilt proneness (TOSCA-3)	NSSI history	$d = .53^*$	Mean difference = 0.30	
Seidlitz et al (2001)	Guilt proneness (NEO-PI-R)	NSSI frequency	$r = 0.14$		
		Suicide attempt history	$d = 0.34$	Mean difference = 1.08	
Taylor et al (2018)	Characterological shame (ESS)	NSSI history	$d = 1.71^*$	Mean difference =	
	Body shame (ESS)	NSSI history	$d = 0.89^*$	13.01	
				Mean difference = 3.24	
Todd (2002)	External shame (OAS)	NSSI history	$d = 0.50^*$	Mean difference = 7.21	
VanDerhei et al (2014)	Shame proneness (TOSCA-3)	NSSI frequency	$r = .16^*$		
	Shame proneness (TOSCA-3)	NSSI history	$r = .19^*$		
	Guilt proneness (TOSCA-3)	NSSI frequency	$r = -.02$		

	Guilt Proneness( TOSCA-3)	NSSI history	$r = -.07^c$	
Weingarden et al (2016)	Shame proneness (TOSCA-4) Guilt proneness (TOSCA-4)	Suicide attempt history Suicide attempt history	$d = 0.61^*$ $d = 0.07$	Mean difference = 19.14
Wiklander et al (2012)	Shame proneness (TOSCA) Guilt proneness (TOSCA)	Suicide attempt history Suicide attempt history	$d = 0.60^*$ $d = 0.16$	Mean difference = 1.31 Mean difference = 5.93 Mean difference = 1.13
Xavier et al (2016)	External shame (OAS-2)	NSSI frequency	$r = .39^*$	

---

Note: NSSI = non-suicidal self-injury; ESS = Experience of Shame Scale; ISS = Internalised Shame Scale; NEO-PI-R = NEO Personality Inventory Revised; OAS = Other As Shamer Scale; OAS-2 = The Other as Shamer Scale-2 Portugese Version ; OBCS = Objectified Body Consciousness Scale; OBCS-Y = Objectified Body Consciousness Scale – Youth; TOSCA = Test of Self-Conscious Affect; TOSCA-SD = Test of Self-Conscious Affect for Socially Deviant populations; \*  $p < .05$ ; <sup>a</sup> Means for those with a history of NSSI are based on aggregating data from individuals who either continued NSSI or stopped NSSI at follow-up; <sup>b</sup> standardized effect size should be treated with caution due to unusually small reported standard deviations; <sup>c</sup> value reported as significant in paper but on inspection coefficient not large enough to be significant given sample size (assuming two-tailed test and  $\alpha = .05$ ).

Supplementary Table 2

Summary of Bivariate Associations between Shame or Guilt Variables and Self-Harm with REML Estimation

Shame or guilt variable	Outcome	N/K	Association	$I^2$
Shame proneness	NSSI frequency	488/4	$r = .24$ (95% CI: .09, .38)	41%
	NSSI history (binary)	493/3	$d = 0.42$ (95% CI: 0.24, 0.60)	0%
	Suicide attempt history (binary)	1306/4	$d = 0.36$ (95% CI: 0.04, 0.68)	82%
Body shame	NSSI frequency	239/2	$r = .07$ (95% CI: -.30, .42)	85%
	NSSI history (binary)	826/5	$d = 1.61$ (95% CI: -.97, 4.20)	>99%
External shame	NSSI history (binary)	105/2	$d = 0.51$ (95% CI: 0.12, 0.90)	0%
Characterological or internal shame	Self-harm history (binary)	119/2	$d = 0.39$ (95% CI: 0.02, 0.77)	0%
State shame	Suicide attempt history (binary)	278/3	$d = 0.58$ (95% CI: -0.13, 1.29)	75%
Guilt proneness	NSSI frequency	386/2	$r = -.01$ (95% CI: -.11, .09)	0%
	Suicide attempt history (binary)	1335/5	$d = 0.10$ (95% CI: -0.02, 0.21)	0%
State guilt	Suicide attempt history (binary)	363/4	$d = 0.59$ (95% CI: 0.25, 0.93)	33%

Note: NSSI = non-suicidal self-injury; *K* refers to independent samples rather than studies; Meta-analysis undertaken where two or more studies available. Effects in bold are significant at  $p < .05$ ;

