
Citation:

Guo, N and Zhao, L and Zhao, Y and Li, Q and Xue, X and Wu, L and Gomez Escalada, M and Wang, K and Peng, W (2020) Comparison of the Chemical Composition and Biological Activity of Mature and Immature Honey : An HPLC/QTOF/MS-Based Metabolomic Approach. Journal of Agricultural and Food Chemistry. ISSN 0021-8561 DOI: <https://doi.org/10.1021/acs.jafc.9b07604>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/6641/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

A comparison of the chemical composition and biological activity of mature and immature honeys: an HPLC/QTOF/MS-based metabolomics approach

Nana Guo^{†,&}, Liuwei Zhao^{†,&}, Yazhou Zhao[†], Qiangqiang Li[†], Xiaofeng Xue[†], Liming Wu^{†,#}, Margarita Gomez Escalada^{††}, Kai Wang^{†,*}, Wenjun Peng^{†,*}

[†] Institute of Apicultural Research, Chinese Academy of Agricultural Sciences, Beijing, 100093, China;

[#] Northwest University, Xi'an 710069, Shanxi, China;

^{††} School of Clinical and Applied Sciences, Leeds Beckett University, Leeds LS1 3HE, UK.

* Corresponding authors:

Dr. Kai Wang. Tel: 15652631288. Fax: +86 10 62594643.

E-mail: kaiwang628@gmail.com

Dr. Wenjun Peng. Tel: 13911120162. Fax: +86 10 62594643.

E-mail: pengwenjun@vip.sina.com

& These authors contributed equally to this work.

1 **Abstract:** Harvesting uncapped immature honey (IMH) followed by dehydration is a
2 typical counterfeit honey production process, but the differences between IMH and
3 capped mature honey (MH) have previously not been well described. In this study, MH
4 and IMH from the *Apis mellifera* colonies in the same rapeseed flower season were
5 compared. MH was found to have lower water content, acidity and higher fructose
6 content. HPLC-Q-TOF/MS based untargeted metabolomic analysis indicated that MH
7 had a distinct metabolite composition to IMH. Targeted metabolomic analysis on 20
8 major polyphenolic constituents showed higher accumulation in MH. MH had greater
9 bacteriostatic effect and stronger free radical scavenging effect. Whilst both honeys
10 mitigated cell damage caused by H₂O₂, the effective dosage of IMH was higher and its
11 inducing effect on the anti-oxidant gene expression was weaker. Overall, MH was
12 shown to be of better quality than IMH not only because of its richer polyphenolic
13 composition, but also due to its stronger biological activity.

14

15 **Keywords:** honey, mature, immature, metabolomic analysis, bioactivity, HPLC-Q-
16 TOF/MS.

17

Introduction

Honey is a miraculous product resulting from millions of years of coevolution between plants and honey bees, *Apis* species.¹ It is a natural sweetener that originates from the plant nectar or honey dew collected by bees and further matured inside the bee hive.² Bees build a band of honeycomb above the brood cells in their nests to store honey and pollen. The mature honey is capped with white wax for long-term storage.² Stored honey and pollens act as food sources, whilst the honeycomb band provides insulation during the winter period or on days without foraging activities.³

The process of honey maturation begins with the forager bees taking the nectar or honey dew to the hives.² The forager transfers these carbohydrates from their stomach to storer bees.⁴ Storer bees normally add their own substances, like enzymes from the hypopharyngeal glands to convert the sucrose into glucose or fructose.⁵ The acids from the bees' stomach lowers the pH of the IMH. At the same time, the drying process by their evaporation behavior further decreases the moisture of the honey.² The duration of honey maturation varies from one to eleven days depending on-colony size, humidity, climatic conditions and the botanical origins of the nectar.⁶ After the honey matures, bees cover the honey with a wax lid as protection and to prevent unwanted fermentation and spoilage.⁷

Due to its great value, honey has been subjected to fraud threat since ancient times. Counterfeit honeys remain a serious threat to the global beekeeping business. Typical frauds may involve diluting honey using a variety of syrups,^{8,9} lightening honey color using ion-exchange resins,⁴ labeling the honey with fraudulent geographical and/or

botanical origins,¹⁰ artificial feeding of bees during a nectar flow and harvesting the immature (uncapped) honey.¹¹ The latter fraud type is quite prevalent, since some beekeepers think this can increase the honey harvest. The unmatured honey then undergoes dehydration with vacuum dryers, resulting in most physiochemical features still falling within regulatory.¹¹ Collecting uncapped honey followed manual dehydration is now regarded as an illicit practice. It is already accepted that water content might be a major difference between mature (capped) and immature (uncapped) honey.¹¹ However, during the natural transformation of nectar into honey, bees can add specific substances. The chemical composition of honey is complex, not only consisting of sugars and water, but also other constituents, including amino acids, vitamins, minerals and plant polyphenolic acids.¹² These components together endow honey with distinct flavors and biological activities.¹³ Nevertheless, it remains to be determined whether these minor substances result in significant differences in chemical compositions and biological activities between mature honey (MH) and immature honey (IMH). To understand these two types of honey better, this study compared the chemical composition and biological activities (anti-oxidative and anti-microbial) of MH and IMH.

Materials and methods

Chemicals and reagents

Methanol (MeOH) and formic acid (FA) were purchased from Fisher Scientific Inc (Pittsburgh, PA, USA). Trolox, ascorbic acid, quercetin, gallic acid and other standards were purchased from Sangon biological engineering co. LTD (Shanghai,

China). Solid-phase extraction (C18) was purchased from Waters scientific Inc. LB Nutrient Agar was purchased from Beijing Aoboxing biotechnology co. LTD. *Escherichia coli*, *Staphylococcus aureus*, and *Bacillus subtilis* were obtained from Institute of microbiology, Chinese Academy of Sciences, China. The solid phase extraction (SPE) cartridges were obtained from Waters (Milford, Mass, USA). All the cartridges contained 500 mg of C₁₈.

Honey sample collection and physicochemical analysis

Raw honey samples were collected from three *A. mellifera* L. colonies in Sichuan, China, during the flower season from March 1st to March 30th, 2019. Three colonies with the same potential were selected from the experimental bee hive. The honey in the colonies was cleared and only a small amount was left for bees to maintain a basic life. The honey collected by bees and brought back to the nest for no more than 24 h was recorded as immature (uncapped) honey (IMH), and the honey stored in the honeycomb with a beeswax seal until the sealed area of beeswax was greater than 70% was recorded as capped mature (capped) honey (MH). Three IMH and MH samples were separately collected from each colony. A total of 18 samples were collected, including 9 MH and 9 IMH, and stored at -20 °C in the dark prior to use.

These 18-batches of rapeseed honey were subjected to chemical analysis. Indicators including water, glucose, fructose, sucrose, acidity and 5-hydroxymethylfurfural (HMF) were determined as previously described.¹⁴

Preparation of active substances

Five grams of honey sample was added into 10 mL deionized water followed by

sonicating—at 60 kHz for 10 min and centrifugation at 8000 r/min for 5 min. The supernatant was collected and added to the SPE cartridges that were preconditioned initially with 5 mL of methanol (MeOH) and then 5mL of water. The supernatant samples passed through the cartridges at a flow rate of approximately 1 mL/min. The analytes were eluted with 8 mL of methanol. The resulting eluate was dried using a nitrogen stream to obtain immature honey extract (IMHE) and mature honey extract (MHE). Both extracts were stored at -20 °C.

HPLC-Q-TOF/MS analysis of honey extract

The honey extracts were re-dissolved to a pre-determined concentration with MeOH. The solution was then filtered with a 0.22 µm nylon membrane and placed in a brown vial. High performance liquid chromatography combined with quadrupole time-of-flight mass spectrometry (HPLC-Q-TOF/MS, 6545) system was used to perform the chromatographic analysis in the negative ionization mode. An Agilent Zorbax Poroshell EC-C18 column (2.1 mm x 100 mm, 2.7 µm) was used to separate the extracted compounds. Analytes were separated by linear gradient elution with ultrapure water containing 0.1% formic acid (v/v) (A) and MeOH (B) at a flow rate of 0.25 mL min⁻¹. The linear gradient elution program was: 0–1 min, 5% B; 1–6 min, 55% B; 6–20 min, 95% B; 20–26 min, 95% B; 26–27 min, 5% B. The column temperature was set to 30 °C with an injection volume of 2 µL. The parameters of ESI source were as follows: a nebulizer pressure of 40 psi, a capillary voltage of 3500 V, a fragmentor voltage of 120 V, a drying gas (N₂) flow rate of 8 L/min, a drying-gas temperature of 320 °C and a mass range of m/z 100–1700.

Determination of total phenolic and flavonoid content

The measurement of total polyphenol content in the honey extracts was determined by the Folinol- Ciocalteu method. 100 μ L of the extract was mixed with 100 μ L of Folin and Ciocalteu's phenol reagent. The mixture was incubated in the dark for 5 min, followed by the addition of 300 μ L sodium carbonate solution (2% w/v) and mixed. The reaction proceeded in the dark for 120 min. The absorbance was measured at 765 nm. Gallic acid was used to calculate the standard curve and the results were expressed as mg of gallic acid equivalents (GAEs) per g of honey extraction.

For the measurement of total flavonoid content, 150 μ L of the sample was mixed with 10 μ L aluminium nitrate (100 g/L), 10 μ L potassium acetate (9.8 g/L) and 330 μ L of distilled water. The reaction proceeded in the dark for 120 min. The absorbance of the product was determined at 415 nm. Quercetin was used to calculate the standard curve and the results were expressed as mg of Quercetin equivalents (QEs) per g of honey extraction.

Antioxidant activity

Free radical scavenging ability

Various concentrations of honey phenolic extracts (0.2 mL) were mixed with 0.2 mL of ethanolic solution containing DPPH radicals. The mixture was shaken vigorously and left to stand for 30 min in the dark or until stable absorption values were obtained. The reduction of the DPPH radicals was determined by measuring the absorption at 517 nm. The concentration of the extract providing 50% of radical scavenging activity (IC₅₀) was determined by a linear curve established by mass concentration and

clearance. The results were expressed as mg of Trolox per g of honey extraction.

Vitamin C was used as the positive control.

Various concentrations of honey phenolic extracts (0.15 mL) were mixed with 0.25 mL of ethanol solution containing ABTS⁺ working liquid. The mixture was shaken vigorously and left to stand for 10 min in the dark until stable absorption values were obtained. The reduction of the ABTS⁺ radical was determined by measuring the absorption at 734 nm. The concentration of the extract providing 50% of radicals scavenging activity (IC₅₀) was determined by a linear curve established by mass concentration and clearance. The results were expressed as mg of Trolox per g of honey extraction. Vitamin C was used as positive control.

Reducing ability

Various concentrations of the honey extracts (0.3 mg) were mixed evenly with 75 µL of sodium phosphate buffer (pH 6.6) and 75 µL of 1% potassium ferricyanide (w/v). The mixture was then incubated at 50 °C for 20 min. After 75 µL of 10% (v/v) trichloroacetic acid was added, the mixture was centrifuged at 2000 rpm for 10 min. The upper layer (300 µL) was mixed with 300 µL of deionized water and 60 µL of 0.1% of ferric chloride (v/v). Then the mixture was shaken, and the absorbance was measured spectrophotometrically at 700 nm. The concentration of the extract providing an absorbance of 0.5 (IC₅₀) was determined by a linear curve established by mass concentration and absorbance. The results were expressed as mg of Trolox per g of honey extraction. Vitamin C was used as the positive control.

Cell culture and cell viability assay

Mouse skin fibrocytes L929 cells were incubated in high-glucose Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% heat-inactivated fetal bovine serum (FBS), 100 µg/mL streptomycin and 100 U/mL penicillin at 37 °C in an incubator with 5% CO₂. Cells were then passaged once every 1.5 days. The toxicity of the honey extract and H₂O₂ was determined by using a CCK-8 kit (Dojindo, Japan) following the manufacturer's instructions. The absorbance was measured at 450 nm using a microplate reader (Bio-Rad Model 550, CA, USA).

Total RNA isolation and quantification

L929 cells were pretreated with designated concentrations of the honey extract for 2 h, then stimulated with 500 µM H₂O₂ for 24 h. Total RNA was collected and extracted using the RNA Pure Kit (Carry Helix Biotechnologies Co., Ltd., Beijing, China). The concentration and purity of the RNA measured using the Nano Drop 2000 ultramicro spectrophotometer. RNA was reverse transcribed by PrimeScript™ RT Master MIX kit (TaKaRa, Dalian, China) and the product stored at -20 °C.

Quantitative real-time PCR was implemented using Bioer LineGene 9600 system (Hangzhou, China) with the SYBR premix EXTaq (TaKaRa, Dalian, China) according to the two-step reaction method. The gene-specific primers of selected cytokines were listed in Supplemental Table 1. The expression of housekeeping gene GAPDH was used to normalize the expression levels of these target genes, the specificity was confirmed by dissociation curve analysis and gel electrophoresis. And the relative expression levels of target genes were calculated using $2^{-\Delta\Delta C_t}$ method.

Anti-microbial activity

Anti-microbial activity was measured by an agar diffusion method. LB agar was sterilized and cooled to 60 °C and 100 µL bacterial solution (10^6 CFU/mL) was added to each 30 mL agar to prepare the bacteria-containing medium. After the plate was set, the sample solution to be tested was evenly added into a sterilized Oxford cup (100 µL/cup). The negative control was deionized water, and the positive control was ampicillin solution (5 g/mL). Plates were incubated at 37 °C for 16 h. A Vernier caliper was used to measure the diameter of the zone inhibition (in mm), and the average values were obtained by repeating the test in triplicate. The results were presented as a mean \pm SD.

Statistical analyses

General analysis

Data was obtained from at least three independent experiments and shown as the mean \pm SD of the indicated replicates. Statistical differences were analyzed using One way ANOVA test followed by Bonferroni post hoc analysis and Student's unpaired *t*-test $P < 0.05$ was accepted as statistically significant.

Untargeted metabolomics statistical analysis

Raw data obtained by HPLC-Q-TOF/MS system was preliminarily processed to provide structured data in an appropriate format for subsequent data analysis. The resulting data was extracted by the Profinder software tool in the MassHunter Qualitative Analysis Software (Agilent Technologies) and converted into CEF files. The list of all possible components, as represented by the full TOF mass spectral data, was created in this way. Each compound was described by mass retention, time, and abundance. Then

data filtering was performed with Mass Profiler Professional (Agilent Technologies) software. Before statistical analysis, filtration of data matrix by sample frequency was also applied. Only substances with a frequency greater than 70% were selected for further analysis. The sample differences were statistically analyzed by using One way ANOVA test followed by Bonferroni post hoc analysis and Student's paired t -test (again, $P < 0.05$ was considered significant).

The materials showing significant difference between groups were matched and analyzed by using Traditional Chinese Medicines (TCM) database (Agilent Technologies). Principal-component analysis (PCA) was also used to analyze the difference between samples, and score plots were produced.

Targeted metabolomics statistical analysis

Targeted compound ion chromatogram was extracted by Mass Hunter Qualitative Analysis software (Agilent Technologies) for all samples. We conducted qualitative analysis according to the retention time, molecular weight and mass spectrometry fragment and quantitative analysis through the external standard method. The peak areas were used to construct standard curves with $R^2 \geq 0.99$. A t -test of the quantitative results was performed to analyze the difference of phenolic substances in MH and IMH samples.

Results

Physicochemical analysis

To study the difference between mature (capped) honey (MH) and immature (uncapped) honey (IMH), physical and chemical indicators were assessed as shown in

Table 1. The indexes of MH were as follows: moisture content $18.31 \pm 1.52\%$, acidity 13.67 ± 1.88 mL/kg, total sugar content $73.22 \pm 2.71\%$, fructose content $36.40 \pm 0.37\%$. IMH: moisture content $31.20 \pm 1.81\%$, acidity 19.9 ± 0.42 mL/kg, total sugar content $61.11 \pm 2.09\%$, fructose content $30.86 \pm 0.64\%$. Compared with IMH, MH samples had lower water content, lower acidity and a higher fructose content.

Metabolomic profiling

Untargeted study

We enriched the active components in the honey and analyzed their differences in the honey extracts by metabolomics using Agilent MPP software. In the first step, the molecular features (MFs) that were present in all injections were retained for each species. The total number of the molecules were 3,751 from all injections, and significantly reduced to 3,060 after the filtering step. The results from the data analysis are represented by a Venn diagram (Figure. 1. A). The results showed that 2,572 chemicals were detected in MH, and 2,686 substances were detected in IMH, with 2,198 substances in common. Secondly, molecular features were further filtered based on p-values calculated by one-way ANOVA. A p-value cutoff of 0.05 was set as the filtering standard to maintain the MFs which differed significantly. The final filtering step was conducted using fold change (FC) analysis (Figure. 1. B). The value of FC was calculated as the MF abundance ratios between each of the two groups. Only the MFs with FC of 2.0 or higher abundance were picked out. As shown in Figure. 1. B, each grey dot represents a chemical while the red dots highlight those substances that were significantly up-regulated in MH group compared with those in IMH group. Equally,

the blue dots highlight those substances that were significantly down-regulated. The substances without significant difference between the two groups are represented by gray dots. To evaluate the variation between the two honey samples and simplify the data management, PCA was used. The raw data of 3,060 MFs were subjected to PCA algorithm in the MPP software (Figure. 1. C). The 2D PCA shown represents 67.99% of the total variation. The first principle component (PC1) accounted for 60.29% of the total data variability, while the second one accounted for 7.7%. The distribution areas of the two samples are clearly differentiated. IMH is mainly distributed in the positive axis of PC1, while MH is mainly in the negative axis of PC1.

Targeted study

Twenty types of phenolic compounds were qualitatively analyzed by HPLC-Q-TOF/MS (Figure. 2 & Table 2). Further quantitative analysis showed that except for vanillic acid and syringic acid, the concentrations of 3, 4-dihydroxybenzoic acid, chlorogenic acid, caffeic acid, p-coumaric acid, ferulic acid, 3-O-acetylpinobanksin, quercetin, hesperitin, pinobanksin, naringenin, galangin, luteolin, kaempferol, apigenin, pinocembrin, 3-(3, 4-Dimethoxyphenyl)-2-propenoic acid, chrysin, caffeic acid, and phenethyl ester in MH was significantly higher than that of IMH. Among them, kaempferol, apigenin, pinocembrin, 3-(3, 4-dimethoxyphenyl)-2-propenoic acid, chrysin and caffeic acid phenethyl ester were only detected in MH.

Comparison on the anti-bacterial activity

We measured the anti-bacterial activity of honey solution against *Escherichia coli*, *Staphylococcus aureus* and *Bacillus subtilis* (Table 3). The result revealed that the zones

of inhibition of MH and IMH on *Escherichia coli* and *Staphylococcus aureus* were 19.47±0.31 mm, 14.13±0.68 mm and 17.29±0.78 mm, 12.80±0.98 mm, respectively. However, neither of them showed any obvious inhibitory effect on *Bacillus subtilis* as no zones were observed. The zones of inhibition were significantly higher for the MH than for the IMH (both $P<0.05$), indicating that MH has a stronger bacteriostatic effect than IMH.

Comparisons of the anti-oxidant activities

The experimental results (Table 4) of oxidation resistance showed the content of total phenolics and total flavonoids in the extracts of honey. The MH and IMH samples contained total phenolics of 12.99 ± 0.19 mg gallic acid equivalent (GAE) per gram and 12.20 ± 0.16 mg gallic acid equivalent (GAE) per gram, respectively, and total flavonoids as 3.53 ± 0.07 mg quercetin equivalent (QE) per gram and 3.41 ± 0.01 mg quercetin equivalent (QE) per gram, respectively. The reduction capacity was 36.97 ± 0.53 mg Trolox equivalent per gram and 28.41 ± 0.76 mg Trolox equivalent per gram in MH and IMH, respectively. DPPH and ABTS⁺ Free radical scavenging power were 21.89 ± 0.08 mg Trolox equivalent per gram and 19.60 ± 0.36 mg Trolox equivalent per gram, 37.82 ± 0.90 mg Trolox equivalent per gram and 32.30 ± 0.81 mg Trolox equivalent per gram.

Subsequently, cell experiments were conducted to further study the anti-oxidant effect of honey. The results showed that when the concentration of H₂O₂ was 500 µM, the anti-oxidative activity of L929 cells was significantly reduced. However, the honey extract had no toxic effect in the range of test concentrations and honey extract

treatment significantly improved the proliferation activity of cells stimulated by H₂O₂ (Figure. 3). MHE concentration of 400 µg/mL significantly increased the expression of anti-oxidant genes HO-1, TXNRD, GCLM and NQO1 (Figure. 4). IMHE was only effective when the concentration of IMHE was 600 µg/mL, and the expression of anti-oxidation gene NQO1 did not significantly promote the effect.

Discussion

To determine whether immaturity of honey might adversely affect honey quality, we performed a comparative study of the differences between the capped and uncapped honeys from the same botanic source (*Brassia campestris L*). We found notable physicochemical and bioactive differences between immature and mature honeys.

The physicochemical indices, including water, sugar, acidity, and HMF, have been previously investigated.¹⁶ These are the basic indicators to characterize the quality of honey. Studies have shown that the physicochemical parameters of honey can be utilized to distinguish between mature and immature acacia honey.¹⁷

The water content of honey represents a highly important quality parameter for the its shelf life during storage. The origin of honey, nectars normally, contain more than 50% water but bees will further dehydrate the honey in the comb environment.¹⁸ Therefore, early harvest of the immature honey leads to high moisture content.¹⁹ High water content increases the possibility of honey being fermented during long storage periods.²⁰ The average water content of our MH samples (18.31 ± 1.52 g per 100 g) were below the required threshold standard of the European Regulations of Quality

(no more than 20 g per 100 g). The average water content of IMH samples was well above this standard (31.20 ± 1.81 g per 100 g) suggesting reduced quality and increased possibility of fermentation.

Increased free acidity is an important indicator of microbial spoilage and freshness of honey. When acidity values are above the standard limits, it indicates sugar fermentation due to the formation of acetic acid by alcohol hydrolysis.²¹ Depending on the flower source or geographic area, the free acidity of honey varies. As reported in a previous study, honey's acidity ranges from 9.7 to 29.5 meq/kg.²² In our study, the free acidity of capped rapeseed honey was 13.67 ± 1.88 meq/kg, whereas for uncapped it one is 19.9 ± 0.42 meq/kg. Both results were below the required standard (less than 50 meq/kg).²³ This shows that, the acidity of honey not only depends on nectar source species but is also affected by maturity.

HMF represents an indicator of honey freshness and authenticity whereas high concentrations of HMF in honey indicates overheating and poor storage conditions or adulteration of the honey with inverted syrup.²⁴ Honey storage at 35°C causes an increase of HMF that exceeds the allowed limit (40 mg/Kg).²⁵ However, HMF was not detected in our study samples, which means that the honey samples were fresh, but the immature uncapped honey would need to be further dehydrated including a heating process which may increase HMF levels.

Sugar is the main ingredient in all honeys, with concentrations of up to 80%, and explains why honey is the oldest natural sweetener.²⁶ The sugar content of honey varies from harvest time, due to the flutter of the wings of the bees or the variance

among the nectars.²⁷ The most abundant sugars in honey are fructose and glucose, with higher quantities of fructose in the majority of honeys.²⁸ The percentage of fructose and glucose in our samples ranged from 15.5-49.3% and 18.2–48.0%, respectively. Sucrose was not detected or detected in very low amounts in the honey samples, this is not surprising since sucrose is broken down into glucose, fructose and other monosaccharide by enzymes secreted by bees during honey maturation process. Rapeseed honeys normally contain more glucose than fructose, but in this study, glucose content was found to be at lower concentrations than fructose. This may be due to the different geographical origin and the local climate. The results also suggest that a higher percentage of fructose may be produced as honey is matured in the hive for a long time. Mature honey has been shown to have lower water content, higher fructose content and lower acidity than immature honey, and therefore MH appears to be of better quality.

Recently, metabolic profiling methods have been robustly applied to detect the intrinsic similarities and differences in metabolites within biological samples.²⁹ In the present study, the Mass Profiler Professional (MPP) software was applied in the analysis of the chromatographic data, which enabled us to compare accurately, comprehensively and quickly the major constituents between MH and IMH samples.³⁰ This MPP analysis has already been used for screening and development of drugs and food inspection, the results of which have been well-recognized in related fields. The method has been confirmed to be precise, accurate and sensitive enough for untargeted analysis.^{30,31} The present study is the first application of MPP technology in determining

honey maturity. We analyzed 18 batches of honey samples collected from three different hives. To ensure comparability each comb analysed contained both immature (uncapped) and mature (capped) honey, during a single rapeseed blossom season. We also performed multivariate statistical analysis for classification, prediction, and characterization of marker compounds. Among them, some metabolites, including organic acids, flavonoids, polyphenols, terpenes and others have been reported from honey.^{32,33} We investigated the differences in the metabolite composition in honeys under different maturation conditions. As seen in PCA-score plots (Figure. 1. C), IMH samples separate from MH samples, indicating a large difference between the two groups of samples. A volcano plot representing the filtered data is shown in Figure. 1. B The compounds found at significantly ($P < 0.05$) higher levels in IMH than MH samples were organic acids (benzoic acid, linalool, sinapic acid and ganoderic acid *etc.*), alcohols, some derivatives of acids (ethyl gallate, levistilide), some glycosides, plant alkaloids, and very small amounts of phenolic compounds such as vanilic acid and eugenol. The compounds found at significantly ($P < 0.05$) lower levels were caffeic acid, 3, 4-dihydroxybenzoic acid, chlorogenic acid and common phenolic substances.

To further understand the material differences between the two kinds of honey, we chose to analyze the polyphenols that are major active ingredients in honey. A total of 29 types of flavonoids and phenolic acids were studied of which 20 were detected. These compounds were selected as they were predominately the active constituents in honey as well as propolis. We have previously established accurate quantification methods for these chemicals.^{14,34} Average concentrations of these polyphenolic

compounds varied greatly among samples (between 0.38 ± 0.01 to 158.09 ± 2.89 $\mu\text{g}/100$ g honey). Six of these compounds were only detected in MH, including kaempferol, apigenin, pinocembrin, 3-(3, 4-dimethoxyphenyl)-2-propenoic acid, chrysin and caffeic acid phenethyl ester. Interestingly, with the exceptions of vanillic acid and syringic acid, the content of the remaining 12 substances in MH were significantly higher than those in IMH. These polyphenols are derived from plants, and are known as the key contributors to the honey's color and taste, as well as its biological activities.³⁵ Of course, phenols can vary depending on nectar plant, bee species and geographic source.^{36,37} It is well documented that plant phenolic metabolites change by the action of bee enzymes in honey. The results from the present study show significant changes, which might be due to interaction with some substances in the beehive, like hydrolysis from glycosides to give rise to aglycones. Nevertheless, we only analyzed honey of unifloral origin and it would be interesting to further compare differences between mature and immature honeys from other nectar sources and over different seasons.

The anti-microbial activity of honey is clearly established and honey could provide a potential alternative to antibiotics.³⁹ The possible underlying mechanism of action relies on the ability of honey to generate hydrogen peroxide by the bee-derived enzyme glucose oxidase.^{40,41} However, other factors may also contribute to its antimicrobial activity such as high osmotic pressure, acidic environment, low protein content, high carbon to nitrogen ratio, low redox potential (due to the high level of reducing sugars), and a level of viscosity that limits dissolved oxygen and other chemical

agents/phytochemicals. Another potential contributor is the complex composition of honey, which has more than 181 constituents.⁴² These include terpenes, pinocembrin, benzyl alcohol, 3, 5-dimethoxy-4-hydroxybenzoic acid (syringic acid), methyl-3, 5-dimethoxy-4-hydroxybenzoate (methyl syringate), 2-hydroxy-3-phenylpropionic acid, 2-hydroxybenzoic acid, 3, 4, 5-trimethoxybenzoic acid, and 1, 4-dihydroxybenzene. Consistent with previous studies, we found that honey exhibits a bacteriostatic against several pathogens, such as *Staphylococcus aureus* and *Escherichia coli*. *E. coli* is a Gram-negative bacterium that is pathogenic to human and animals and can cause diarrhea and sepsis in children, travelers, piglets and chickens.⁴³ *S. aureus* is a representative of gram-positive bacteria and infection can cause serious illness in humans.⁴⁴ Our results showed that the zones of inhibition of *E.coli* and *S. aureus* by mature rapeseed honey were bigger than the immature rapeseed honeys, thus demonstrating that MH has a stronger bacteriostatic effect than IMH. Nevertheless, both samples had no obvious inhibitory effect on *Bacillus subtilis*. *B. subtilis* is a multifunctional probiotic and is beneficial for human digestion and absorption. It produces subtilisin, polymyxin and other active substances to inhibit intestinal pathogenic bacteria.⁴⁵

Honey works as an abundant source of natural anti-oxidants which play an important role in food preservation and human health.¹² Anti-oxidant substances have different mechanisms, such as reducing the damaging effects of reactive oxygen and reactive nitrogen species, inhibiting the effects of enzymes that produce superoxide anions, promoting metal chelation and free radical chain reaction, and inhibiting the

414 formation of active oxidants.³⁷ In the present study, three standard spectrophotometric
415 methods are used for comparing the *in vitro* anti-oxidant effects of MH and IMH
416 samples: The DPPH test and ABTS⁺ test for radical scavenging activity and the Ferric
417 reducing ability of plasma (FRAP) method for their reducing power.¹⁵ The main anti-
418 oxidants in honeys are polyphenols, including phenolic acids and flavonoids. According
419 to previous studies, the total phenolic content of honey is uncertain, ranging from 0.205
420 mg GAE/g to 1.877 mg GAE/g honey, while among rapeseed honey, it ranges from
421 0.205 mg GAE/g to 0.311 mg GAE/g honey.^{15,40} In this study, we studied the total
422 phenolic content of honey extracts, producing results of 12.99 ± 0.19 and 12.20 ± 0.16
423 mg GAE/g honey extracts for MH and IMH, respectively. Although these values are
424 fall within a certain range with previous literature, our results are significantly higher
425 than previously published data.⁴⁶ An explanation for this may be that we extracted the
426 honey before testing it. The content of total phenols in the honey polyphenol extract of
427 mature rapeseed honey (12.99 ± 0.19 mg GAE/g extract) was significantly higher than
428 that of immature rapeseed honey (12.20 ± 0.16 mg GAE/g extract). However, there was
429 no significant difference in the content of total flavonoids. Rapeseed honey from
430 different geographical sources has been shown to possess different anti-oxidant
431 capacities. Piotr Marek Kuś et al.⁴⁷ studied the anti-oxidant capacity of 10 kinds of
432 rapeseed honey from 8 regions in Poland, finding that the FRAP level was 1.0-1.8
433 (mmol Fe²⁺/kg), and the average level was 1.3 ± 0.3 . DPPH level was 0.3-0.5 (mmol
434 TEAC/kg), average level was 0.4 ± 0.1 (mmol TEAC/kg). The FRAP and DPPH values
435 of MH and IMH samples in our study were smaller than Piotr Marek Kuś et al reported,

but the MH has a stronger anti-oxidant activity than the IMH.

Honey has a regulatory effect on cell growth and proliferation, metabolism and anti-oxidant enzymes, and has a protective effect on cell damage caused by adverse stimulation.¹¹ The mechanisms by which honey influences the biological activity of cells is complex.^{49,50} In this study, a cell oxidative stress model was applied in mouse fibroblasts (L929) stimulated by hydrogen peroxide as previously established.⁵¹ Firstly, the concentration of H₂O₂ was determined by toxicity testing, as shown in the figure 3. The reproductivity of the cultured cells can be significantly reduced when treated with 500 µM H₂O₂, but the honey extract had no toxic effect on cells in the range of tested concentrations. Then cells were pretreated with honey extract prior to 500 µM H₂O₂ treatment. Our results demonstrated that the honey extract could significantly improve the cell growth activity. MH showed a positive effect at the concentration of 400 µg/mL, while for IMH is the required concentration was 600 µg/mL. This suggests that honey can counteract the cell damage caused by oxidative stress, with the effect of mature honey more potent.

In the meantime, we examined the expression of antioxidant genes (*HO-1*, *TXNRD*, *GCLM*, and *NQO1*) in cells. The results showed significantly increased expression of anti-oxidant genes in the MH-pretreated cells. However, the effect of IMH was weak and had no significant effect on *NQO1* gene expression. Heme oxygenase 1 (HO-1), catalyzes the decomposition of heme into a series of anti-oxidant and anti-inflammatory molecules that prevent oxidation;⁵² NQO1 catalyzes double electron reduction to reduce oxidative damage;⁵³ GCLM is a subunit of glutamic acid and cysteine synthase,

the most important genes in the cellular anti-oxidant defense mechanism;⁵⁴ TXNRD (thioredoxin reductase) is involved in many redox reactions in vivo.⁵⁵ These anti-oxidant genes are important regulators of NRF-2 signaling pathway.⁵⁶ The NRF-2 signaling pathway regulates the transcriptional expression of many proteins with detoxification and anti-oxidant defense functions. Our results suggest that honey may affect the cellular oxidative stress response by affecting the NRF-2 signaling pathway.

This study performed analysis of mature and immature honey using untargeted and targeted methods, and determined their anti-bacterial and anti-oxidant activity *in vitro*. The results demonstrated that the harvest of honey before the maturity stage can have profound impacts upon its quality. Our study demonstrated using metabolomics data analysis the possibility to that mature honey and immature honey could be distinguished by the metabolite differences between them by means of metabolomics data analysis. Untargeted substance analysis based on Mass Profiler Professional software explains the difference between the two from a macro perspective. Further in-depth analysis of target substance research indicates that effective and beneficial substances are more abundant in mature honey than in immature honey. This is the first time that metabolomics analysis technology was applied to the study of honey quality. Results from in vitro anti-bacterial and anti-oxidant experiments showed that mature capped honey is more effective in inhibiting proliferation of *E. coli* and *S. aureus*, and may protect mice skin fibroblast L929 cells from the damage of free radicals by enhancing the expression of anti-oxidant related genes after H₂O₂ stimulation. In conclusion, mature honey has a greater value.

Acknowledgement

We wish to thank Mr. Ran Liu for his kind assistance during honey sample collection process.

Funding information

This work was supported by the National Natural Science Foundation of China under Grant (31972628); The Agricultural Science and Technology Innovation Program under Grant (CAAS-ASTIP-2019-IAR), and the earmarked fund for Modern Agroindustry Technology Research System from the Ministry of Agriculture of China under Grant (CARS-44).

Conflict of interest

The authors declare that they have no conflict of interest.

Supporting Information.

Supplemental Table 1: Sequences of the primers used for qRT-PCR

Reference

1. Finola, M. S.; Lasagno, M. C.; Marioli, J. M., Microbiological and chemical characterization of honeys from central Argentina. *Food Chemistry*. **2007**, *100*(4), 1649-1653.
2. Eyer, M.; Neumann, P.; Dietemann, V., A look into the cell: honey storage in honey bees, *apis mellifera*. *PloS one*. **2016**, *11*(8), e0161059.
3. Zhang, C.; Pokhrel, S.; Wu, Z. H.; Miao, X. Q.; Huang, Z. Y.; Yang, W. Y., Longevity, food consumption, and foraging performance of *Apis cerana* and *Apis mellifera* in mixed colonies. *Apidologie*. **2019**, *50*(2), 153-162.
4. García, N. L., The current situation on the international honey market. *Bee World*. **2018**, *95*(3), 89-94.
5. Olofsson, T.; Vásquez, A., Honeybee-Specific Bifidobacteria and Lactobacilli, The Bifidobacteria and Related Organisms. *Academic Press*. **2018**, 235-241.
6. Murilhas, A. M., Varroa destructor infestation impact on *Apis mellifera carnica* capped worker brood production, bee population and honey storage in a Mediterranean climate. *Apidologie*. **2002**, *33*(3), 271-281.
7. Bromenshenk, J. J.; Gudatis, J. L.; Carlson, S. R.; Thomasb, J.M.; Simmonsb, M.A., Population dynamics of honey bee nucleus colonies exposed to industrial pollutants. *Apidologie*. **1991**, *22*(4), 359-369.
8. Locher, C.; Neumann, J.; Sostaric, T., Authentication of honeys of different floral origins via high-performance thin-layer chromatographic fingerprinting. *JPC-Journal of Planar Chromatography-Modern TLC*. **2017**, *30*(1), 57-62.

- 518 9. Melucci, D.; Zappi, A.; Bolelli, L.; Corvucci, F.; Serra, G.; Boi, M.; Grillenzoni, F.
519 V.; Fedrizzi, G.; Menotta, S.; Girotti, S., Checking syrup adulteration of honey using
520 bioluminescent bacteria and chemometrics. *European Food Research and Technology*.
521 **2019**, 245(2), 315-324.
- 522 10. Bougrini, M.; Tahri, K.; Saidi, T.; El Alami El Hassani, N.; Bouchikhi, B.; El Bari,
523 N., Classification of Honey According to Geographical and Botanical Origins and
524 Detection of Its Adulteration Using Voltammetric Electronic Tongue. *Food Analytical*
525 *Methods*. **2016**, 9(8), 2161-2173.
- 526 11. Semkiw, P.; Skowronek, W.; Skubida, P., Changes in water content of honey during
527 ripening under controlled condition. *Journal of Apicultural Science*. **2008**, 52(1), 57-
528 63.
- 529 12. Meo, S. A.; Al-Asiri, S. A.; Mahesar, A. L.; Javed, M. J., Role of honey in modern
530 medicine. *Saudi journal of biological sciences*. **2017**, 24(5), 975-978.
- 531 13. Alvarez-Suarez, J.; Gasparrini, M.; Forbes-Hernández, T.; Mazzoni, L.; Giampieri,
532 F., The Composition and Biological Activity of Honey: A Focus on Manuka Honey.
533 *Foods*. **2014**, 3(3), 420-432.
- 534 14. Wang, K.; Wan, Z. R.; Ou, A. Q.; Liang, X. W.; Guo, X. W.; Zhang, Z. Y.; Wu, L.
535 M.; Xue, X. F., Monofloral honey from a medical plant, *Prunella Vulgaris*, protected
536 against dextran sulfate sodium-induced ulcerative colitis via modulating gut microbial
537 populations in rats. *Food & function*. **2019**.
- 538 15. Vasić, V.; Gašić, U.; Stanković, D.; Lušić, D.; Vukić-Lušić, D.; Milojković-Opsenica,
539 D.; Tešića, Ž.; Trifkovića, J., Towards better quality criteria of European honeydew

- 540 honey: Phenolic profile and antioxidant capacity. *Food chemistry*. **2019**, 274, 629-641.
- 541 16. De-Melo, A. A. M.; Almeida-Muradian, L. B. D.; Sancho, M. T.; Pascual-Maté, A.,
542 Composition and properties of *Apis mellifera* honey: A review. *Journal of Apicultural*
543 *Research*. **2018**, 57(1), 5-37.
- 544 17. Ma, T. C.; Zhao, H. A.; Liu, C. Y.; Zhu, M.; Gao, H.; Cheng, N.; Cao, W.,
545 Discrimination of Natural Mature Acacia Honey Based on Multi-Physicochemical
546 Parameters Combined with Chemometric Analysis. *Molecules*. **2019**, 24(14), 2674.
- 547 18. Nicolson, S. W.; Human, H., Bees get a head start on honey production. *Biology*
548 *Letters*. **2008**, 4(3), 299-301.
- 549 19. Kamal, M. M.; Rashid, M. H. U.; Mondal, S. C.; El Taj, H. F.; Jung, C.,
550 Physicochemical and microbiological characteristics of honey obtained through sugar
551 feeding of bees. *Journal of Food Science and Technology*. **2019**, 56 (4), 2267-2277.
- 552 20. Iqbal, S.; Sukhmeet, S., Honey moisture reduction and its quality. *Journal of Food*
553 *Science & Technology*. **2018**, 55(10), 3861-3871.
- 554 21. Al-Farsi, M.; Al-Belushi, S.; Al-Amri, A.; Al-Hadhrami, A.; Al-Rusheidi, M.; Al-
555 Alawi, A., Quality evaluation of Omani honey. *Food Chemistry*. **2018**, 262, 162-167.
- 556 22. Geană, E. I.; Ciucure, C. T.; Costinel, D.; Ionete, R. E., Evaluation of honey in terms
557 of quality and authenticity based on the general physicochemical pattern, major sugar
558 composition and $\delta^{13}\text{C}$ signature. *Food Control*. **2020**, 109, 106919.
- 559 23. Council, E. U., Council Directive 2001/110/EC of 20 December 2001 relating to
560 honey. *Official Journal of the European Communities L*. **2002**, 10, 47-52.
- 561 24. Wu, L.; Du, B.; Vander Heyden, Y.; Chen, L.; Zhao, L.; Wang, M.; Xue, X., Recent

562 advancements in detecting sugar-based adulterants in honey – A challenge. *Trac Trends*
563 *in Analytical Chemistry*. **2017**, 86, 25-38.

564 25. Mouhoubi-Tafnine, Z.; Ouchemoukh, S.; Louaileche, H.; Tamendjari, A., Effect of
565 storage on hydroxymethylfurfural (HMF) and color of some Algerian honey.
566 *International Food Research Journal*. **2018**, 25(3).

567 26. Kuropatnicki, A. K.; Kłósek, M.; Kucharzewski, M., Honey as medicine: historical
568 perspectives. *Journal of Apicultural Research*. **2018**, 57(1), 113-118.

569 27. Michael, E.; Peter, N.; Vincent, D.; Olav, R., A Look into the Cell: Honey Storage
570 in Honey Bees, *Apis mellifera*. *Plos One*. **2016**, 11(8), e0161059.

571 28. Mart N, I. G.; As, E. M. M.; Sánchez, J. S.; Rivera, B. G., Detection of honey
572 adulteration with beet sugar using stable isotope methodology. *Food Chemistry*. **1998**,
573 61(3), 281-286.

574 29. Hong, E.; Lee, S. Y.; Jeong, J. Y.; Park, J. M.; Kim, B. H.; Kwon, K.; Chun, H. S.,
575 Modern Analytical Methods for the Detection of Food Fraud and Adulteration by Food
576 Category. *Journal of the Science of Food & Agriculture*. **2017**, 97(12), 3877-3896.

577 30. Gao, W.; Yang, H.; Qi, L.; Liu, E.; Ren, M.; Yan, Y.; Chen, J.; Li, P., Unbiased
578 metabolite profiling by liquid chromatography–quadrupole time-of-flight mass
579 spectrometry and multivariate data analysis for herbal authentication: Classification of
580 seven *Lonicera* species flower buds. *Journal of Chromatography A*. **2012**, 1245: 109-
581 116.

582 31. Li, T.; Shuai, W.; Meng, X.; Bao, Y.; Guan, S.; Bo, L.; Lu, C.; Lei, W.; Ran, X.,
583 Metabolomics Coupled with Multivariate Data and Pathway Analysis on Potential

584 Biomarkers in Gastric Ulcer and Intervention Effects of *Corydalis yanhusuo* Alkaloid.
 585 *Plos One*. **2014**, 9.

586 32. Gheldof, N.; Wang, X. H.; Engeseth, N. J., Identification and Quantification of
 587 Antioxidant Components of Honeys from Various Floral Sources. *Journal of*
 588 *Agricultural & Food Chemistry*. **2002**, 50(21), 5870-5877.

589 33. Samborska, K.; Jedlińska, A.; Wiktor, A.; Derewiaka, D.; Wołosiak, R.; Arkadiusz
 590 Matwiczuk, A.; Jamróz, W.; Skwarczyńska-Maj, K.; Kiełczewski, D.; Błażowski,
 591 Ł.; Tułodziecki, M.; Rajchert, D. W., The Effect of Low-Temperature Spray Drying
 592 with Dehumidified Air on Phenolic Compounds, Antioxidant Activity, and Aroma
 593 Compounds of Rapeseed Honey Powders. *Food and Bioprocess Technology*. **2019**,
 594 12(6), 919-932.

595 34. Jin, X. L.; Wang, K.; Li, Q. Q.; Tian, W. L.; Xue, X. F.; Wu, L. M.; Hu, F. L.,
 596 Antioxidant and anti-inflammatory effects of Chinese propolis during palmitic acid-
 597 induced lipotoxicity in cultured hepatocytes. *Journal of Functional Foods*. **2017**, 34,
 598 216-223.

599 35. Marco, C.; Nadia, S.; Maria, P.; Gavino, S., Recent advances in the analysis of
 600 phenolic compounds in unifloral honeys. *Molecules*. **2016**, 21(4), 451.

601 36. Alvarez-Suarez, J. M.; Giampieri, F.; Brenciani, A. Mazzoni, L.; Gasparrini, M.;
 602 González-Paramás, A. M.; Santos-Buelga, C.; Morroni, G.; Simoni, S.; Forbes-
 603 Hernández, T. Y.; Afrin, S.; Giovanetti, E.; Battino, M., *Apis mellifera* vs *Melipona*
 604 *beecheii* Cuban polyfloral honeys: A comparison based on their physicochemical
 605 parameters, chemical composition and biological properties. *LWT - Food Science and*

606 *Technology*. **2018**, 87, 272-279.

607 37. Machado De-Melo, A. A.; Almeida-Muradian, L. B.; Sancho, M. T.; Pascual-Maté,
608 A., Composition and properties of *Apis mellifera* honey: A review. *Journal of*
609 *Apicultural Research*. **2018**, 57(1), 5-37.

610 38. Zhu, Z. Y.; Zhang, Y.; Wang, J. H.; Li, X.; Wang, W.; Huang, Z. P., Sugaring-out
611 assisted liquid-liquid extraction coupled with high performance liquid chromatography-
612 electrochemical detection for the determination of 17 phenolic compounds in honey.
613 *Journal of Chromatography A*. **2019**, 1601, 104-114.

614 39. Samarghandian, S.; Farkhondeh, T.; Samini, F., Honey and health: A review of
615 recent clinical research. *Pharmacognosy research*. **2017**, 9(2), 121.

616 40. Girma, A.; Seo, W.; She, R. C., Antibacterial activity of varying UMF-graded
617 Manuka honeys. *Plos One*. 2019, 14(10).

618 41. Sindi, A.; Chawn, M. V. B.; Hernandez, M. E.; Green, K.; Islam, M. K.; Locher,
619 C.; Hammer, K., Anti-biofilm effects and characterisation of the hydrogen peroxide
620 activity of a range of Western Australian honeys compared to Manuka and multifloral
621 honeys. *Scientific Reports*. 2019, 9(1), 1-17.

622 42. Vallianou, N. G.; Gounari, P.; Skourtis, A.; Panagos, J.; Kazazis, C., Honey and its
623 anti-inflammatory, anti-bacterial and anti-oxidant properties. *General Medicine: Open*
624 *Access*. **2014**, 2(132): 1-5.

625 43. Pontrelli, S.; Chiu, T. Y.; Lan, E. I.; Chen, F.; Chang, P.; Liao, J., Escherichia coli
626 as a host for metabolic engineering. *Metabolic Engineering*. **2018**, 50: 16-46.

627 44. Lakhundi, S.; Zhang, K., Methicillin-resistant Staphylococcus aureus: molecular

628 characterization, evolution, and epidemiology. *Clinical microbiology reviews*. **2018**,
629 *31*(4): e00020-18.

630 45. Suva, M. A.; Sureja, V. P.; Kheni, D. B., Novel insight on probiotic *Bacillus subtilis*:
631 mechanism of action and clinical applications. *Journal of Current Research in Scientific*
632 *Medicine*. **2016**, *2*(2): 65.

633 46. Dżugan, M.; Tomczyk, M.; Sowa, P.; Grabek-Lejko, D., Antioxidant activity as
634 biomarker of honey variety. *Molecules*. **2018**, *23*(8): 2069.

635 47. Kuś, P. M.; Congiu, F.; Teper, D.; Sroka, Z.; Jerković, I.; Tuberoso, C. I. G.,
636 Antioxidant activity, color characteristics, total phenol content and general HPLC
637 fingerprints of six Polish unifloral honey types. *LWT - Food Science and Technology*.
638 **2014**, *55*(1), 124-130.

639 48. Minden-Birkenmaier, B. A.; Meadows, M. B.; Cherukuri, K.; Smelthew, M. P.;
640 Smith, R. A.; Radic, M. Z.; Bowlin, G. L., The Effect of Manuka Honey on dHL-60
641 Cytokine, Chemokine, and Matrix-Degrading Enzyme Release under Inflammatory
642 Conditions. *Med One*. **2019**, *4*(2).

643 49. Afrin, S.; Giampieri, F.; Gasparrini, M.; Forbes-Hernández, T. Y.; Cianciosi, D.;
644 Reboredo-Rodriguez, P.; Amici, A.; Quiles, J. L.; Battino, M., The inhibitory effect of
645 Manuka honey on human colon cancer HCT-116 and LoVo cell growth. Part 1: the
646 suppression of cell proliferation, promotion of apoptosis and arrest of the cell cycle.
647 *Food and Function*. **2018**, *9*(4): 2145-2157.

648 50. Afrin, S.; Giampieri, F.; Gasparrini, M.; Forbes-Hernández, T. Y.; Cianciosi, D.;
649 Reboredo-Rodriguez, P.; Manna, P. P.; Zhang, J. J.; A.; Quiles, J. L.; Battino, M., The

inhibitory effect of Manuka honey on human colon cancer HCT-116 and LoVo cell growth. Part 2: Induction of oxidative stress, alteration of mitochondrial respiration and glycolysis, and suppression of metastatic ability. *Food and Function*. **2018**, 9(4), 2158-2170.

51. Cao, X. P.; Chen, Y.; Zhang, J.; You, M.; Wang, K.; Hu, F., Mechanisms underlying the wound healing potential of propolis based on its in vitro antioxidant activity. *Phytomedicine*. **2017**, 34, 76-84.

52. Hseu, Y. C.; Chou, C.; Kumar, K. J. S.; Fu, K.; Wang, H.; Hsu, L.; Kuo, Y.; Wu, C.; Chen, S.; Yang, H., Ellagic acid protects human keratinocyte (HaCaT) cells against UVA-induced oxidative stress and apoptosis through the upregulation of the HO-1 and Nrf-2 antioxidant genes. *Food and Chemical Toxicology*. **2012**, 50 (5), 1245-1255.

53. Luo, S.; Lei, K.; Xiang, D.; Ye, K., NQO1 Is Regulated by PTEN in Glioblastoma, Mediating Cell Proliferation and Oxidative Stress. *Oxidative Medicine & Cellular Longevity*. **2018**.

54. Jinhwan, L.; N., N. B.; Isaac, M.; J., K. T.; Ulrike, L., Glutamate Cysteine Ligase Modifier Subunit (Gclm) Null Mice Have Increased Ovarian Oxidative Stress and Accelerated Age-Related Ovarian Failure. *Endocrinology*. **2015**, (9), 9.

55. Ingold, I.; Conrad, M., Oxidative Stress, Selenium Redox Systems Including GPX/TXNRD Families. *Selenium. Springer, Cham*. **2018**, 111-135.

56. Mohammadzadeh, M.; Halabian, R.; Gharehbaghian, A.; Amirizadeh, N.; Jahanian-Najafabadi, A.; Roudkenar, A. M. R. A. Nrf-2 overexpression in mesenchymal stem cells reduces oxidative stress-induced apoptosis and cytotoxicity. *Cell Stress and*

672 *Chaperones*. **2012**, 17 (5), 553-565.

673

Figure Captions

Figure 1 Discrimination of mature capped honey and immature uncapped honey (MH represents mature honey; IMH represents immature honey). A: Venn diagram of untargeted analysis of MH and IMH with a filtration of samples frequency (70%). The number in the picture represents the number of species of matter. B: Volcano plot of the honey different metabolites for group MH vs IMH ($P < 0.05$). C: PCA scores plot of MH and IMH.

Figure 2 Total ion chromatography of honey extracts with negative scanning mode in HPLC-Q-TOF-MS. Red line represents mature honey (MH); Green line represents immature honey (IMH). The samples for 0-1 min are discarded without mass spectrometry.

Figure 3 Effect of H_2O_2 and honey extracts on L929 cells viability. (A). Cells were pretreated with/without the indicated concentrations of H_2O_2 (300 μ M-600 μ M) and honey extracts (0 μ g/mL-600 μ g/mL) for 24 h. (B)(C). Cells were pretreated with/without the different concentrations of MHE/IMHE for 2 h and then stimulated with 500 μ M H_2O_2 for 24 h. ★ indicates the control group for significance analysis. Each result was expressed as the mean \pm SD ($n = 3$); * $P < 0.05$ versus the control group (★); ** $P < 0.01$ versus the control group (★); *** $P < 0.001$ versus the control group (★).

Figure 4 Effect of honey extracts on the expression of antioxidant related genes in H_2O_2 stimulated cells. L929 cells were pretreated with or without the indicated concentrations of MHE/IMHE for 2 h and were then stimulated with 500 μ M H_2O_2 for 6 h. The relative mRNA expression of *HO-1*(A), *TXNRD* (B), *GCLM* (C) and *NQO1* (D) were

696 determined using qRT-PCR. Each result was shown as the mean \pm SD (n = 3). **P <
697 0.01 versus the untreated group (★), ***P < 0.001 versus the untreated group (★).
698




Tables






Table 1 Parameters of Mature and Immature Honey^{*}.






Parameter	MH	IMH
Fructose%	36.40± 0.37 ^a	30.86 ± 0.64 ^b
Glucose%	34.49± 2.17 ^a	30.14 ± 0.72 ^a
Sucrose%	2.33 ± 0.17 ^a	1.11 ± 0.73 ^a
Water%	18.31 ± 1.52 ^a	31.20 ± 1.81 ^b
Acidity		
meq/kg	13.67 ± 1.88 ^a	19.9 ± 0.42 ^b
HMF	ND	






^{*} In each column, different letters (a, b) mean significant differences ($p < 0.05$). ND means not detected.



Table 2 HPLC-Q-TOF MS Analysis of Major Phenolic Compounds and Relative Occurrence in MH and IMH*.

Compounds	Molecular formula	Molecular weight	Molecular formulae	[M-H] ⁻ , <i>m/z</i>	RT/min	μg/100gMH	μg/100gIMH	LOD μg/100g	R ²
3,4-Dihydroxybenzoic acid	C ₇ H ₆ O ₄	154.12		153.0193	4.596	7.20±0.05 ^a	3.05±0.05 ^b	0.025	0.999
Chlorogenic acid	C ₁₆ H ₁₈ O ₉	354.31		353.0878	5.510	11.70±0.20 ^a	4.80±0.09 ^b	0.006	0.990
Caffeic acid	C ₉ H ₈ O ₄	180.16		179.0350	6.035	22.86±0.88 ^a	5.37±0.05 ^b	0.023	0.992

Vanillic acid	C ₈ H ₈ O ₄	168.15		167.0350	5.950	51.17±0.58 ^b	60.10±0.25 ^a	0.250	0.99 5
Syringic acid	C ₉ H ₁₀ O ₅	198.17		197.0455	6.169	6.81±0.04 ^b	14.68±0.12 ^a	0.147	0.99 0
p-Coumaric acid	C ₉ H ₈ O ₃	164.16		163.0401	6.804	9.51±0.52 ^a	1.38±0.04 ^b	0.013	0.99 0
Ferulic Acid	C ₁₀ H ₁₀ O ₄	194.18		193.0506	6.950	19.84±1.10 ^a	4.36±0.13 ^b	0.043	0.99 8
3-O-Acetylpinobanksin	C ₁₇ H ₁₄ O ₆	314.00		313.0718	11.561	13.53±0.40 ^a	0.38±0.01 ^b	0.008	0.99 4

Quercetin	C ₁₅ H ₁₀ O ₇	302.24		301.0354	8.743	134.16±2.41 ^a	58.94±0.63 ^b	0.057	0.99	1
Hesperitin	C ₁₆ H ₁₄ O ₆	302.28		301.0718	8.939	158.09±2.89 ^a	76.32±0.83 ^b	0.012	0.99	1
Pinobanksin	C ₁₅ H ₁₂ O ₅	272.25		271.0612	8.928	25.74±0.20 ^a	0.84±0.01 ^b	0.027	0.99	3
Naringenin	C ₁₅ H ₁₂ O ₅	272.25		271.0612	9.098	23.39±0.33 ^a	0.88±0.01 ^b	0.027	0.99	3
Galangin	C ₁₅ H ₁₀ O ₅	270.24		269.0455	9.122	8.42±0.24 ^a	3.63±0.01 ^b	0.018	0.99	0

Luteolin	C ₁₅ H ₁₀ O ₆	286.24		285.0405	9.952	67.32±1.09 ^a	34.82±0.57 ^b	0.062	0.99 0
Kaempferol	C ₁₅ H ₁₀ O ₆	286.24		285.0405	11.391	25.44±0.46	ND	0.078	0.99 8
Apigenin	C ₁₅ H ₁₀ O ₅	270.24		269.0455	7.792	15.24±0.19	ND	0.039	0.99 3
Pinocembrin	C ₁₅ H ₁₂ O ₄	256.25		255.0663	12.037	10.95±0.31	ND	0.036	0.99 2
3-(3,4-Dimethoxyphenyl)-2-propenoic Acid	C ₁₁ H ₁₂ O ₄	208.21		207.0663	12.184	19.60±0.23	ND	0.166	0.99 1

Chrysin	C ₁₅ H ₁₀ O ₄	254.24		253.0506	12.612	18.83±0.97	ND	0.011	0.99
									8
Caffeic acid phenethyl ester	C ₁₇ H ₁₆ O ₄	284.31		283.0976	9.793	12.89±0.26	ND	0.025	0.99
									8

* Detected in negative ionization mode. In each column different letters (a, b) mean significant differences ($p < 0.05$). ND means not detected.

Table 3 Zones of inhibition of MH and IMH*.

	Zones of inhibition (mm)				
	Ampicillin(5µg/mL)	Phenol(10%)	50%MH	50%IMH	Water
<i>E.coli</i>	17.69±0.43	15.40±0.57	19.47 ^a ±0.31	17.29 ^b ±0.78	--
<i>S.aureus</i>	29.23±0.62	14.40±0.69	14.13 ^a ±0.68	12.8 ^b ±0.98	--
<i>B.subtilis</i>	0.50±0.08	20.90±0.29	--	--	--

* In each column different letters (a, b) mean significant differences ($p < 0.05$). -- means that there is no observed bacteriostatic zone.

Table 4 Antioxidant Activity of Mature Honey and Immature Honey. Including Radical Scavenging Capacity, Reducing Power, and Total Phenols and Flavone of MH and IMH*.

	FRAP		ABTS		DPPH		Total phenols	Total flavonoids
	IC50	mg Trolox	IC50		IC50			
	mg/mL	/g	mg/mL	mgTrolox/g	mg/mL	mgTrolox/g	mgGAE/g	mgQE/g
MH	1.69±0.02	36.97±0.53 ^a	0.86±0.02	37.82±0.90 ^a	2.26±0.01	21.89±0.08 ^a	12.99±0.19 ^a	3.53±0.07 ^a
IMH	2.21±0.06	28.41±0.76 ^b	1.01±0.06	32.30±0.81 ^b	2.53±0.05	19.60±0.36 ^b	12.20±0.16 ^b	3.41±0.01 ^a
Vc (µg/mL)	35.07±0.02		16.76±0.06		28.01±0.02			

* In each column different letters (a, b) mean significant differences ($p < 0.05$). IC50 means the sample concentration providing 0.5 of absorbance was determined by a linear curve established by mass concentration and absorbance. Meanwhile all the results were expressed as equivalent of the corresponding standard reference (mg Trolox equalation per gram (mg Trolox/g); mg gallic acid equivalent per gram (mg GAE/g); mg quercetin equivalent per gram (mg QE/g)).

Figure graphics

Figure 1




Figure 2




Figure 3




Figure 4




Table of Contents (TOC) Art

