



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Avramidis, S and McKenna, J and Long, JA and Butterly, R and Llewellyn, DJ (2010) Application for the 4W Model of Drowning for Prevention, Rescue and Treatment, Research and Education.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/674/>

Document Version:

Article (Accepted Version)

---

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

## Applications of the 4W Model of Drowning for Prevention, Rescue and Treatment, Research and Education

<sup>1,2,3</sup>Stathis Avramidis, <sup>2</sup>Jim McKenna, <sup>2</sup>Jonathan Long, <sup>2</sup>Ronald Butterly, & <sup>4</sup>David J. Llewellyn

<sup>1</sup>Hellenic Center for Disease Control and Prevention, <sup>2</sup>Leeds Metropolitan University, <sup>3</sup>Lifesaving Foundation, <sup>4</sup>University of Cambridge

Previous research has been published about the 4W model of drowning and its four constituent variables (Avramidis, Butterly & Llewellyn, 2007; 2009a; 2009b; 2009c; 2009d; Avramidis, McKenna, Long, Butterly, & Llewellyn, 2010). We presently summarize and suggest applications of the model for the general public, aquatic safety professionals, injury epidemiologists and policy makers. The method consisted of a major literature review of quantitative research that was undertaken to identify potential risk factors of drowning, a qualitative content analysis that was used to analyze publicly available drowning incident videos ( $n = 41$ ,  $M = 345.0$  secs,  $SD = 2.8$ ), and a semi-structured interview of 34 individuals who were involved in drowning incidents (30 males age 16–65 years,  $M = 28.4$ ,  $SD = 11.3$ ; 4 female age 19–65 years,  $M = 37.5$ ,  $SD = 19.5$ ). Results confirm that the model has numerous applications in terms of prevention, rescue, treatment, research and education. In terms of prevention and education, it is suggested that water safety organizations should establish new testing criteria for qualifying pool and beach lifeguards (e.g. “100 m run–50 m swim–100 m run” for open water, “50 m run–20 m swim–50 m run” for pool/water parks and “early approach” criteria). In addition, emphasis needs to be given during training on the 3 dimensions that constitute each drowning problem (i.e. width, length and height or depth). Moreover, this stresses the need, from an educational point of view, for better public awareness regarding water safety prevention in people who engage not only in aquatic activities but also in non-aquatics in or around the water, for them to know how to swim and be able to survive in an aquatic emergency. In terms of rescue and treatment, consideration of the 3 dimensions that synthesize the rescue and its implications for the outcome of the rescue and the first aid treatment will help lifeguards and lifesavers to be faster, more effective and able to avoid potential problems that might delay their attempt. Finally, in terms of research, the 3 dimensions of the model reveal that the numerous “hidden” drowning incidents that contemporary injury epidemiology classifies under different codes, should count as “drownings” for better describing the real mortality rates. By doing this, the increased rate of the drowning figure will act as a positive force that will positively influence decision making, research funding, public awareness and lifeguard preparedness. Collectively, the 4W model has a wide range of lifesaving related applications.

## References

- Avramidis, S., Butterly, R., & Llewellyn, D.J. (2007). The 4W Model of Drowning. *International Journal of Aquatic Research and Education*, 1(3), 221-230.
- Avramidis, S., Butterly, R., & Llewellyn, D.J. (2009a). Drowning Incident Rescuer Characteristics: Encoding the First Component of the 4W Model. *International Journal of Aquatic Research and Education*, 3(1), 66-82.
- Avramidis, S., Butterly, R., & Llewellyn, D.J. (2009b). Who Drowns? Encoding the Second Component of the 4W Model. *International Journal of Aquatic Research and Education*, 3(3), 224-235.
- Avramidis, S., Butterly, R., & Llewellyn, D.J. (2009c). Where do People Drown? Encoding the Third Component of the 4W Model. *International Journal of Aquatic Research and Education*, 3(3), 236-254.
- Avramidis, S., Butterly, R., & Llewellyn, D.J. (2009d). Under What Circumstances Do People Drown? Encoding the Fourth Component of the 4W Model. *International Journal of Aquatic Research and Education*, 3(4), 406-421.
- Avramidis, S., McKenna, J., Long, J., Butterly, R. & Llewellyn, D. (2010). Crucial Findings from the 4W Model of Drowning for Practical and Teaching Applications. In P-L. Kjendlie, R.K. Stallman, & J. Cabri, (Eds.), *Proceedings of the 11<sup>th</sup> International Symposium for Biomechanics and Medicine in Swimming 2010* (pp. 354-356). Oslo: Norwegian School of Sport Sciences.