
Citation:

Morris, R and Myers, T and Emmonds, S and Singleton, D and Till, K (2021) Does resisted sled towing improve the physical qualities of elite youth soccer players of differing maturity status? Journal of Science in Sport and Exercise. ISSN 2096-6709 DOI: <https://doi.org/10.1007/s42978-020-00087-w>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/7053/>

Document Version:

Article (Published Version)

Creative Commons: Attribution 4.0

© The Author(s) 2021

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.



Does Resisted Sled Towing Improve the Physical Qualities of Elite Youth Soccer Players of Differing Maturity Status?

Rhys Morris¹ · Tony Myers² · Stacey Emmonds³ · Dave Singleton⁴ · Kevin Till^{3,5} 

Received: 23 April 2020 / Accepted: 10 September 2020
© The Author(s) 2021

Abstract

Purpose Sled towing has been shown to be an effective method to enhance the physical qualities in youth athletes. The aim of this study was to evaluate the impact of a 6-week sled towing intervention on muscular strength, speed and power in elite youth soccer players of differing maturity status.

Method Seventy-three male elite youth soccer players aged 12–18 years (Pre-Peak Height Velocity [PHV] $n=25$; Circa-PHV $n=24$; Post-PHV $n=24$) from one professional soccer academy participated in this study. Sprint assessments (10 m and 30 m), countermovement jump and isometric mid-thigh pull were undertaken before (T1) and after (T2) a 6-week intervention. The training intervention consisted of 6 weeks (2× per week, 10 sprints over 20 m distance) of resisted sled towing (linear progression 10%–30% of body mass) during the competitive season. Bayesian regression models analysed differences between T1 and T2 within each maturity group.

Results There were minimal changes in strength, speed and power ($P=0.35$ – 0.80) for each maturity group across the 6-week intervention. Where there were changes with greater certainty, they are unlikely to represent real effect due to higher regression to the mean (RTM).

Conclusion It appears that a 6-week sled towing training programme with loadings of 10%–30% body mass only maintains physical qualities in elite youth soccer players pre-, circa-, and post-PHV. Further research is required to determine the effectiveness of this training method in long-term athletic development programmes.

Keywords Youth development · Long-term athletic development · Maturation · Strength · Speed · IMTP

Introduction

Many soccer academies are investing resources into the physical development of their players [43]. However, the large discrepancy of facilities and resources available

throughout academy structures may be a limiting factor that hinders the physical development of such players. As such, not all academies have sports science support to facilitate physical development across the entire pathway, which may have a knock-on effect on weightlifting competency as the players get older, which may hinder development. From a research perspective, many resistance training studies have failed to equate training volume and load across gym-based interventions, making it difficult to understand the differences in development between maturity groups. However, sled towing may offer a cheap and effective alternative to enhancing strength, speed and power; which are essential physical qualities for soccer [14], while also offering the opportunity to undertake controlled and balanced interventions across maturity groups to understand the influence of maturation on training adaptations. While the primary focus of this study was to offer an alternative for clubs with limited facilities, this is not to say, sled towing cannot supplement

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s42978-020-00087-w>) contains supplementary material, which is available to authorized users.

✉ Kevin Till
k.till@leedsbeckett.ac.uk

¹ School of Life Sciences, Coventry University, Coventry, UK

² Sport, Exercise and Health Research Center, Newman University Birmingham, Birmingham, UK

³ Carnegie School of Sport, Leeds Beckett University, Leeds, UK

⁴ Birmingham City Football Club, Birmingham, UK

⁵ Leeds Rhinos RLFC, Leeds, UK

training interventions for those practitioners that have the luxury of gym facilities and staffing.

Sled towing is a popular training method in adult populations [12, 23, 34] and has been demonstrated to improve acceleration, maximum velocity and its underlying mechanisms (e.g., maximal net horizontal ground reaction force, [29]). Lighter loads are reported to improve maximum velocity capabilities [17, 24] while heavier loads target acceleration [20]. More recently, the use of heavier loadings that cause a velocity loss of 50% [12], is believed to target training conditions that produce maximal power output (optimal force and optimal velocity, [11]) to enhance performance [10, 19, 42]. However, limited evidence exists using this method in youth populations and the impact it may have on physical development, especially considering maturity status.

Within youth athletes, the evidence based on the use of sled towing is less developed [5, 6, 36]. Recently, Cahill et al. [7] demonstrated improved physical performance following a sled pushing intervention, however, the authors suggested that many differences exist between sled pushing and pulling, which likely result in unique kinematic and kinetic changes, and sled pushing should be viewed as a unique and specialised form of horizontal resistance training. With this in mind, only one study to date [36] has investigated sled towing (pulling) in elite youth athletes across a range of maturity statuses. Rumpf [36] reported a 6-week sled intervention improved the speed, stride length, stride frequency, force, and power in pubertal boys, but it had no benefit for prepubertal boys. These findings suggested sled towing may improve other physical qualities and not just speed. Therefore, sled towing might be a practical method to enhance relative force production across maturity groups. However, Rumpf et al.'s [36] study had limitations including, (1) the utilisation of a non-motorized treadmill which has been shown to place an inherent resistance, reducing the maximal speed obtained, compared with overground sprinting [12, 30], (2) the combination of the Circa- and Post-peak height velocity (PHV) groups makes it difficult to partition out the

maturation effect, and (3) their statistical approach should be interpreted with caution given the growing concerns when using a *P*-value to make inferential assumptions on group differences [41] and the reporting of a percentage change pre and post-intervention, rather than the inclusion of baseline performance as a covariate [37], which may inflate the findings.

To summarize, maturity status is an important consideration when aiming to optimize youth athlete development. However, practitioners working with youth athletes may experience challenges (i.e., time, facilities, athlete competence) for optimizing strength, speed and power development and researchers may fail to implement controlled and balanced training programmes across differing maturity groups. This is apparent given the limited studies investigating across a range of maturity statuses in elite youth soccer players [27]. Although Rumpf et al. [36] investigated, a sled towing intervention in elite youth athletes' limitations exist in their study. Therefore, the purpose of this study was to evaluate a 6-week sled towing intervention using relative loadings (10%, 20% and 30% body mass) on speed, strength and power performance in youth soccer athletes across maturity statuses (pre-, circa- and post-PHV).

Methods

Subjects

Seventy-three male elite youth soccer players aged 12–18 years (pre-PHV $n=25$; circa-PHV $n=24$; post-PHV $n=24$) were recruited from one professional soccer academy (see Table 1 for descriptive data). The pre-PHV and circa-PHV participants trained on average four football sessions per week, with one competitive match and the post-PHV group trained on average, six football training sessions and one competitive match per week. The sled towing intervention replaced their usual strength and conditioning

Table 1 Descriptive parameters of anthropometric measurements pre and post sled towing intervention for all maturity groups

Anthropometry Parameters	Pre-PHV ($n=25$)		Circa-PHV ($n=24$)		Post-PHV ($n=24$)	
	Pre	Post	Pre	Post	Pre	Post
Age (years)	12.19 ± 0.51	12.38 ± 0.51	14.30 ± 0.84	14.42 ± 0.85	16.34 ± 1.18	16.47 ± 1.18
Years from PHV (years)	−1.97 ± 0.5	−1.88 ± 0.47	−0.15 ± 0.58	−0.11 ± 0.57	2.25 ± 1.01	2.27 ± 1.00
Height (cm)	150.88 ± 7.99	151.94 ± 7.75	164.87 ± 6.99	165.23 ± 6.82	179.38 ± 7.1	179.46 ± 7.02
Body mass (kg)	41.53 ± 6.22	41.95 ± 6.16	53.92 ± 5.31	54.1 ± 5.28	71.74 ± 9.11	71.65 ± 9.29
Sitting height (cm)	75.2 ± 3.24	75.63 ± 3.1	82.01 ± 3.08	82.16 ± 2.96	91.85 ± 4.95	91.89 ± 4.54
Leg length (cm)	75.6 ± 5.73	76.3 ± 5.6	82.85 ± 4.93	83.07 ± 4.95	87.53 ± 4.28	87.57 ± 4.33

PHV peak height velocity

training practices. All experimental procedures gained institutional ethics approval with informed and parental written consent obtained.

Experimental Approach to the Problem

A pre (T1) and post (T2) intervention study design was used to assess the impact of a 6-week sled towing intervention on speed, power and strength performance in elite youth soccer players. The study was implemented in-season, and testing took place at week 1 and week 8 with the intervention implemented on weeks 2–7. Before data collection, the players attended a familiarisation session that consisted of 10×20 m sled towing sprints (FH Pro Mini Speed Sledge Team Series; dimensions; 53 cm length 38 cm width 22 cm height, 300 g) with no additional weight attached. All testing and sled sessions were conducted on a 3G surface for consistency and took place at least 48 h post competitive match-play for all participants (see Supplementary Video 1). Before testing and the training intervention, all subjects performed a standardized 10-min warm-up consisting of jogging, dynamic stretching and acceleration drills. The players were accustomed to the physical tests as they were a part of their standard testing battery.

Procedures

Anthropometry

Height and sitting height were measured to the nearest 0.1 cm using a Seca Alpha stadiometer. Body mass was determined from body weight and taken to be $BW \text{ (kg)}/g$ with g = acceleration due to gravity measured on a commercially available portable force platform (AccuPower, AMTI, ACP, Watertown, MA) using a sampling rate of 1000 Hz then multiplied by 9.81 to convert to kg.

Maturity Offset

Age at PHV was estimated by the Mirwald prediction equation [25]. Years from PHV (YPHV) was calculated for each subject by subtracting the age at PHV from chronological age with a ± 6 months error rate. Subjects were allocated to either pre-PHV (offset < -1 years), circa-PHV (between -0.99 and $+0.99$ years) or post-PHV ($> +1$ years) groups in relation to their YPHV [31].

Sprint Performance

For sprint performance, distances of 10 and 30 m were assessed using Brower photocell timing gates (model number BRO001; Brower, Draper, UT, USA). All subjects

performed two trials, with 3–5 min of rest between trials. Athletes started 0.5 m behind the first gate from a two-point staggered start [38]. The best performance from each of the two trials was used for analysis. Intraclass correlation coefficient (ICC) and coefficient of variation (CV) for 10 m were $r=0.84$ and $CV=3.6\%$ and 30 m were $r=0.81$ and $CV=2.5\%$.

Isometric Mid-Thigh Pull (IMTP)

The IMTP was utilised as a measure of lower body strength. The IMTP was performed on a commercially available portable force platform (AccuPower, AMTI, ACP, Watertown, MA) and recorded vertical force at 1000 Hz. The data was not filtered. Subjects performed the IMTP on a customized pull rack with their shoulders placed over the bar in a position similar to 1 that of the second pull of a power clean [16]. Subjects performed two IMTP trials, each lasting 6 s (the pull lasting 5 s), with 3 min' rest between trials. The IMTP start was identified using a 5 SD threshold that was calculated from 1 s of quiet standing force recorded before the start of each pull [32]. Participants were instructed to push as "fast and hard" as possible and received loud verbal encouragement [32]. Each participant's best trial, as determined by the highest peak force (PF), was selected for analysis. Relative peak force (rPF) and relative impulse at 100 (rImp100) and 300 ms (rImp300), was calculated as $PF/\text{body mass}$, $\text{impulse}/\text{body mass}$ respectively. ICC and CV for the IMTP variables were as follows, PF and rPF were $r=0.98$ and $CV=4.91\%$. For Imp100 and Imp300, ICC and CV were $r=0.88$, $CV=5.0\%$, $r=0.84$, $CV=8.5\%$, respectively. Relative Imp100 and rImp300 were $r=0.88$, $CV=9.6\%$, $r=0.84$, $CV=9.8\%$, respectively.

Countermovement Jump (CMJ)

The CMJ was utilised as a measurement for lower body power. The CMJ was performed on the same portable force platform sampling vertical force (F_z) at 1000 Hz, which was not filtered. After a 1 s quiet standing period to identify initiation of movement, using the same 5SD threshold [8], CMJ was performed utilising a standard technique with arms akimbo [18], with no attempts made to control the depth of the countermovement [31]. Each participant performed two jumps interspersed with 3 min rest. Jump height was calculated using the velocity at take-off method [31] whilst net force was integrated with respect to time to obtain net impulse, which was summed over the propulsion phase. ICCs and CVs for the CMJ jump height (JH) were $r=0.86$, and $CV=7.5\%$ and CMJ impulse (CMJ Imp) were $r=0.94$ and $CV=6.7\%$.

Training Intervention

The training intervention consisted of 6 weeks of resisted sled towing, two sessions per week with a total of 12 sessions during the competitive season. The players had to have completed ten or more sessions (out of a total of 12), which is above an adherence rate of 80% to be included in the T2 post-testing session. From the start of the intervention, there were 102 players, across the age groups, with 73 players completing 10 or more sessions. This is an adherence rate of 72%. Each session consisted of 10 sprints over 20 m distance. The recovery between sets was approximately 90 s. Sessional training volume (calculated by number of sprints \times distance \times load) as reported by Rumpf et al. [36], which was altered every two weeks (every four sessions) by increasing the relative loads on the sleds from 10% (weeks 1–2; session load 20,000 AU), 20% (weeks 3–4; session load 40,000 AU) and 30% (weeks 5–6; session load 60,000 AU). No other load was monitored was monitored during this intervention.

Statistical Analysis

Descriptive statistics are reported as mean and standard deviation (SD) and median and median absolute deviation (MAD). To model pairwise differences between pre and post-scores within each maturity group, Bayesian regression models were fitted using a Student-*t* distribution [1] and allowing unequal variances between conditions. Weakly informative priors were used which included an improper flat prior over the reals for *b* values, a half Student-*t* prior with 3 degrees of freedom and a scale parameter of 10 for sigma with gamma on nu (shape = 2 and rate = 0.1). To illustrate the uncertainty around the estimation, lower and upper

95% Higher Density Intervals (HDI) are reported. Probability values of a change being greater or less than 0 ($P > 0$ or < 0) are provided and a standardized effect size calculated from the posterior estimates along with lower and upper 95% HDIs. For purposes of comparison, the effect sizes (ES) 0.2, 0.5 and 0.8 were considered to represent small, moderate and large differences, respectively [9]. The percentage of regression to the mean (RTM) was calculated. This was done because RTM can make a natural variation in repeated measures data look like real change. All models were checked for convergence ($\hat{r} = 1$), visual posterior predictive checks were conducted no systematic discrepancies between the predictive distribution and observed data *y* [15]. All analyses were conducted using R (Version 3.5. 2, R Foundation for Statistical Computing, Vienna, Austria, 2018) and with the Bayesian Regression Models in Stan (brms) package [4] which uses Stan (Version 2, 2016) for Hamiltonian Markov Chain Monte Carlo (MCMC). ICC's and CV's were analyzed for the performance variables described above during T1 and between repetitions.

Results

The mean and SD for anthropometric and maturation characteristics are displayed in Table 1. Tables 2 and 3 present the mean (SD) and median (MAD) descriptive statistics for performance measurements at T1 and T2. Tables 4, 5 and 6 display the inferential statistics for each maturity group.

Pre-PHV

The means for T1 and T2 suggest minimal improvements occurred across all performance variables apart from 10 m

Table 2 Pre and post descriptive statistics (mean \pm SD) for the performance measures across maturity groups

	Pre-PHV (<i>n</i> = 25)		Circa-PHV (<i>n</i> = 24)		Post-PHV (<i>n</i> = 24)	
	Pre	Post	Pre	Post	Pre	Post
10 m (s)	1.94 \pm 0.11	1.96 \pm 0.10	1.84 \pm 0.1	1.83 \pm 0.11	1.75 \pm 0.06	1.75 \pm 0.05
30 m (s)	4.90 \pm 0.30	4.87 \pm 0.30	4.53 \pm 0.25	4.50 \pm 0.27	4.20 \pm 0.13	4.18 \pm 0.11
CMJ Imp (N·s)	99.7 \pm 15.2	102.4 \pm 27.5	139.2 \pm 19.7	139.3 \pm 21.2	196.7 \pm 32.5	196.6 \pm 31.9
JH (m)	0.28 \pm 0.06	0.28 \pm 0.07	0.31 \pm 0.04	0.31 \pm 0.05	0.36 \pm 0.04	0.35 \pm 0.03
Imp100 (N·s)	20.7 \pm 6.8	21.3 \pm 7.0	32.8 \pm 14.1	34.1 \pm 13.8	50.0 \pm 12.5	50.5 \pm 19.3
rImp100 (N·s/kg)	0.5 \pm 0.2	0.5 \pm 0.2	0.6 \pm 0.3	0.6 \pm 0.3	0.7 \pm 0.2	0.7 \pm 0.2
Imp300 (N·s)	138.3 \pm 42.2	150.6 \pm 43.2	218.7 \pm 75.9	218.9 \pm 58.4	271.3 \pm 115.7	303.6 \pm 66.1
rImp300 (N·s/kg)	3.4 \pm 1.1	3.6 \pm 1.0	4.1 \pm 1.4	4.1 \pm 1	3.8 \pm 1.6	4.2 \pm 0.8
PF (N)	1197.0 \pm 157.7	1239.8 \pm 174.1	1638.6 \pm 193.6	1669.3 \pm 226.7	2318.0 \pm 356.9	2296.3 \pm 375.8
rPF (N/kg)	28.9 \pm 3.3	30.1 \pm 4.6	30.5 \pm 2.8	31.0 \pm 3.8	32.3 \pm 3.7	32.2 \pm 4.3

CMJ counter movement jump, CMJ Imp CMJ impulse, Imp100 impulse at 100 ms, Imp300 impulse at 300 ms, JH jump height, PF peak force, rImp100 relative impulse at 100 ms, rImp300 relative impulse at 300 ms, rPF relative peak force

Table 3 Pre and post descriptive statistics (median \pm MAD) for the performance measures across maturity groups

	Pre-PHV ($n=25$)		Circa-PHV ($n=24$)		Post-PHV ($n=24$)	
	Pre	Post	Pre	Post	Pre	Post
10 m (s)	1.95 \pm 0.09	1.94 \pm 0.1	1.84 \pm 0.09	1.81 \pm 0.08	1.76 \pm 0.04	1.74 \pm 0.04
30 m (s)	4.89 \pm 0.27	4.81 \pm 0.3	4.52 \pm 0.27	4.46 \pm 0.21	4.21 \pm 0.16	4.18 \pm 0.11
CMJ Imp (N·s)	99.2 \pm 14.0	95.1 \pm 14.3	139.2 \pm 18.8	138.0 \pm 19.5	185.1 \pm 23.7	192.4 \pm 27.6
JH (m)	0.28 \pm 0.06	0.27 \pm 0.08	0.31 \pm 0.06	0.31 \pm 0.04	0.35 \pm 0.04	0.35 \pm 0.03
Imp100 (N·s)	19.1 \pm 6.1	21.8 \pm 5.4	29.7 \pm 11.4	34.1 \pm 14.5	51.2 \pm 11.0	47.7 \pm 15.2
rImp100 (N·s/kg)	0.5 \pm 0.2	0.5 \pm 0.2	0.6 \pm 0.2	0.7 \pm 0.2	0.7 \pm 0.2	0.7 \pm 0.2
Imp300 (N·s)	144.3 \pm 38.7	151.3 \pm 58.5	211.8 \pm 71.9	214.5 \pm 55.8	303.2 \pm 74.9	317.6 \pm 62.8
rImp300 (N·s/kg)	3.5 \pm 0.9	3.7 \pm 1.1	4.2 \pm 1.1	3.9 \pm 1.0	4.3 \pm 0.8	4.1 \pm 0.8
PF (N)	1248.1 \pm 133.5	1249.9 \pm 108.4	1591.2 \pm 186.1	1618.2 \pm 220.2	2313.2 \pm 316.3	2272.1 \pm 346.4
rPF (N/kg)	28.5 \pm 3.3	30.2 \pm 3.9	30.3 \pm 3.5	30.9 \pm 4.6	32.0 \pm 2.4	31.4 \pm 4.2

CMJ counter movement jump, CMJ Imp CMJ impulse, Imp100 impulse at 100 ms, Imp300 impulse at 300 ms, JH jump height, MAD median absolute deviation, PF peak force, rImp100 relative impulse at 100 ms, rImp300 relative impulse at 300 ms, rPF relative peak force

Table 4 Bayesian *T*-test comparing pre and post-scores for pre-PHV ($n=25$) maturity group following a 6-week sled intervention with estimated mean data

Measure	Estimated difference	Estimated sigma	HDI's	$P > 0$	$P < 0$	ES	RTM (%)
10 m (s)	0.01	1.08	-0.05, 0.08	0.69	0.31	0.15 (-0.45, 0.68)	56
30 m (s)	-0.03	1.30	-0.21, 0.15	0.37	0.63	-0.11 (-0.69, 0.49)	56
CMJ Imp (N·s)	-0.75	12.31	-9.25, 7.95	0.42	0.58	-0.1 (-0.76, 0.58)	26
JH (m)	0.00	1.27	-0.04, 0.04	0.44	0.56	-0.05 (-0.64, 0.53)	18
Imp100 (N·s)	0.56	1.27	-3.39, 4.56	0.61	0.39	0.07 (-0.47, 0.67)	55
rImp100 (N·s/kg)	0.01	1.11	-0.1, 0.13	0.59	0.41	0.08 (-0.49, 0.68)	43
Imp300 (N·s)	10.84	1.27	-14.56, 36.98	0.80	0.20	0.27 (-0.29, 0.86)	40
rImp300 (N·s/kg)	0.25	1.95	-0.33, 0.82	0.80	0.20	0.26 (-0.35, 0.86)	51
PF (N)	40.19	150.99	-55.99, 132.86	0.79	0.21	0.25 (-0.33, 0.82)	30
rPF (N/kg)	1.16	4.56	-1.09, 3.42	0.85	0.15	0.31 (-0.31, 0.87)	54

CMJ counter movement jump, CMJ Imp CMJ impulse, ES standardized effect size, Estimated difference estimated difference between pre and post-performance, Estimated sigma combined pre and post variation of estimated difference, HDI's 95% higher density intervals, Imp100 impulse at 100 ms, Imp300 impulse at 300 ms, JH jump height, $P > 0$ probability greater than 0, $P < 0$ probability less than 0, PF peak force, rImp100 relative impulse at 100 ms, rImp300 relative impulse at 300 ms, RTM % regression to the mean, rPF relative peak force

sprint performance and CMJ jump height (Table 2, Figs. 1, 2 and 3). The median suggests all performance measures improved except CMJ impulse and jump height (Table 3). The estimated differences suggest any improvement post-intervention for the pre-PHV are highly uncertain (Table 4). There was also high RTM across measures, implying that even where differences were associated with less uncertainty, they are unlikely to represent real effect. The standardized differences (ES) across all measures, at best, represent small effects, and again these are highly uncertain. Overall, the results suggest the 6-week sled intervention did not change physical qualities within the pre-PHV group.

Circa-PHV

The descriptive statistics (mean and median) suggest minor improvements across most performance variables except CMJImp and rImp300 where performance declined (Tables 2 and 3, Figs. 1, 2 and 3). Modelled population differences for the Circa-PHV group were highly uncertain in terms of the intervention improving performance (Table 5). The standardized differences (ES) across all measures represent small effects at best and are again highly uncertain. Overall, the results suggest no change in physical qualities within the Circa-PHV group.

Table 5 Bayesian *T*-test comparing pre and post-scores for circa-PHV ($n=24$) maturity group following a 6-week sled intervention with estimated mean data

Measure	Estimated difference	Estimated sigma	HDI's	$P>0$	$P<0$	ES	RTM (%)
10 m (s)	-0.01	1.22	-0.07, 0.05	0.40	0.6	-0.09 (-0.66, 0.51)	56
30 m (s)	-0.03	1.32	-0.18, 0.13	0.36	0.64	-0.12 (-0.77, 0.46)	45
CMJ Imp (N·s)	-0.04	20.17	-11.86, 12.39	0.49	0.51	-0.01 (-0.59, 0.61)	30
JH (m)	0.00	1.13	-0.02, 0.03	0.63	0.37	0.08 (-0.46, 0.69)	23
Imp100 (N·s)	1.63	14.72	-6.56, 9.8	0.65	0.35	0.12 (-0.48, 0.77)	43
rImp100 (N·s/kg)	0.03	1.23	-0.13, 0.18	0.63	0.37	0.12 (-0.51, 0.67)	45
Imp300 (N·s)	-3.7	68.82	-41.83, 35.99	0.42	0.58	-0.05 (-0.67, 0.54)	38
rImp300 (N·s/kg)	-0.11	2.06	-0.85, 0.62	0.80	0.2	-0.1 (-0.75, 0.48)	36
PF (N)	29.13	189.81	-98.18, 153.61	0.68	0.32	0.14 (-0.43, 0.73)	20
rPF (N/kg)	0.46	4.14	-1.53, 2.55	0.67	0.33	0.14 (-0.46, 0.72)	25

CMJ counter movement jump, CMJ Imp CMJ impulse, ES standardized effect size, Estimated difference estimated difference between pre and post-performance, Estimated sigma combined pre and post variation of estimated difference, HDI's 95% higher density intervals, Imp100 impulse at 100 ms, Imp300 impulse at 300 ms, JH jump height, $P>0$ probability greater than 0, $P<0$ probability less than 0, PF peak force, rImp100 relative impulse at 100 ms, rImp300 relative impulse at 300 ms, RTM % regression to the mean, rPF relative peak force

Table 6 Bayesian *T*-test comparing pre and post-scores for post-PHV ($n=24$) maturity group following a 6-week sled intervention

Measure	Estimated difference	Estimated sigma	HDI's	$P>0$	$P<0$	ES	RTM (%)
10 m (s)	0.00	0.90	-0.03, 0.03	0.51	0.49	0.01 (-0.58, 0.6)	36
30 m (s)	-0.01	0.97	-0.09, 0.06	0.35	0.65	-0.12 (-0.72, 0.47)	13
CMJ Imp (N·s)	-0.21	31.85	-19.18, 18.9	0.49	0.51	-0.08 (-0.66, 0.52)	4
JH (m)	0.00	0.80	-0.03, 0.02	0.36	0.64	-0.12 (-0.75, 0.53)	30
Imp100 (N·s)	-0.88	13.20	-9.93, 8.2	0.42	0.58	-0.09 (-0.69, 0.56)	95
rImp100 (N·s/kg)	-0.02	1.18	-0.15, 0.11	0.38	0.62	-0.08 (-0.71, 0.5)	97
Imp300 (N·s)	24.41	108.36	-29.16, 79.86	0.81	0.19	0.19 (-0.44, 0.82)	97
rImp300 (N·s/kg)	0.32	2.00	-0.4, 1.09	0.78	0.22	0.14 (-0.54, 0.77)	81
PF (N)	-19.37	366.09	-240.24, 203.69	0.42	0.58	-0.08 (-0.66, 0.52)	27
rPF (N/kg)	-0.29	4.57	-2.46, 1.96	0.38	0.62	-0.09 (-0.69, 0.56)	53

CMJ counter movement jump, CMJ Imp CMJ impulse, ES standardized effect size, Estimated difference estimated difference between pre and post-performance, Estimated sigma combined pre and post variation of estimated difference, HDI's 95% higher density intervals, Imp100 impulse at 100 ms, Imp300 impulse at 300 ms, JH jump height, $P>0$ probability greater than 0, $P<0$ probability less than 0, PF peak force, rImp100 relative impulse at 100 ms, rImp300 relative impulse at 300 ms, RTM % regression to the mean, rPF relative peak force

Post-PHV

Data suggest minor improvements across all performance variables apart from CMJImp, JH, rImp100, PF and rPF (Table 2). The median suggests improvements in only 10 m and 30 m sprints, CMJImp, and Imp300, the remaining variables suggested decreases from T1 to T2 (Table 3, Figs. 1, 2 and 3). As with the other two groups, the estimated differences for the Post-PHV group suggest that most differences are highly uncertain for the intervention being successful in improving performance (Table 6). While there are performance increases for Imp300 and rImp300, it appears less uncertain these variables had very high levels of RTM and are unlikely to represent a real effect of the intervention as a result. Again, the ES

across all measures demonstrated small and highly uncertain effects. Overall, the results show high levels of uncertainty and RTM, and no change within Post-PHV players.

Discussion

This study aimed to evaluate the effect of a 6-week sled towing intervention on the physical qualities of youth soccer players across pre-, circa-, and post-PHV maturity groups. The current findings suggest that the sled towing intervention had minimal impact on the physical qualities of youth soccer players across all maturity groups. The result showed uncertainty and RTM and therefore observing a real effect

Fig. 1 Relative strength pre- and post-scores for pre-, circa- and post-PHV maturity groups following a 6-week sled intervention




Fig. 2 Countermovement jump height pre and post-scores for pre-, circa- and post-PHV maturity groups following a 6-week sled intervention





Fig. 3 The 10 m sprint times pre- and post-scores for pre-, circa- and post-PHV maturity groups following a 6-week sled intervention



was unlikely across all groups. These findings suggest that a bi-weekly sled towing training intervention for 6 weeks may not change the physical performance of youth soccer players. Still, there may be multiple explanations for such findings.

Sled towing training interventions have been widely used in senior populations and are associated with enhancing speed development [2, 12] along with other physical attributes (e.g., force and velocity, [12, 29]). However, the literature on the effect of sled towing in youth athletes is less developed. To the authors' knowledge, only one other study has considered maturity status and a 6-week sled intervention in elite youth athletes [36]. The current findings for pre-PHV are consistent with Rumpf et al. [36] where no changes were observed following the 6-week intervention. At the same time, the *p*-values (probability values) associated with the pre-PHV are around 0.80 which can be interpreted as 80% chance that the variables were greater in T2 and 20% chance they were lower. In its first instance, this seems like a positive change, however, all the variables with a *p*-value around 0.8 are associated with high uncertainty, and therefore, a real effect following the intervention is unlikely (associated high percentages of RTM) for the pre-PHV group. However, Rumpf et al. [36] reported significant improvements for the combined group (circa-post-PHV), which is in contrast to the current study's findings. The authors failed to discuss mechanisms as to why the pre-PHV failed to show any improvements but elude to a potential maturity-related response for the combined group.

The current findings appear to suggest that prepubertal boys may not improve physical qualities by sled towing that uses maximum BM loads of 30%. The loads used in the present study were heavier than those of the Rumpf et al. [36] (10% vs. 30% respectively) and were heavily influenced by the recent shift in philosophy for increased loading paradigms [12]. As such, very heavy sled loads have been shown to enhance athletic performance in senior athletes [12, 29] with loads around 69%–89% BM to be used. The current study aimed to advance from the loads used in the Rumpf et al. [36], given their non-significant findings for the pre-PHV group, while considering there may be an increased risk in injury with loads as high as 89% BM in youth athletes. While position statements advocate for the use of heavy loads in youths [21], given the limited empirical evidence surrounding sled towing in youth athletes, caution must proceed until evidence is provided demonstrating the use of heavier loads is safe. However, the sled loads used for the pre-PHV group may have not been appropriate, and, likely, the training impetus was not maturity-specific. For instance, when we consider the impact maturation has on physical development, it is believed that maturity-dependent response may occur following a specific intervention, which may be indicative of "synergistic adaptation" [35]. A "synergistic adaptation" refers to the symbiotic relationship

between specific adaptations of an imposed training demand and concomitant growth and maturity-related adaptations [22]. Whilst, Van Hooren et al. [39] question the validity of "sensitive periods" for specific physical qualities, it certainly does not detract from the notion of a "synergistic adaptation" following interventions in youth athletes. For example, it is not uncommon for prepubertal athletes to improve physical performance (mainly) through neuromuscular improvements [33] due to the plasticity of the central nervous system [40]. As such, plyometric training is believed to induce changes in motor unit recruitment, contraction velocity, preactivation and a greater reliance on the short-latency stretch reflex, resulting in a more feed-forward SSC function in prepubertal youth athletes [35]. Thus, plyometric training creates the ideal maturity-dependent impetus to achieve the desired "synergistic adaptation" specific for this maturity group.

Although sled towing may be a cyclical movement with SSC function (similar to that of a plyometric), adaptations following plyometric training are determined by the rate of the pre-stretch (i.e. the eccentric phase) of the activity. Therefore, it is likely the sled loadings were too heavy, increasing ground contact time (GCT) [2], which would reduce the pre-stretch (eccentric phase) and therefore limit the neurological signal sent from the muscle spindle reducing the shortening cycle of the plyometric movement [13]. This primarily denotes the term plyometric and changes the exercise impetus and is not the desired effect to achieve the "synergistic adaptation" for the pre-PHV boys [35]. Although GCT was not measured in the current study, it is plausible the lack of observed effects are a consequence of the loads used along with a range of other factors (e.g., concurrent training of soccer practice, increase stride length as a consequence of the sled load) and inherently do not create the ideal training impetus to reap the benefits for a "synergistic adaptation". It has also been established that youths of differing maturity status could respond to sprint training [26, 27]. A recent meta-analytical review showed far larger effects in postpubertal ($d = 1.39$) and pubertal ($d = 1.15$) athletes than in prepubertal ($d = -0.18$) boys [27], indicating a maturity-specific adaptation.

Similar to the pre-PHV group, both the circa- and post-PHV groups failed to improve following the sled intervention. Although high *P*-values (for example $P = 0.80$) have been reported for rImp300, this 80% probability is associated with a higher percentage of RTM of 36%. Therefore, there is uncertainty associated with these positive improvements. These findings are different from those reported by Rumpf et al. [36] who reported a moderate improvement of around 6% in the combined circa- and post-PHV in 30 m sprint time. However, the methodologies employed by Rumpf et al. [36] to investigate the impact of sled towing on other qualities should be scrutinised and have been discussed previously. Finally, the combined group is not a true representation of a

"synergistic adaptation" as both circa- and post-PHV groups will respond differently to a given bout of training or specifically to sprint training [27]. For example, Morris et al. [31] demonstrated a circa-PHV group had superior physical development across a football season, compared to the pre- and post-PHV groups. By combining maturity groups for Rumpf et al. [36], information around maturity specific adaptations and the potential "synergistic adaptation" is not clear. Although the current study's findings reveal no impact, it has demonstrated those findings for each maturity group.

While the current study examined the impact of sled towing across a range of maturity status' using a variety of practical assessment methods (IMTP, CMJ) including gold-standard techniques (e.g., force platforms), thus adding practical value to the paediatric literature. It is not without limitations. Firstly, limitations can exist for the classification of maturity status using the Mirwald equation [22]. As such, some of the players may have been misclassified into the incorrect maturity group given the associated error and close overlap of the group bandwidths. The authors acknowledge these limitations, but it was the most practical assessment available. Secondly, the lack of improvements observed in the current findings across the maturity groups, and the conflicting evidence reported by Rumpf et al. [36], may be a consequence of the methodology used to determine the sled towing loadings in the current study. Although the prescription of %BM method is practical and heavily cited [34], it is now under scrutiny [5, 12]. When using %BM prescription, although the load is standardized using BM, everyone will display different velocity decrements based on the same %BM loading [6, 12]. For example, one athlete may show a decrement of 20% compared to a 40% decrement in another athlete with the same loading creating a non-standardized effect [5]. In addition, Cahill et al. [6] investigated the load-velocity relationship and established that 50% decrement is achieved ~89% BM to target the Pmax zone in high-school athletes but reports a large variance (95% CI 71%–107% BM). This concept is derived from senior populations [12, 29], and Cahill et al. [6] follow the recommendations that a decrement of 50% is the optimum range to enhance accelerative capacity (Pmax zone), but warrants further investigation when we consider maturity status. Furthermore, the positive findings from Rumpf et al. [36], compared to the current study, may be a consequence of the specificity of testing to training. This principle has been shown in youth athletes [3] and could explain why the current study has not demonstrated a positive effect following the intervention. Also, the length of the intervention (minimum of 10 sessions) may have also contributed to the lack of findings. For example, Moran et al. [28] has demonstrated an increase in performance is attributed to where interventions last longer than 8 weeks [28]. It is plausible the length of the current studies intervention was inadequate. The limited evidence

in elite youth populations makes it hard to understand the maturational effect on sled performance. The recent study by Cahill et al. [6] identifies a load-velocity relationship with a population of Post-PHV athletes and in high-school males. Although, the authors use the narrative 'youth athletes' their population has a mean age of 16.7 ± 0.9 years and a mean YPHV of 1.8 ± 0.8 years. Therefore, the broader application of the load-velocity relationship to pre- and circa-PHV requires investigation. It is also a limitation for this study not to have a control group. Inclusion of a control group would substantiate the findings (or lack of) for those who have received the treatment (in this instance, the sled towing intervention). Finally, understanding kinematic and kinetic changes during sprinting would provide greater clarity surrounding the specific maturity-related changes more so than the generic testing battery currently used in the study. Future studies should look to include specific testing-training protocols [3] when monitoring changes from a sled towing intervention.

Conclusion

This study evaluated the impact a 6-week sled towing intervention on speed, lower body power and strength in youth soccer players across pre-, circa- and post-PHV maturity groups. Findings demonstrated that the sled intervention had minimal impact on the performance measures irrespective of maturity group. However, the results also revealed that none of the performance measures declined over the six weeks. Practically, sled towing may not be an effective strategy to improve physical performance, regardless of maturity status when used in isolation. These findings are likely to be a consequence of the loadings used (and the methodology used to determine such loadings; testing to training was not specific), coupled with the influence of maturation and the length of the intervention (only 6-weeks). For example, the desired 'synergistic adaptation' or maturity-related changes were not responsive to the loadings used (10%–30% BM). The pre-PHV group did not elicit any change which may be due to the inherent nature and reliance of force production required during sled towing which would not elicit changes in the neuromuscular properties, which are the maturity associated improvements in the group. Equally, those who were more mature also failed to improve performance with the increased loadings, which can increase strength, speed and power via neural and morphological adaptations. This study does warrant further investigation into heavier loadings for the circa- and post-PHV groups, while the pre-PHV group should focus on interventions that aim to decrease GCT, which may be more related to

the maturity-specific adaptations, such as plyometric training for long-term athletic development of youth athletes.

Funding Not applicable.

Data availability Raw data can be provided upon request.

Code availability Code for Bayesian modelling can be provided upon request (R Studio, R script software).

Compliance with Ethical Standards

Conflict of interest Not applicable.

Ethics approval Institutional Ethics was granted.

Consent to participate Parental assent and participant consent was obtained for all involved.

Consent for publication Club, parental assent and participant consent was obtained for all involved in the project.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aas K, Haff IH. The generalized hyperbolic skew student's t-distribution. *J Financ Econom*. 2006;4(2):275–309.
2. Alcaraz PE, Carlos-Vivas J, Oponjuru BO, Martinez-Rodriguez AJSM. The effectiveness of resisted sled training (RST) for sprint performance: a systematic review and meta-analysis. *Sports Med*. 2018;48(9):2143–65.
3. Behm DG, Young JD, Whitten JHD, Reid JC, Quigley PJ, Low J, Li Y, Lima CD, Hodgson DD, Chaouachi A, Prieske O, Granacher U. Effectiveness of traditional strength vs power training on muscle strength, power and speed with youth: a systematic review and meta-analysis. *Front Physiol*. 2017;8:423. <https://doi.org/10.3389/fphys.2017.00423>.
4. Bürkner P-C. brms: an R package for Bayesian multilevel models using Stan. *J Stat Softw*. 2017;80(1):1–28.
5. Cahill MJ, Cronin JB, Oliver JL, Clark KP, Lloyd RS, Cross MRJS, Journal C. Sled pushing and pulling to enhance speed capability. *Strength Cond J*. 2019;41(4):94–104.
6. Cahill MJ, Oliver JL, Cronin JB, Clark KP, Cross MR, Lloyd RSJS. Sled-pull load-velocity profiling and implications for sprint training prescription in young male athletes. *Sports*. 2019;7(5):119. <https://doi.org/10.3390/sports7050119>.
7. Cahill MJ, Oliver JL, Cronin JB, Clark KP, Cross MR, Lloyd RS. Influence of resisted sled-push training on the sprint force-velocity profile of male high school athletes. *Scand J Med Sci Sports*. 2019;30(3):545. <https://doi.org/10.1111/sms.13600>.
8. Chavda S, Bromley T, Jarvis P, Williams S, Bishop C, Turner AN, Lake JP, Mundy P. Force-time characteristics of the counter-movement jump: analyzing the curve in Excel. *Strength Cond J*. 2018;40(2):67–77.
9. Cohen J. Statistical power analysis. *Current directions in psychological science*. 1992;1(3):98–101.
10. Cormie P, McCaulley GO, Triplett NT, McBride JM. Optimal loading for maximal power output during lower-body resistance exercises. *Med Sci Sports Exerc*. 2007;39(2):340–9. <https://doi.org/10.1249/01.mss.0000246993.71599.bf>.
11. Cormie P, McGuigan MR, Newton RUJ. Developing maximal neuromuscular power: part 2—training considerations for improving maximal power production. *Sports Med*. 2011;41(2):125–47.
12. Cross MR, Brughelli M, Samozino P, Brown SR, Morin J-B. Optimal loading for maximizing power during sled-resisted sprinting. *Int J Sports Physiol Perform*. 2017;12(8):1069–77.
13. Davies G, Riemann BL, Manske R. Current concepts of plyometric exercise. *Int J Sports PhysTher*. 2015;10(6):760.
14. Deprez D, Coutts A, Fransen J, Deconinck F, Lenoir M, Vaeyens R, Philippaerts R. Relative Age, Biological Maturation and Anaerobic Characteristics in Elite Youth Soccer Players. *Int J Sports Med*. 2013;34(10):897–903.
15. Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. Visualization in Bayesian workflow. *J R Stat Soc Ser A*. 2019;182(Pt.2):389–402.
16. Haff GG, Ruben RP, Lider J, Twine C, Cormie PJT, Research C. A comparison of methods for determining the rate of force development during isometric midhigh clean pulls. *J Strength Cond Res*. 2015;29(2):386–95.
17. Harrison AJ, Bourke G. The effect of resisted sprint training on speed and strength performance in male rugby players. *J Strength Cond Res*. 2009;23(1):275–83.
18. Hori N, Newton RU, Andrews WA, Kawamori N, McGuigan MR, Nosaka K. Does performance of hang power clean differentiate performance of jumping, sprinting, and changing of direction? *J Strength Cond Res*. 2008;22(2):412–8.
19. Kawamori N, Haff GG. The optimal training load for the development of muscular power. *J Strength Cond Res*. 2004;18(3):675–84. [https://doi.org/10.1519/1533-4287\(2004\)18<675:Totlft>2.0.Co;2](https://doi.org/10.1519/1533-4287(2004)18<675:Totlft>2.0.Co;2).
20. Kawamori N, Newton RU, Hori N, Nosaka K. Effects of weighted sled towing with heavy versus light load on sprint acceleration ability. *J Strength Cond Res*. 2014;28(10):2738–45. <https://doi.org/10.1519/JSC.0b013e3182915ed4>.
21. Lesinski M, Prieske O, Granacher UBJSM. Effects and dose-response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br J Sports Med*. 2016;50(13):781–95.
22. Lloyd RS, Radnor JM, Croix MBDS, Cronin JB, Oliver JL. Changes in sprint and jump performances after traditional, plyometric, and combined resistance training in male youth pre-and post-peak height velocity. *J Strength Cond Res*. 2016;30(5):1239–47.
23. Lockie RG, Murphy AJ, Schultz AB, Knight TJ, de Jonge XAKJ. The effects of different speed training protocols on sprint acceleration kinematics and muscle strength and power in field sport athletes. *J Strength Cond Res*. 2012;26(6):1539–50. <https://doi.org/10.1519/JSC.0b013e318234e8a0>.
24. Lockie RG, Murphy AJ, Spinks CD. Effects of resisted sled towing on sprint kinematics in field-sport athletes. *J Strength Cond Res*. 2003;17(4):760–7.
25. Mirwald RL, Baxter-Jones AD, Bailey DA, Beunen GP. An assessment of maturity from anthropometric measurements. *Med Sci Sports Exerc*. 2002;34(4):689–94.

26. Moran J. The effectiveness of resistance, plyometric and sprint training at different stages of maturation in male youth athletes. PhD diss.: University of Essex; 2017.
27. Moran J, Parry DA, Lewis I, Collison J, Rumpf MC, Sandercock GRJ. Maturation-related adaptations in running speed in response to sprint training in youth soccer players. *J Sci Med Sport*. 2018;21(5):538–42.
28. Moran J, Sandercock G, Rumpf MC, Parry DAJ. Variation in responses to sprint training in male youth athletes: a meta-analysis. *Int J Sports Med*. 2017;38(1):1–11.
29. Morin JB, Petrakos G, Jimenez-Reyes P, Brown SR, Samozino P, Cross MR. Very-heavy sled training for improving horizontal-force output in soccer players. *Int J Sports Physiol Perform*. 2017;12(6):840–4. <https://doi.org/10.1123/ijsp.2016-0444>.
30. Morin JB, Seve P. Sprint running performance: comparison between treadmill and field conditions. *Eur J Appl Physiol*. 2011;111(8):1695–703. <https://doi.org/10.1007/s00421-010-1804-0>.
31. Morris R, Emmonds S, Jones B, Myers TD, Clarke ND, Lake J, Ellis M, Singleton D, Roe G, Till K. Seasonal changes in physical qualities of elite youth soccer players according to maturity status: comparisons with aged matched controls. *J Sci Med Footb*. 2018;2(4):272–80.
32. Morris RO, Jones B, Myers TD, Lake J, Emmonds S, Clarke ND, Till KJ. Isometric mid-thigh pull characteristics in elite youth male soccer players: comparisons by age and maturity offset. *J Strength Cond Res*. 2020;34(10):2947–55. <https://doi.org/10.1519/JSC.0000000000002673>.
33. Peña-González I, Fernández-Fernández J, Cervelló E, Moya-Ramón M. Effect of biological maturation on strength-related adaptations in young soccer players. *PLoS ONE*. 2019;14(7):e0219355.
34. Petrakos G, Morin J-B, Egan B. Resisted sled sprint training to improve sprint performance: a systematic review. *J Sports Med*. 2016;46(3):381–400.
35. Radnor JM, Oliver JL, Waugh CM, Myer GD, Moore IS, Lloyd RS. The influence of growth and maturation on stretch-shortening cycle function in youth. *J Sports Med*. 2018;48(1):57–71.
36. Rumpf MC, Cronin JB, Mohamad IN, Mohamad S, Oliver JL, Hughes MG. The effect of resisted sprint training on maximum sprint kinetics and kinematics in youth. *Eur J Sport Sci*. 2015;15(5):374–81.
37. Senn S. Change from baseline and analysis of covariance revisited. *Statistics in medicine*. 2006;25(24):4334–44.
38. Thomas C, Comfort P, Chiang C-Y, Jones PA. Relationship between isometric mid-thigh pull variables and sprint and change of direction performance in collegiate athletes. *J Trainol*. 2015;4(1):6–10.
39. Van Hooren B, Croix MDS. Sensitive Periods to Train General Motor Abilities in Children and Adolescents. *Strength Cond J*. 2020. Publish Ahead of Print.
40. Viru A, Loko J, Harro M, Volver A, Laaneots L, Viru M. Critical periods in the development of performance capacity during childhood and adolescence. *Eur J Phys Educ*. 1999;4(1):75–119.
41. Wasserstein RL, Lazar NA. The ASA's statement on p-values: context, process, and purpose. *Am Stat*. 2016;70(2):129–33.
42. Wilson GJ, Newton RU, Murphy AJ, Humphries BJ. The optimal training load for the development of dynamic athletic performance. *Med Sci Sports Exerc*. 1993;25(11):1279–86.
43. Wrigley R, Drust B, Stratton G, Atkinson G, Gregson W. Long-term Soccer-specific Training Enhances the Rate of Physical Development of Academy Soccer Players Independent of Maturation Status. *Int J Sports Med*. 2014;35(13):1090–4.