



LEEDS
BECKETT
UNIVERSITY

Citation:

Christian, MS and Evans, CE and Ransley, JK and Greenwood, DC and Thomas, JD and Cade, JE (2012) Process evaluation of a cluster randomised controlled trial of a school-based fruit and vegetable intervention: Project Tomato. *Public health nutrition*, 15 (3). 459 - 465. ISSN 1368-9800
DOI: <https://doi.org/10.1017/s1368980011001844>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/710/>

Document Version:

Article (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Process evaluation of a cluster randomised controlled trial of a school based fruit and vegetable intervention: Project Tomato

M.S. Kitchen (PhD student),¹ C.E. L. Evans (Research Fellow),¹ J.K. Ransley (Lecturer in Nutritional Epidemiology), D.C. Greenwood Senior(Lecturer in Medical Statistics)², J.D. Thomas(Database Manager), J.E. Cade (Professor of Nutritional Epidemiology and Public Health)

¹*Nutritional Epidemiology Group, Centre for Epidemiology and Biostatistics, University of Leeds, Room 8.001, The Worsley Building, University of Leeds, Leeds, UK, LS2 9JT, United Kingdom*

²*Centre for Epidemiology and Biostatistics, University of Leeds, Room 8.001, The Worsley Building, University of Leeds, Leeds, UK, LS2 9JT, United Kingdom*

*Correspondence to: Meaghan Kitchen, Nutritional Epidemiology Group, Centre for Epidemiology and Biostatistics, University of Leeds, Room 8.001, The Worsley Building, University of Leeds, Leeds, UK, LS2 9JT, United Kingdom; Tel: 0113 343 8907
Fax: 0113 343 4877 m.s.kitchen@leeds.ac.uk*

Keywords: Fruit, vegetable, interventions, process measures, children, family meal, UK

2990 words excluding titles, authors and addresses, abstract and references

ABSTRACT

Background Multi component interventions delivered at school may affect children's dietary behaviour. Where intervention trials are not successful the reasons why need to be explored.

This analysis evaluates the overall appreciation and implementation of an intervention designed to maintain fruit and vegetable intake in children aged 8-9 years.

Method A random sample was selected using schools as clusters across England; each group consisted of 27 primary schools. The intervention group received a multi-component programme delivered in school by teachers and items sent home for parents/children. Of the 311 children that completed the trial, 261 children participated in the process evaluations.

Dietary measurements were collected at baseline and follow-up. The intervention participants completed questionnaires on the intervention materials, to identify implementation and appreciation of the intervention, and other environmental mechanisms.

Results Implementation of the intervention was low, 21.3% of school items and 56% of home items were implemented. The intervention materials were well received by teachers, parents and children. Other mechanism that affect with fruit and vegetable intake, revealed parents who eat their main meal with their children 3-4 nights or 5-7 nights a week, on average had 48gram (95% CI: -15.4 to 111.7) and 34gram (95% CI: 6.4 to 62.0) respectively more fruit and vegetables intake, compared to parents who eat with their children less than three times per week.

Conclusion Implementation of the trial components was poor. However, the results identified the importance of parental environment and mealtime structure on children's fruit and vegetable intake.

INTRODUCTION

Children's health and nutritional intake remain a national priority in the UK. Nutritional education programmes have been developed for schools, home and community settings in an attempt to improve children's diets; however, it is still unclear which methodologies increase or maintain children's fruit and vegetable consumption.

As part of the UK's "5 A DAY" programme in 2004, the Department of Health launched the School Fruit and Vegetable Scheme (SFVS), aimed at increasing children's fruit and vegetable consumption by providing a free piece of fruit or vegetable every school day for children in Reception to Year 2 (aged 4-6 years). Three academic studies have evaluated the SFVS¹⁻³, suggesting that whilst the scheme appears to increase children's fruit intake, this increase is only short-term. The flexible multi-component intervention, Project Tomato, was designed to tackle this issue, aiming to encourage children to maintain fruit and vegetable consumption once eligibility for the SFVS ceased.

The Project Tomato intervention focused on curriculum based activities and tasting sessions to improve children's knowledge and awareness of dietary fruit and vegetables. However, the home environment and mealtime structure are vital components to a child's eating pattern.⁴ The Project Tomato intervention incorporated fun activities for parents and children to do together at home; as parents play a direct role in their child's eating behaviours and attitudes towards food.^{4,5}

Several systematic reviews have summarised existing research on school interventions to increase children's fruit and vegetable intake⁶⁻⁸ and these reviews suggest multi-component

interventions are the most successful in terms of improving fruit and vegetable consumption,⁹⁻
¹⁷ supporting the design methodology used to create the Project Tomato intervention. With
such complex interventions, identifying the combination of components that are the most
effective at increasing children's fruit and vegetable intake is difficult, illustrating the
importance of exploring the process evaluations for each study. Process evaluations are used
to improve the understanding of successful or unsuccessful health interventions; to identify the
key components that make an intervention successful, for boys, girls or both, and determine
which environment/conditions lead to these particular components facilitating a successful
outcome.^{18 19}

Despite the promising theory-based design of the Project Tomato intervention, it was not
successful at maintaining children's fruit and vegetable intake (unpublished data). The aim of
the present study is to investigate the level of implementation and appreciation of the different
elements involved in the Project Tomato intervention, to identify to which elements worked
and which did not. As well as exploring other possible factors within the home environment
that could be associated with fruit and vegetable intake.

METHODS

Population and study design

A nationally representative sample of 1031 children from 54 schools were recruited into the
trial.²⁰ Schools were randomised to receive the Project Tomato intervention or control.

Further detail on the sampling methodology is reported elsewhere²⁰. Ethical approval was
obtained through the University of Leeds Central Research Ethics Committee.

Process Evaluation

Twenty-four schools and 311 children completed the intervention. These schools were asked to complete a set of questionnaires on the intervention materials, sent to them at different periods during the intervention, for the teacher, parents or children to complete. The teachers were asked to complete a questionnaire about the school items, by ticking when they completed a task/item and commenting on a rating scale for each item. For example with the teacher lesson plans teachers were asked to tick if they did or did not use the lesson plan, and then rate their appreciation of the lesson plan on a Likert scale of 1-5. The parent questionnaire asked the parents to tick boxes if their child had received different intervention materials (yes or no), and to confirm if they used these materials. Parents were also asked to comment on whether they sent their children to school with fruit or vegetables, along with other family eating pattern questions; for example how many nights a week they ate their main meal with their child at a table. How many days a week parents sent their children to school with a piece of fruit or vegetables was categorised into 0-1, 2-3, 4-5 per week, and how many nights a week they ate their main meal together was categorised into 0-2, 3-4 and 5-7 nights per week. The children were asked to complete three questionnaires. Each questionnaire asked the children to comment on the intervention items sent in the three kit bags, using a Likert scale, with the children asked to tick one of the following options 'love it, like it, don't like it, hate it, don't know.' The children were also asked to tick if they had or had not used each item.

The intervention

The intervention was designed using the theories of maintenance of health behaviour as a basis.²¹ To apply the theories, underlying factors that have an effect on initiating and maintaining change in people's eating behaviour were described using the following acronym – FRAME: familiarisation, repetition, activities, modelling and environment^{7 20 21}. The

intervention consisted of core and customised materials designed for the whole school, the Year 3 classroom and for the children to take home. The core elements consisted of the following; the Project Tomato manual and 12 curriculum related lesson plans. The customised elements were: cooking (designed by the British Nutrition Foundation); growing club information (designed by the Royal Horticultural Society), funding support; and information to set up a Project Tomato Team. The customised elements were tailored to meet each school's needs. The head teacher could decide which of these elements were included in their school's package. The intervention also involved items designed for the parents and children to complete consisting of: three Project Tomato kit bags, two Project Tomato newsletters, and parent handouts. The duration of the intervention was 10 months from July 2007 to April 2008, and these materials were sent out at appropriate intervals throughout the school year.

Dietary Assessment

Dietary intake was collected using a 24 hour dietary assessment tool 'Child And Diet Evaluation Tool, (CADET)' diary.²² To complete the diary the participant has to tick each item consumed, under the appropriate meal time heading within the 24-hour period. For this analysis a National Foundation for Educational Research (NFER) field worker filled in the CADET diary for each child during school hours, and parents were asked to complete the evening and morning food consumption for their child.

Statistical analysis

To assess potential bias amongst parents and children who returned a process measures questionnaire and those that did not, multi-level regression model was conducted on school and child characteristics, to explore if there were any differences between the two groups.

Descriptive statistics for implementation and appreciation of school and home items were calculated for teachers, children and parents. Participants implementation and appreciation of either set of materials (school, home) was re-coded into variables with a range from 0 to 100 (zero meaning no items used or low implementation/appreciation; 100 meaning all items used/implemented or high appreciation). The total scores for implementation of school items; curriculum lessons, tasting sessions, cooking club and growing club were categorised into binary codes; no and yes. The total scores for implementation of parents was coded into tertiles; low, medium and high. The total scores for appreciation for all teachers, and parents was categorised into low and high.

To assess the association of different intervention materials (appreciation and implementation) on the difference in children's total fruit and vegetable intake, a two level multilevel regression model was used with total weight of fruit and vegetable as the predictor. This model took into consideration the hierarchical structures of the data caused by randomising by cluster (school).²⁰ Two models were assessed, the first was unadjusted and the second was adjusted for age, gender, ethnicity and Index of Multiple Deprivation (IMD) score²³. The mean and 95% confidence intervals were generated for all models.

RESULTS

The main outcome of the Project Tomato intervention has been reported elsewhere (Refer main paper). Of the 24 schools that completed the interventions, 79%, (19 teachers) completed the teacher questionnaire. The number of children that participated in the intervention with a completed baseline and follow-up CADET was 311. Of these children 84%

(261) responded to one of the three children's questionnaires. 38% (120) completed the parents' questionnaire.

A sensitivity analysis conducted to determine differences between children who completed a process measure questionnaire and children who did not, this analysis revealed there were no differences for child or school characteristics.

[INSERT TABLE 1 HERE]

Implementation and Appreciation of the Intervention for Teachers

Of the 24 intervention schools that completed the process measures questionnaires, eight schools did not implement any of the school items stated above. On average, schools implemented 21 percent of all school items. Table 2 describes the mean implementation, appreciation, and 95 percent confidence intervals for the school items. School curriculum lessons and tasting sessions were the most widely implemented items, whilst the most appreciated items were the tasting sessions and the cooking club lessons.

[INSERT TABLE 2 HERE]

Implementation and Appreciation of the Intervention for Parents and Children

The parents and children on average had low implementation scores of the home items, with the parents implementing 35% (N =57, 95%; CI: 30 to 40), and the children implementing 56% (N=115, 95% CI= 53 to 60). Both parents' and children's appreciation of the Project Tomato items was high, with a mean for parents of 76% (N=112, 95% CI= 73 to 81), and for

children 73% (N=101, 95% CI= 71 to 76). From the three child process measure questionnaires, the children were asked to comment if they used each item sent to them in one of the three kit bags, the results are presented in Table 3. The items that were, on average, used the most were the fruit and veg portion game, fruit ‘n’ veg snack box for fruit, the Eat5! Quiz book, and the balance of good health quiz book. The children were also asked to rate each intervention item from the kit bags, on a five-point scale (love it, like it, don’t know, don’t like it, hate it) this was re-coded into love/like it and don’t know, don’t like or hate it. On average, the Project Tomato items that the children liked/loved the most were; the fruity face, the Christmas cake recipe, eat 5 balloon, project tomato pencils, portion game, and fruit “n” veg snack box.

[INSERT TABLE 3 HERE]

Intervention implementation and the association with total fruit and vegetable intake

A two level multilevel regression model was conducted to explore the relationship between children’s total follow-up fruit and vegetable intake and overall implementation of school items. The unadjusted and adjusted (gender, ethnicity, baseline intake and IMD score) models are presented in Table 4.

[INSERT TABLE 4 HERE]

Regression models revealed that for total implementation of all school items (curriculum lessons and tasting sessions) and home items (see items included in Table 3), there was no significant difference between intervention implementation levels and follow-up fruit and vegetable intake in the control and intervention groups, after adjusting for baseline fruit and

vegetable intake, gender, ethnicity and IMD score. The unadjusted and adjusted models are presented in Table 4. Whilst Parents' level of implementation was not significantly associated with improvements in fruit and vegetable consumption, after adjusting for possible confounders, parents with a medium implementation of the intervention items were 40% more likely to have a higher fruit and vegetable intake, whereas parents with a high implementation level were 66% more likely to have children who consumed more fruit and vegetables compared to parents who had a low implementation rating. This suggests that children of parents who had a high or medium intervention implementation consumed more fruit and vegetables compared to those with a low level of implementation.

Parents were also asked "*How many days per school week do you send your child to school with fruit or vegetables?*" After adjusting for baseline fruit and vegetable intake, gender, ethnicity and IMD score, there was no significant difference in how many days per week children were sent to school with a piece of fruit or vegetable and their overall fruit and vegetable intake (N=111, 2-3days per week -6.1g 95% CI: -104.9 to 92.7, 4-5days per week 23.9g 95% CI: 25.0 to 72.3. p=0.3). Parents were also asked "*How many nights a week do you eat an evening meal with your child, at a table?*" The multi-level regression model revealed that there was a significant relationship between eating at a table and children's fruit and vegetable intake. Parent's who ate their evening meal at a table 5-7 nights a week were 34g (95% CI: 6.4 to 62.0) and parents who ate their evening meal at a table with their children 3-4 nights a week were 48g (N=102 95% CI: -15.4 to 111.7, p=0.03) more likely to have children with higher fruit and vegetable intake than children who ate their evening meals with their parents zero to two nights per week, after adjustments were taken into consideration.

Intervention appreciation and the association with total fruit and vegetable intake

Table 5 shows that although the appreciation of various aspects of the project was high it was not significantly associated with change in children's fruit and vegetable intake.

[INSERT TABLE 5 HERE]

DISCUSSION

The evaluation of the impact of the Project Tomato intervention indicated that a flexible multi-component intervention had little effect on maintaining or increasing children's fruit and vegetable intake in the UK. The components designed and used in this programme are similar to those used in other multi-component interventions which have been shown to be effective^{5 10 14 17 24 25}. However, implementation rate was a vital component in the success of these studies. Te Velde et al¹⁷ achieved long term change in children's fruit and vegetable intake in the country that had the highest implementation rate. Similarly, a successful intervention reported by Story et al²⁴ reported high adherence to the intervention materials in their process evaluation.

The total implementation of the intervention for teachers, parents and children was low; this implies that the effect of the intervention on the main outcome, fruit and vegetable intake, might have improved with stronger implementation. These results are similar to previous studies^{11 26} which had low implementation levels, and reported process measures stated that the primary barrier to teaching the intervention was preparation time. Multi-component interventions are often seen as labour intensive, and if the teacher-parent relationship is poor or the parents are rarely involved in school activities, barriers may exist to implementing

nutritional interventions⁷. The intervention was designed to be pragmatic and not relying on external agencies for its delivery. This was deliberate, since had the intervention been successful the approach would have been readily transferable between schools. Whereas the majority of successful fruit and vegetable interventions have had external assistance in training, delivering and running of the intervention.^{9 10 12 13}

Children's and parents' implementation levels of the home items provided a positive association with children's fruit and vegetable intake. Whilst this difference was not significant, parents who implemented more intervention items were more likely to have children with higher fruit and vegetable consumption. The analysis identified the "*fruit and veg portion game*" as one of the most used and liked items by children and parents. This activity involves each member of the household recording their daily intake of fruit and vegetables, to see who has eaten the most at the end of the week. It is a very simple concept that could easily be implemented, in public health initiatives.

Another important finding that was exploring other possible mechanisms associated with children's fruit and vegetable intake was the meal time environment. Results revealed that parents who eat their main meal with their children at a table at least three times per week, on average have higher fruit and vegetable intake compared to parents who eat together less often. This suggests that programmes encouraging parents to eat together with their children at a table, may be a more effective way of increasing fruit and vegetable intake. These findings support previous research that parents' influence is an important determinant of children's diet^{5 27}. Eating together at a table provides the perfect environment for parents to model good nutritional behaviour.²⁸ Food preparation, such as planning, writing a shopping list has been

associated with higher fruit and vegetable intake from a study of women aged 18-65 years.²⁹ Also Traveras et al,³⁰ found that eating a family meal together was inversely associated with obesity in children aged 9 to 14 years old. Children need to see adults eating fruit and vegetables, to help demonstrate that eating fruit and vegetables is beneficial.⁶

Teachers, parents and children had high total appreciation scores for the different intervention materials. The results identified that there was no association between the teachers, parents or children's appreciation scores and fruit and vegetable consumption.

Validity and reliability of the process measures questionnaires has not been tested, however this is a common weakness with health interventions, as limited resources are allocated to process evaluations^{5 13}. Teachers, parents and children might be inclined to give socially desirable answers, leading to overestimation of the intervention effect.

Another limitation is that the study is subject to multiple comparisons or testing, as the more statistical analysis conducted in this case, on the intervention group, such as categorising implementation and appreciation levels, the more likely differences amongst the comparison groups would be found³¹. This study was powered to analyse the main outcome change in fruit and vegetable intake, and as a consequence it may not be adequately powered for this analysis. The main strengths of the present study are that it provides information on process measures of the first multi-component intervention trial to promote fruit and vegetables conducted across England and used advanced statistical techniques.

Conclusion

The Project Tomato intervention was poorly implemented by teachers. The analysis revealed that the school items need improvement, to have a positive association with children's diets. Future intervention research should design activities that involve little preparation time for the classroom teacher. This study also confirmed the importance of parents' involvement, and the home environment. Further research should be conducted to review family eating behaviours and the mealtime environment to facilitate change in children's dietary behaviours. Interventions need activities that parents and children will enjoy together to change children's diets. The next step in research into school interventions is to continue using and validating process evaluations to identify the best methods for conducting health interventions.

What is already known on this subject?

School based fruit and vegetable interventions can increase children's consumption levels, however not all interventions are successful. Process evaluations can be used to improve our understanding of successful health interventions; to identify the key components that achieve an intervention successful, which theoretical constructs made a difference and what environment/conditions lead to these particular components facilitating a successful outcome.

What does this study add?

- This study is one of the few clustered randomised controlled trials, of a multi-component intervention designed to increase children's fruit and vegetable intake conducted in England.

- Parents who eat their main meal with their children at a table at least 3 times per week, have higher fruit and vegetable intakes compared to parents who eat with their children less often.

Acknowledgements

University of Leeds

Dr Siobhan Hugh-Jones, Dr Fiona Jones, Dr Rebecca Lawton, Dr Sinead Boylan, and Miss Jennifer Lewis

National Foundation for Educational Research

Jennifer Jupp, Christine Webster, Frances Reed, Janice Walker and Ben Styles for their work at The National Foundation for Educational Research (NFER) in the sample selection, liaising with the schools and in the collection, entry and distribution of the data.

Completing Interests: None to declare.

Reference List

1. Wells L, Nelson M, Wells L, Nelson M. The National School Fruit Scheme produces short-term but not longer-term increases in fruit consumption in primary school children. *British Journal of Nutrition* 2005;93(4):537-42.
2. Ransley JK, Greenwood DC, Cade JE, Blenkinsop S, Schagen I, Teeman D, et al. Does the school fruit and vegetable scheme improve children's diet? A non-randomised controlled trial. *Journal of Epidemiology & Community Health* 2007;61(8):699-703.
3. Fogarty AW, Antoniak M, Venn AJ, Davies L, Goodwin A, Salfield N, et al. Does participation in a population-based dietary intervention scheme have a lasting impact on fruit intake in young children? *International Journal of Epidemiology* 2007;36(5):1080-5.
4. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL. Parental influences on young girls' fruit and vegetable, micronutrient, and fat intakes. *Journal of the American Dietetic Association* 2002 102(1):58-64.
5. Wind M, Bjelland M, Perez-Rodrigo C, Te Velde SJ, Hildonen C, Bere E, et al. Appreciation and implementation of a school-based intervention are associated with

- changes in fruit and vegetable intake in 10- to 13-year old schoolchildren--the Pro Children study. *Health Education Research* 2008;23(6):997-1007.
6. French SA, Stables G. Environmental interventions to promote vegetable and fruit consumption among youth in school settings. *Preventive Medicine* 2003;37(6 Pt 1):593-610.
 7. Knai C, Pomerleau J, Lock K, McKee M, Knai C, Pomerleau J, et al. Getting children to eat more fruit and vegetables: a systematic review.[see comment]. *Preventive Medicine* 2006;42(2):85-95.
 8. Ciliska D, Miles E, Brien MA, Turl C, Tomasik HH, Donovan U. The effectiveness of community interventions to increase fruit and vegetable consumption in people four years of age and older. *Ontario Ministry of Health Department*. Canada, 1999.
 9. Auld G, C O, Endinger E, M A. Outcomes from a school-based nutrition education program using resource teachers and cross-disciplinary models. . *J. Nutr. Educ* 1998; 30:268-280.
 10. Perry CL, Bishop DB, Taylor G, Murray DM, Mays RW, Dudovitz BS. Changing fruit and vegetable consumption among children: the 5-a-Day Power Plus program in St. Paul, Minnesota. *Am J Public Health* 1998 88:603-9.
 11. Baranowski J, Doyle C, Lin LS, Smith M, Wang DT. Gimme 5 fruit, juice, and vegetables for fun and health: outcome evaluation.[erratum appears in Health Educ Behav 2000 Jun;27(3):390]. *Health Education & Behavior* 2000;27(1):96-111.
 12. Reynolds KD, Franklin FA, Binkley D, Raczynski JM, Harrington KF, Kirk KA, et al. Increasing the fruit and vegetable consumption of fourth-graders: results from the high 5 project. *Preventive Medicine* 2000;30(4):309-19.
 13. Sahota P, Rudolf MC, Dixey R, Hill AJ, Barth JH, Cade J, et al. Randomised controlled trial of primary school based intervention to reduce risk factors for obesity.[see comment]. *BMJ* 2001;323(7320):1029-32.
 14. Perry CL, Bishop DB, Taylor GL, Davis M, Story M, Gray C, et al. A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. *Health Education & Behavior* 2004;31(1):65-76.
 15. Cullen KW, Watson K, Baranowski T, Baranowski JH, Zakeri I, Cullen KW, et al. Squire's Quest: intervention changes occurred at lunch and snack meals. *Appetite* 2005;45(2):148-51.
 16. Bere E, Veierod MB, Klepp K-I. The Norwegian School Fruit Programme: evaluating paid vs. no-cost subscriptions. *Preventive Medicine* 2005;41(2):463-70.
 17. Te Velde SJ, Brug J, Wind M, Hildonen C, Bjelland M, Perez-Rodrigo C, et al. Effects of a comprehensive fruit- and vegetable-promoting school-based intervention in three European countries: the Pro Children Study. *British Journal of Nutrition* 2008;99(4):893-903.
 18. Baranowski T, Anderson C, Carmack C. Mediating variables framework in physical activity interventions: How are we doing? How might we do better? . *America Journal of Preventive Medicine* 1998;15 (4): 266-297.
 19. Steckler A, Linnan L. *Process Evaluation for public health interventions and research* United States of America: Jossey-Bass, 2002.
 20. Kitchen MS, Ransley JK, Greenwood DC, Clarke GP, Conner MT, Jupp JE. Study protocol: a cluster randomised controlled trial of a school based fruit and vegetable intervention - Project Tomato. *BMC Health Services Research* 2009;9(101).
 21. Conner M, Sparks P. The theory of planned behaviour and health behaviour. In: Conner M, Norman P, editors. *Predicting Health behaviour*. Buckingham: Open University Press, 2005:170-222.

22. Cade JE, Frear L, Greenwood DC, Cade JE, Frear L, Greenwood DC. Assessment of diet in young children with an emphasis on fruit and vegetable intake: using CADET-- Child and Diet Evaluation Tool. *Public Health Nutrition* 2006;9(4):501-8.
23. Noble M, Wright G, Dibben C. Indices of Deprivation 2004: Report to the Office of the Deputy Prime Minister. London: Neighbourhood Renewal Unit. , 2004.
24. Story M, Mays RW, Bishop DB, Perry CL, Taylor G, Smyth M, et al. 5-a-day Power Plus: process evaluation of a multicomponent elementary school program to increase fruit and vegetable consumption. *Health Education & Behavior* 2000;27(2):187-200.
25. Tak NI, Te Velde SJ, Brug J, Tak NI, Te Velde SJ, Brug J. Ethnic differences in 1-year follow-up effect of the Dutch Schoolgruitem Project - promoting fruit and vegetable consumption among primary-school children. *Public Health Nutrition* 2007;10(12):1497-507.
26. Bere E, Veierod MB, Bjelland M, Klepp KI. Outcome and process evaluation of a Norwegian school-randomized fruit and vegetable intervention: Fruits and Vegetables Make the Marks (FVMM). *Health Education Research* 2006;21(2):258-67.
27. Blanchette L, Brug J. Determinants of fruit and vegetable consumption among 6-12-year-old children and effective interventions to increase consumption.[see comment]. *Journal of Human Nutrition & Dietetics* 2005;18(6):431-43.
28. Patrick H, Nicklas TA, Patrick H, Nicklas TA. A review of family and social determinants of children's eating patterns and diet quality. *Journal of the American College of Nutrition* 2005;24(2):83-92.
29. Crawford D, Ball K, Mishra G, Salmon J, Timperio A. *Which food-related behaviours are associated with healthier intakes of fruits and vegetables among women?:* Public Health Nutrition. 10(3)(pp 256-265), 2007. Date of Publication: Mar 2007., 2007.
30. Taveras EM, Rifas-Shiman SL, Berkey CS, Rockett HR, Field AE, Frazier AL, et al. Family dinner and adolescent overweight. *Obesity Research* 2005;13(5):900-6.
31. Miller RG. *Simultaneous Statistical Inference* New York: Springer Verlag 1981 .

Table 1. *Intervention demographic characteristics of the respondent and non-respondent participants: Process Measurement Questionnaires*

Characteristics	Completed a questionnaire	95%CI	Did not completed a questionnaire	95%CI
Child characteristics				
Age (years baseline)	7.0	(6.9, 7.0)	7.0	(6.9, 7.1)
%Sex (boys)	132/261 (51%)	(45, 57)	28/50 (56%)	(42, 70)
%Ethnicity (non-white)	33/246 (13%)	(9, 18)	5/45 (11.0%)	(2, 21)
Weight (kg baseline)	24.6	(24.1, 25.1)	24.8	(23.4, 26.3)
Height (cm baseline)	122.5	(121.9, 123.1)	123.3	(121.6, 124.9)
School characteristics				
% Free school meal eligibility*	6.6	(6, 8)	17.6	(13, 25)
IMD score*	13.7	(12.5, 15.1)	23.8	(18.9, 30.0)
Responded group $n = 261$; non-responds group $n = 50$ * $p < 0.001$ (%free school meals eligibility and IMD score are transformed using natural log)				

Table 2 *Teacher's implementation and appreciation of the Project Tomato items (range 0-100)*

Intervention items	Implementation		Appreciation	
	Mean %	95% CI	Mean %	95% CI
Curriculum lessons	45%	(40, 50)	65.8%	(61, 71)
Tasting Sessions	25%	(21, 30)	85.7%	(83, 87)
Cooking Club	8%	(5, 10)	84.5%	(81, 87)
Gardening Club	8%	(6, 11)	46.7%	(42, 51)
All school intervention items	21%	(19, 24)	65.7%	(60, 71)
$N = 24$				

Table 3. Children's implementation and appreciation of home intervention items

Intervention items	Percentage of children who implemented each item			Percentage of children who liked or loved each item		
			95% CI			95% CI
Kit 1						
Bean growing kit	74/181	41%	(34, 49)	121/177	68%	(62, 75)
<i>Juicy science</i>						
Electric Apples	60/181	33%	(26, 40)	87/177	49%	(42, 57)
Fruity Sugars from Apples	63/181	35%	(28, 49)			
Eat5! Quiz book	97/181	54%	(46, 61)	122/180	68%	(61, 75)
Fruity face	148/181	82%	(76, 88)	159/171	93%	(89, 97)
<i>Recipes</i>						
Fruity flapjacks	65/181	36%	(29, 43)	83/179	46%	(38, 54)
Crumpet pizzas	54/181	30%	(23, 37)			
Kit 2						
Balance of good health quiz book	131/236	56%	(49, 62)	156/218	72%	(66, 78)
Christmas cracker	78/213	37%	(30, 43)	177/218	81%	(76, 86)
<i>Juicy science:</i>						
Bad apples	70/218	32%	(26, 38)	110/214	51%	(45, 58)
Scary heads	65/217	30%	(24, 36)			
<i>Christmas recipes:</i>						
Baked vegetables	51/219	23%	(18, 29)	107/214	50%	(43, 57)
Christmas cake	64/219	29%	(23, 35)			
Cabbage and bacon	24/219	11%	(7, 15)			
Healthy apple muffins	46/219	20%	(16, 26)			
Kit 3						
Fruit n veg portion game	178/221	81%	(75, 86)	184/217	85%	(80, 90)
Cress egg head kit	65/221	29%	(23, 36)	139/216	64%	(58, 71)
Fruit "n" veg snack box for fruit	127/221	58%	(50, 64)	192/218	88%	(84, 92)
Fruit "n" veg snack box for vegetables	65/221	29%	(23, 36)			
Easter Bunny carrot cake recipe	51/221	23%	(22, 24)	72/211	36%	(30, 43)
Cheat Easter carrot cake recipe	40/216	19%	(13, 27)	57/204	28%	(22, 34)

Table 4. Implementation of the Project Tomato intervention for Teachers, Children and Parents and the association with total fruit and vegetable intake at follow-up.

Degree of implementation	Coefficient	95% CI	p-value	Coefficient	95% CI	p-value	N
<i>Teachers</i>							
All School intervention items	7.34	(-24.5, 39.1)	0.63	-2.44	(-38.3, 33.4)	0.88	274
Curriculum lessons	-0.84	(-34.4, 32.7)	0.95	-11.54	(-50.1, 26.9)	0.54	274
Tasting Sessions	3.56	(-32.4, 39.5)	0.83	-12.25	(-50.1, 25.5)	0.50	274
<i>Children</i>							
Total Implementation	0.87	(-0.4, 2.1)	0.16	1.10	(-0.2, 2.4)	0.08	115
<i>Parents</i>							
Total Implementation							
Medium	48.63	(-56.4, 153.7)		40.78	(-67.9, 149.5)		
High	56.42	(-35.4, 208.3)	0.34	66.10	(-71.4, 203.7)	0.58	57

Adjusted for baseline fruit and vegetable intake, gender, IMD, and ethnicity.

Table 5. Appreciation of the Project Tomato intervention for Teachers, Children and Parents and the association with total fruit and vegetable intake at follow-up.

Degree of appreciation	Coefficient	95% CI	p-value	Coefficient	95% CI	p-value	N
<i>Teachers</i>							
Curriculum lessons	2.74	(-85.8, 91.3)	0.94	26.20	(-27.2, 79.6)	0.26	75
Tasting Sessions	38.52	(-6.7, 83.7)	0.08	33.12	(-24.5, 90.7)	0.23	145
<i>Children</i>							
Appreciation score	1.51	(-0.6, 4.6)	0.31	0.99	(-1.4, 3.4)	0.38	101
<i>Parents</i>							
Appreciation score							
Medium	-32.36	(-112.3, 47.6)		-17.7	(-80.5, 45.0)		
High	-28.71	(-84.5, 27.1)	0.53	-39.3	(-97.4, 18.7)	0.36	112

Adjusted for baseline fruit and vegetable intake, gender, IMD score, and ethnicity.