

Citation:

Ispoglou, T and Witard, O and Duckworth, L and Lees, M (2020) The efficacy of essential amino acid supplementation for augmenting dietary protein intake in older adults: implications for skeletal muscle mass, strength and function. Proceedings of the Nutrition Society. ISSN 0029-6651 DOI: https://doi.org/10.1017/S0029665120008010

Link to Leeds Beckett Repository record: https://eprints.leedsbeckett.ac.uk/id/eprint/7317/

Document Version: Article (Published Version)

Creative Commons: Attribution 4.0

Copyright © The Authors 2020.

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please contact us and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

CrossMark

Proceedings of the Nutrition Society, Page 1 of 13 doi © The Authors 2020. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nutrition Society Live 2020 was held virtually on 14-15 July 2020

Symposium five: Protein nutrition and ageing

The efficacy of essential amino acid supplementation for augmenting dietary protein intake in older adults: implications for skeletal muscle mass, strength and function

Theocharis Ispoglou¹* ^(D), Oliver C. Witard², Lauren C. Duckworth¹ and Matthew J. Lees¹

¹Carnegie School of Sport, Leeds Beckett University, Leeds LS6 3QS, UK

²School of Basic and Medical Biosciences, King's College London, London, UK

The primary aim of this review is to evaluate the efficacy of essential amino acid (EAA) supplementation as a strategy to increase dietary protein intake and improve muscle mass, strength and function in older adults. A sufficient daily protein intake is widely recognised to be fundamental for the successful management of sarcopenia in older undernourished adults. In practice, optimising protein intakes in older adults is complex, requiring consideration of the dose and amino acid composition (i.e. a complete EAA profile and abundant leucine content) of ingested protein on a per meal basis, alongside the age-related decline in appetite and the satiating properties of protein. Recent studies in older adults demonstrate that EAA-based supplements are non-satiating and can be administered alongside food to enhance the anabolic properties of a meal containing a suboptimal dose of protein; an effect magnified when combined with resistance exercise training. These findings support the notion that EAA supplementation could serve as an effective strategy to improve musculoskeletal health in older adults suffering from non-communicable diseases such as sarcopenia. Compliance is critical for the long-term success of complex interventions. Hence, aspects of palatability and desire to eat are important considerations regarding EAA supplementation. In conclusion, EAA-based supplements enriched with L-leucine offer an alternative strategy to whole protein sources to assist older adults in meeting protein recommendations. In practice, EAA supplements could be administered alongside meals of suboptimal protein content, or alternatively between meals on occasions when older adults achieve their per meal protein intake recommendations.

Leucine: Sarcopenia: Muscle protein synthesis: Ageing

The maintenance of skeletal muscle mass and function with advancing age is fundamental to metabolic health across the lifespan⁽¹⁾, and is primarily achieved through the prevention or improved management of muscle wasting, termed sarcopenia. In essence, sarcopenia is defined as a reduction in muscle mass, strength and function⁽²⁾, and is prevalent in older adults^(3,4). Skeletal muscle also acts as the primary site for insulin-stimulated glucose uptake, and therefore the dysregulation of this body tissue contributes to the development of metabolic syndrome, type II diabetes and increased risk of CVD⁽⁵⁾. Sarcopenia also underpins frailty and the decline in

Downloaded from https://www.cambridge.org/core. IP address: 217.35.243.161, on 05 Feb 2021 at 13:51:51, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/S0029665120008010

Abbreviations: EAA, essential amino acid; MPS, muscle protein synthesis; mTOR, mammalian target of rapamycin; mTORC1, mTOR complex 1. *Corresponding author: Theocharis Ispoglou, email t.ispoglou@leedsbeckett.ac.uk

functional capacity that compromises quality of life⁽⁶⁾, and is associated with a range of clinical disorders, such as osteoporosis and obesity⁽⁷⁾. Although the aetiology of sarcopenia is clearly multifactorial and not fully understood⁽⁸⁾, key underlying factors include morphological changes in skeletal muscle⁽⁹⁾, the loss of motor units⁽¹⁰⁾, physical inactivity⁽¹¹⁾ and, particularly prudent to this review, dietary protein and energy deficiencies⁽¹²⁾.

The combination of adequate dietary protein intake and muscle loading (i.e. resistance) exercise serves as the cornerstone for the management of sarcopenia in both institutionalised and non-institutionalised older adults⁽¹²⁻¹⁷⁾. Another key component of an effective countermeasure for sarcopenia includes adequate energy intake⁽¹⁸⁾, which reduced with ageing⁽¹⁹⁾. Although typically is protein-energy malnutrition and undernutrition are often associated with institutionalised settings⁽²⁰⁻²²⁾</sup> wherein the majority of residents are older adults, protein-energy undernutrition is also prevalent in community-dwelling older populations. According to recent reports, about 8 % of community-dwelling older adults are classified as malnourished and 35 % are at risk of malnutrition⁽²³⁾, whereas 66% of those admitted to a hospital are at risk of malnutrition⁽²⁴⁾. Dependent on the degree of deficiency, protein-energy undernutrition positions older adults at greater risk of other co-morbidities and mortality^(25,26). Given that immune cell function is dependent on energy availability and amino acid provision, protein-energy undernutrition is also implicated in impaired immunity and increased susceptibility to viral infections with advancing age⁽²⁷⁾. Thus, older adults should ensure adequate provision of dietary protein and energy to maintain good health.

Alongside the importance of consuming sufficient daily protein intake, the effective management of sarcopenia also depends on the quantity and quality of protein consumed on a more refined 'per meal' basis⁽²⁸⁾. In this regard, older adults require a higher per meal dose of protein (about 0.4 g/kg body mass) compared to their younger counterparts (about 0.24 g/kg body mass) to maximally stimulate postprandial muscle protein synthesis (MPS) rates⁽²⁹⁾. Protein quality is traditionally assessed by the protein digestibility-corrected amino acid $\text{score}^{(30,31)}$, whereby the digestion and absorption kinetics of essential amino acids (EAA) determine the nutritional quality of a protein source. A more contemporary, and increasingly preferred, tool for assessing protein quality is the digestible indispensable amino acid score since the protein digestibility-corrected amino acid score tends to overestimate the value of lower quality proteins⁽³¹⁾. Of all the EAA, leucine plays a prominent role in regulating muscle metabolism via direct action on the mammalian target of rapamycin (mTOR) complex 1 (mTORC1) signalling pathway that switches on $MPS^{(32-34)}$, an effect that is potentiated by the other EAA⁽³⁴⁾. At the metabolic level, the maintenance of skeletal muscle mass is contingent on muscle protein balance, whereby the rates of muscle protein breakdown are offset against the rates of $MPS^{(35,36)}$. Hence, on a per meal basis, the consumption of adequate, high-quality protein

sources that comprise the full complement of EAA represents an intuitive strategy for the management of sarcopenia. Moreover, there is extensive evidence that combining protein supplementation with resistance exercise training exhibits a synergistic effect on skeletal muscle mass, strength and functional outcomes in older adults^(37,38), with exercise sensitising skeletal muscle to the anabolic action of EAA^(34,39,40). On this basis, lifestyle strategies designed to combat sarcopenia and promote skeletal muscle health in older adults require a holistic approach encompassing both exercise and protein nutrition interventions (Fig. 1).

Protein is the most satiating of all macronutrients⁽⁴¹⁾. Given that older adults commonly experience a reduction in appetite⁽¹⁹⁾, difficulties with chewing, dysphagia⁽⁴²⁾ and suppression of taste and smell⁽⁴³⁾, it may be considered paradoxical to recommend increased protein intakes, potentially at the expense of overall energy intake. Moreover, ageing is associated with reductions in gastric emptying and the splanchnic uptake of amino acids⁽⁴³⁾ that can be further compromised depending on the type of protein ingested $^{(44)}$. As such, older adults are presented with additional challenges in terms of achieving higher daily and per meal protein intakes, which tend to be particularly low at breakfast and lunch⁽⁴⁵⁾. Thus, it is imperative that interventions targeted at increasing total dietary protein intake are not counter-productive in exacerbating energy deficiencies in older adults at risk of negative energy balance, or who would benefit from increasing energy intake.

The aims of this review are 2-fold. First, to critically evaluate the existing evidence base that has explored the efficacy of EAA ingestion as a strategy to increase daily and per meal dietary protein intakes, without compromising intake of other nutrients (macro- and micro-nutrients) in older adults. Secondly, to critically evaluate findings from experimental studies investigating the impact of EAA supplementation on acute (i.e. muscle protein turnover) and chronic (i.e. body composition, strength and muscle function) musculoskeletal health outcomes in older adults, with a view to advancing evidence-based practical guidelines for the management of sarcopenia. By way of introduction, we provide a brief overview regarding the evolution of dietary protein recommendations in the context of managing sarcopenia.

Dietary protein intake and sarcopenia

Re-evaluating the protein RDA in older adults

There is general consensus among the scientific community that daily protein intakes should exceed the current RDA (0.8 g/kg body mass per day) for the management of sarcopenia^(13,15–17) and preservation of muscle mass and function in older adults^(46–50). Consistent with this notion, achieving the current RDA may result in a daily protein intake below the acceptable macronutrient distribution range of about 15–35 %^(51,52). Multiple lines of evidence^(48,53–55) substantiate the recommendation to increase protein intakes above the RDA in older adults. First, although cause and effect cannot be established S Proceedings of the Nutrition Society



Fig. 1. A holistic approach to addressing the challenge of age-associated sarcopenia and optimising skeletal muscle health with reference to dietary protein intake (food and supplementation) and physical activity/exercise.

based on observational studies, Houston *et al.*⁽⁵⁴⁾ documented that healthy older adults consuming a protein intake in excess of the protein RDA were associated with an attenuated decline in muscle mass over a 3 year follow-up period. However, older adults that consumed a protein intake only marginally in excess of the RDA (i.e. up to 1·1 g/kg per day) still suffered a decline in muscle mass. Secondly, a recent carefully controlled study by Mitchell *et al.*⁽⁵⁵⁾ demonstrated that increasing protein intake to twice the RDA (i.e. 1·6 g/kg per day) resulted

in significant gains in lean tissue mass in healthy older men. Furthermore, older individuals have been shown to require a higher per meal protein intake because they are less sensitive to anabolic stimuli such as protein; a concept termed skeletal muscle 'anabolic resistance'^(56,57). Short-term periods of reduced physical activity or muscle disuse (i.e. bed rest or limb immobilisation) have been shown to induce anabolic resistance in older individuals, as indicated by reduced skeletal muscle amino acid transporter expression, impaired mTORC1 signalling, and

blunted postprandial MPS rates in response to EAA⁽⁵⁸⁻⁶⁰⁾. In this regard, and when expressed relative to body mass, older adults require a protein dose of about 0.4 g/kg body mass per meal (i.e. about 30 g protein for an average male) for the maximal postprandial stimulation of MPS, which is about 67% higher than the relative protein dose required for younger adults⁽²⁹⁾. Accordingly, reputable international expert groups such as the European Society for Clinical Nutrition and Metabolism have called for higher protein intakes in older adults⁽⁴⁸⁾ with recommendations set in relation to the current protein RDA: a 25-50% increase for healthy individuals, a 50-90% increase those suffering from acute or chronic disease, and greater than 50% increase above the RDA for those experiencing severe illness or injury. Notwithstanding these latest guidelines, a large proportion of older adults fail to meet the existing RDA in terms of habitual protein intake. For example, in a large cohort of older adults who participated in the National Health and Nutrition Examination Survey, over 30% failed to meet the current protein RDA⁽⁶¹⁾. This observation highlights the practical issues associated with older adults successfully meeting the recommended protein intakes for the management of sarcopenia.

Rationale for dietary protein supplementation in older adults

Achieving a protein intake in excess of the RDA and in accordance with the European Society for Clinical Nutrition and Metabolism guidelines⁽⁴⁸⁾ presents a profound challenge for older adults, given that this population commonly suffers from acute or chronic illness. This concern is exemplified by the high prevalence of malnutrition and anorexia in these compromised populations^(23,24,62–64). In theory, increasing daily protein intake to twice the RDA translates to an 80 kg older adult consuming about 130 g protein daily. Given that protein increases satiety^(65,66) in a dose-dependent manner^(67,68), and appetite is dysregulated in older adults⁽¹⁹⁾, meeting this target daily protein intake is not pragmatic for all older adults. Therefore, strategies to increase protein intake either through whole food or nutritional supplementation should not be at the expense of other nutrients and/or contribute to a reduction in overall energy intake.

The recommendation to increase protein intake while simultaneously maintaining, and in many cases increasing energy intake, can present a 'protein paradox' (Fig. 2). In this regard, dietary supplementation strategies to increase protein intake may unintentionally result in partial energy redistribution^(63,64), which may negatively affect both protein and energy intakes. A reduction in energy intake may be desirable for certain groups (i.e. individuals with obesity), however the focus of this review is on increasing protein intake in older adults without affecting their ability to maintain energy balance.

Protein supplementation alone or in combination with resistance exercise in older adults

Despite compelling proof-of-concept evidence^(53,69) from acute metabolic studies that increasing dietary protein

intake may counter the age-related decline in muscle mass, the efficacy of chronic nutritional interventions to increase dietary protein intake is weak. This observation is particularly relevant when considering studies of protein-based supplementation in the absence of prescribed exercise training. Moreover, limited evidence exists regarding the benefits of protein supplementation v. placebo for maintaining muscle mass in healthy older adults⁽⁷⁰⁾. Multiple factors likely explain these surprising findings, including the duration of the intervention, habitual daily protein intake, and sample size. However, when combining the anabolic stimuli of protein supplementation and resistance exercise training, findings are more consistent in demonstrating improvements in outcome measurements of muscle health in older adults. The capacity of resistance exercise to sensitise muscle to the anabolic potential of dietary protein is primarily achieved through a timely supply of $EAA^{(34,39,40)}$. Accordingly, a meta-analysis by Cermak et al.⁽³⁷⁾ revealed that gains in lean tissue mass were of greater magnitude in both younger and older adults when combining resistance training with protein supplementation v. resistance training alone. Consistent with this observation, Morton *et al.*⁽³⁸⁾ reported that protein supplementation augmented muscle growth during resistance training when habitual dietary protein intakes were, on average, below 1.6 g/kg body mass per day in younger adults. However, the impact of protein supplementation on muscle mass was reduced with advancing age. Interestingly, Cermak *et al.*⁽³⁷⁾ demonstrated similar increases in type II muscle fibre cross-sectional area in older adults following resistance training, irrespective of whether the study administered protein or a placebo. This observation is physiologically relevant, given that type II muscle fibres exhibit the greatest capacity for hypertrophy following resistance training⁽⁷¹⁾, with studies consistently demonstrating smaller type II fibres in both healthy and frail older adults compared with younger adults⁽⁷¹⁻⁷⁴⁾. Moreover, the increased excitation threshold of type II fibres due to their greater innervation by large motor neurons⁽⁷⁵⁾ supports the utility of resistance training⁽⁷⁶⁾ as an exercise mode to recruit these large motor neurons and potentially minimise type II muscle fibre atrophy in older individuals. However, in practice, data from the Health Survey for England and Scotland (1994–2008, *n* 80 306) reported that only 3.4% of individuals meet the strength-based exercise recommendations⁽⁷⁷⁾. Therefore, in addition to optimising protein intake, it is important for older adults to engage in strength-based exercise training to prolong independence, functional capacity and quality of life⁽⁷⁸⁻⁸⁰⁾.

Protein quality and amino acid composition

The absolute and relative intake of dietary protein is an important consideration for supporting skeletal muscle hypertrophy. However, the quality of a given protein source is of additional importance for maintaining muscle mass in the context of healthy ageing and is defined by EAA content and digestibility of the ingested protein^(30,31). Moreover, protein quality as assessed using



Fig. 2. A 'protein paradox' whereby increased protein intake through food or supplementation may not result in an increase in protein intake and compromise energy intake in older adults suffering from protein and energy deficiencies.

the digestible indispensable amino acid score accounts for each individual amino acid as a nutrient in its own right⁽³¹⁾. The amino acid composition of a given protein source can influence the extent and amplitude of postprandial MPS, and induce varying patterns of aminoacidaemia^(81,82). As previously outlined, the leucine content of a given protein source is particularly important in attenuating declines in muscle mass when consumed alongside other EAA⁽⁸³⁾. It follows that older adults require higher amounts of leucine to facilitate the maximal stimulation of MPS in the presence of other EAA⁽⁵⁰⁾. Therefore, it is important that older adults consume high-quality proteins⁽⁸⁴⁾ that provide a full complement of EAA for skeletal muscle health.

Optimising protein intake requires consideration of the following key points: (a) daily protein intake; (b) protein intake on a per meal basis and (c) protein quality as determined by a full and optimal complement of EAA. The broader role of EAA as key regulators of MPS is highlighted by Park *et al.*⁽⁸⁵⁾ who compared the response of MPS to a low dose EAA mixture (3·2 g EAA; 2·4 g protein), a high EAA mixture (6·4 g EAA; 4·4 g protein) and an intact protein supplement containing 17·9 g whey. The combined ingestion of EAA with a low whey protein dose stimulated a greater anabolic response *v*. whey protein alone, a finding partially attributed to the rapid aminoacidaemia observed in the high EAA condition. This observation further supports the notion that

provision of ample dietary EAA and leucine are necessary to support a skeletal muscle anabolic response in older adults. From a practical standpoint, the strategic nutritional supplementation with EAA and leucine alongside meals containing suboptimal protein content (i.e. breakfast and lunch) could assist older adults in achieving their per meal protein intake and EAA profile objectives. Nevertheless, it must also be noted that skeletal muscle is refractory to the anabolic effects of leucine during the postprandial 'muscle-full' period^(86,87). Therefore, in cases where older adults receive adequate high-quality protein with their main meals, it would be prudent that protein-based snacks or supplements are administered between meals when additional nutritional supplementation is required to reach their daily protein goal.

Effects of essential amino acid supplementation on dietary protein intake

In addition to the aforementioned nutritional plan considerations, the age-related anorexia and satiating properties of protein must be taken into account. Although it has been proposed that EAA are not satiating^(88–92), limited studies have investigated the acute^(62,93) or longterm impact of EAA supplementation on dietary intake^(91,94). Acute studies investigating the efficacy of

EAA focus on the measurement of MPS under laboratory conditions where dietary intakes are strictly controlled^(90,92), whereas longer term studies may^(88,91,94) or may not⁽⁸⁹⁾ report habitual protein intakes. To our knowledge, only two studies have investigated the acute effects of EAA-based nutritional supplements, containing an optimum ratio of EAA and leucine, on postprandial rates of MPS^(50,95), appetite and dietary protein intake in concurrent⁽⁹³⁾ or subsequent meals^(62,93). The findings demonstrated that EAA-based supplementation may be an effective strategy to facilitate an increased protein intake without compromising intakes of other macronutrients and ultimately energy intake. Ingestion of a gel-based EAA (7.5 g EAA) $1 h^{(62,93)}$ or immediatelv⁽⁹³⁾ before an *ad lib* breakfast meal significantly increased both protein and energy intakes at the subsequent meal compared to a control condition. Moreover, appetite ratings were not significantly sup-pressed^(62,93) compared to the control condition, while this was not the case for a 15 g EAA-matched whey protein isolate condition⁽⁶²⁾. Energy intake was also increased in the EAA condition compared to the whey protein isolate⁽⁶²⁾, which resulted in an increase of protein intake at the *ad lib* meal but at the expense of energy intake. The postprandial increase in peptide YY in the whey protein condition offers a potential explanation for the suppression of appetite and ultimately reduced energy intake. This observation may be attributed to a greater increase in amino acid concentration in the gastrointestinal tract⁽⁹⁶⁾ compared to the EAA condition, which did not contain non-EAA.

Previous research demonstrates that test meal palatability influences food intake to a greater extent when it is followed by disguised high-energy preloads rather than low energy preloads⁽⁹⁷⁾. Other studies⁽⁹⁸⁾ have also provided insights into the influence of sensory inputs, including palatability, on food intake control. It follows that high-energy supplements administered in close proximity or alongside meals should be palatable to allow for additional food consumption and avoid inadvertent energy redistribution. Compliance to any long-term nutritional supplementation regimen is also critical and will be discussed briefly in the section 'Effects of essential amino acid supplementation alone or in combination with resistance exercise on muscle mass, strength and function'. Ispoglou *et al.*⁽⁹³⁾ considered practical aspects of palatability following different delivery methods of EAA (i.e. a gel or a bar). Participants in their study⁽⁹³⁾ rated both EAA containing supplements (i.e. 7.5 g EAA) as palatable, suggesting that such nutritional prototypes may be adopted long term by older adults who wish to increase their protein intake without reducing food consumption^(97,99). Readily administered forms of supplements (i.e. in gel format) are also critical for enhancing protein intakes in older adults with proteinenergy malnutrition, which appears to exacerbate the rates of dysphagia with ageing⁽⁴²⁾. These findings suggest that palatable forms of EAA supplements may be an efficacious mode of ingestion for enhancing protein intake in older adults with suboptimal protein intakes, a need to maintain or increase protein intake, difficulties

with chewing and dysphagia⁽⁴²⁾ and/or suppression of taste and smell⁽⁴³⁾.

A paucity of studies have investigated the chronic effect of EAA supplementation in augmenting protein intake, and inconsistent approaches have been adopted in relation to the reporting of habitual dietary intakes and habits^(88,91,94). For example, in a 3-month EAA intervention in older obese women, participants were instructed to maintain dietary habits, but dietary intakes were not reported or monitored⁽⁸⁹⁾. Conversely, studies involving EAA supplementation, such as a 10-d bed rest intervention in healthy older adults with strict dietary controls⁽⁸⁸⁾, precludes meaningful conclusions being drawn regarding the efficacy of EAA since the methodological approaches followed in controlled laboratory studies cannot always be applied to real-world settings. It is also not clear if converting nitrogen or amino acid content to protein content is consistently applied. For example, Rondanelli *et al.*⁽⁹¹⁾ reported that the daily protein intake of older institutionalised adults >75 years was not increased with EAA supplementation when accounting for the nitrogen content from EAA (i.e. 8 g daily), despite improvements in nutritional status including a small but non-significant increase in energy intake. Nevertheless, it is unclear how nitrogen content was calculated. It would be judicious to convert nitrogen to protein content whereby this methodological step would have highlighted a substantial increase in the theoretical protein intake (i.e. using a default factor of 5.6, 1 g nitro-gen would require intake of 5.6 g protein)⁽¹⁰⁰⁾. Adding supplemental EAA to total daily protein content⁽⁹¹⁾ is also an inherently flawed approach given that twice the amount of a high-quality protein would be required to supply a given amount of $EAA^{(101)}$. Hence, it could be speculated that daily protein intake as assessed at the end of this 8-week, dietary intervention⁽⁹¹⁾ was probably underestimated if neither nitrogen to protein nor total EAA to protein conversion was implemented. Similarly, daily supplementation with approximately 15 g EAA alongside two main meals for a period of 12 weeks⁽⁹⁴⁾ did not alter habitual energy intakes of healthy older adults. However, accounting for the EAA content in this well-nourished group of older adults, who exceeded recommendations for the current RDA, would have resulted in a daily protein intake closer to 1.5 g/kg/ d⁽¹⁰¹⁾. These studies did not report per meal protein intakes, hence further research is warranted to explore whether concurrent or subsequent mealtime intakes, following EAA supplementation, are impacted over the long term. In summary, the few studies that have investigated the efficacy of EAA to augment protein intake provide preliminary evidence that EAA supplementation does not reduce habitual daily dietary intakes, whereas the increase in daily protein intake is due to the supplement administered. It is currently not known how daily EAA supplementation may affect per meal dietary intakes, since most studies administer supplements between meals, while addressing suboptimal per meal protein intakes is likely to address daily protein deficiencies and ultimately augment skeletal muscle responses and adaptations. Since attrition to a nutritional EAA

NS Proceedings of the Nutrition Society

ly attribu- term response

intervention involving capsules may be primarily attributed to palatability and acceptability issues⁽⁹⁴⁾, special consideration should be given regarding administration methods (i.e. gels, bars and crystalline EAA) that can safeguard the long-term success of the proposed supplementation regimen.

Effects of essential amino acid supplementation alone or in combination with resistance exercise on muscle mass, strength and function

The acute effects of EAA ingestion on muscle protein turnover have been extensively studied^(85,95,102-106) and provide robust evidence for the role of EAA as key regulators of MPS. However, limited information exists regarding the chronic impact of EAA supplementation in modulating muscle mass, strength and function in older adults. Consistent with this notion, two recent systematic reviews^(70,107) summarised the longterm effects of protein-based supplementation, identifying only seven eligible intervention studies using $EAA^{(88,89,91,94,108,109)}$. As such, these systematic reviews reveal equivocal findings regarding the impact of EAAbased supplementation on muscle mass, strength and function in healthy older adults. Methodological differences inevitably exist between studies with regard to the provision of intact protein sources, individual amino acids or amino acid metabolites that may explain this discord. For instance, two studies administered leucine, which in isolation appears to be ineffective at promoting gains in lean tissue mass^(110,111). Crucially, the anabolic potential of leucine is only observed in the presence of the other amino acids and in particular the EAA. since a sustained increase in MPS requires the availability of all EAA^(34,112). Nevertheless, it is vital that EAA mixtures include a relatively high ratio of leucine to achieve maximal stimulation of MPS in older adults⁽⁹⁵⁾.

Most interventions involving EAA only supplementation that report a beneficial effect on muscle mass and functional outcomes have been conducted in patients with obesity and/or sarcopenia^(89,91,109,113,114), with a single study conducted in glucose-intolerant older men and women⁽¹¹⁵⁾. This observation emphasises the need for targeted interventions in clinical older populations with underlying sarcopenia and/or obesity. A shared methodological aspect across the majority of these studies is that EAA supplements are administered between meals rather than alongside meals. A major practical implication of this approach may be that older adults with suboptimal per meal protein intakes still fail to meet their per meal protein recommendations, thus the degree and rate of skeletal muscle adaptations may be negatively influenced over the long term. However, in cases where older adults meet per meal protein recommendations, it would be practical to ingest supplements between meals, since rates of MPS would not be increased further due to the 'muscle-full' $effect^{(87)}$.

In view of the fact that none of the studies discussed here have reported per meal protein intakes, we can only speculate regarding the potential impact on longterm responses and adaptations. Accordingly, in older institutionalised adults, twice-daily supplementation with 4 g EAA (8 g in total, with EAA given as betweenmeal snacks) was associated with improvements in quality of life, muscle function and diet profile as assessed by the mini nutrition assessment tool⁽⁹¹⁾. Consistent with this finding, whole body lean tissue mass was increased following 18 months of daily supplementation with 16 g EAA $(2 \times 8 \text{ g as between-meal snacks})^{(113)}$ in older adults with sarcopenia, whereas insulin sensitivity was improved and insulin-like growth factor 1 was significantly increased. Although MPS was not measured in this study⁽¹¹³⁾, we speculate that the increase in insulinlike growth factor 1 with EAA supplementation likely increased the stimulation of MPS. Support for this notion also stems from a study by Dillon *et al.*⁽⁸⁹⁾, sedentary older women with obesity, who showed that enhanced rates of basal MPS in an EAA-treated group were consistent with increases in lean body mass and protein expression of skeletal muscle insulin-like growth factor 1. The supplementation regimen consisted of a twice-daily (7.5 g) dose of EAA (15 g total daily) containing 18.6% L-leucine, and resulted in a marked increase in lean body mass (3.9%) following 3 months of EAA supplementation⁽⁸⁹⁾, despite the absence of strength gains. In a study focusing on women with sarcopenia⁽¹⁰⁹⁾, supplementation with 3 g leucine-enriched EAA twice-daily (6 g in total) for a period of 3 months did not result in strength or functional performance improvements, other than walking speed, compared to baseline or a placebo condition. In a similar investigation, with strict dietary controls in place, ingestion of 10 g EAA as snacks twice-daily (20 g in total) over 28 months was shown to increase appendicular skeletal muscle index in patients with sarcopenia and obesity⁽¹¹⁴⁾, however no functional outcome measures were reported. Furthermore, Borsheim et al.⁽¹¹⁵⁾, demonstrated that twice-daily supplementation with 11 g EAA including L-arginine (in total 22 g/d) between meals over a 16 week period improved performance and physical function in older adults with glucose intolerance and adequate daily protein intake. In this study, lean body mass was increased by about 1 kg compared to baseline at week 12, but not at week 16 of the intervention. Habitual energy and protein intakes remained unchanged compared to baseline. Nevertheless, the addition of the EAA supplement would have resulted in a daily increase in total protein intake to a level considered optimal for this population. It is unknown whether supplementation enabled participants to meet their per meal protein recommendations, since supplements were ingested between meals rather than alongside meals.

Despite an overall positive impact of supplementation with EAA on muscle mass in populations with obesity, sarcopenia, or compromised health status, positive changes in strength and functional performance are not always evident. Methodological differences (i.e. supplementation between meals), ratio and amount(s) of amino acids used, and lack of exercise interventions alongside EAA supplementation are all potential factors that can explain the divergent findings.

Fewer studies have been conducted in healthy older adults than older adults with sarcopenia or obesity, and these studies tend to utilise more diverse methodological approaches. Nonetheless, preliminary evidence suggests that supplementation with EAA may contribute to functional and strength gains, whereas evidence is weaker with regard to muscle mass increases. For instance, Markofski et al.⁽¹¹⁶⁾ supplemented healthy older adults with EAA over a 24 week period and failed to detect any increases in total lean body mass, regardless of an increase before and at the end of the intervention in acute EAA-stimulated rates of MPS. Furthermore, neither functional nor strength gains were observed, giving support to the notion that supplementation of EAA alone does not confer any additional benefits. In this study, although daily (15 g) EAA supplementation was strategically timed between meals, it remains unknown whether participants met recommendations for protein intake on a daily and/or per meal basis, since these outcomes were not measured. Conversely, during a 10 d bed rest study in healthy older adults when dietary protein intake was increased to about 1.4 g/kg per day $(3 \times 15 \text{ g})$ EAA) during energy balance⁽⁸⁸⁾, muscle function was maintained. Nevertheless, muscle mass was reduced despite an increase in protein intake and maintenance of MPS rates, which supports the idea that preservation of muscle function may be due to the maintenance of more functional fibres during inactivity. A study in older adults with no underlying health conditions demonstrated both functional performance gains and an increase in lean tissue mass during 3 months of EAA supplementation while adopting a slightly different methodological approach; participants consumed about 7.5 g leucine-enriched (40%) EAA twice daily (total of 0.21 g/kg per day or about 15 g daily)⁽⁹⁴⁾. Previous research has demonstrated that physical (in)activity and protein intake may contribute to a blunted response of MPS^(58,117,118). Thus, it is intuitive that when dietary protein intake is sufficient to maximise MPS, additional supplementation may not confer any muscle anabolic advantage due to a 'muscle-full' effect⁽⁸⁷⁾. In practice, if participants were already receiving adequate high-quality protein with their main meals, nutritional supplementation would not have been of additional benefit, since the required daily protein goal was already reached. Furthermore, in terms of compliance, participants in the study by Ispoglou *et al.*⁽⁹⁴⁾ managed only 75–85 % of the prescribed dose of EAA due to palatability issues and the practicality of ingesting multiple (about 30) capsules on a daily basis. Compliance issues associated with tolerating high numbers of capsules raises important questions regarding long-term compliance to EAA supplementation studies and protein supplementation trials in general, whereby compliance has been shown to be as low as 43 $\%^{(119)}$. Strength and functional performance improvements have also been observed in older adults with reduced physical activity levels following a daily supplementation regimen of 3×4 g EAA over a 3 month period⁽¹⁰⁸⁾. It is worth acknowledging that benefits associated with EAA supplementation may not be manifested at a skeletal muscle level but rather accrue

in other physiological domains (i.e. immunity) and conditions (i.e. energy deficit, catabolic state and injury) that are beyond the scope of this review⁽¹²⁰⁾.

It should be reiterated that sensitivity to the anabolic potential of EAA increases in the presence of exercise, largely through the synergistic influence of mTOR⁽¹²¹⁻¹²³⁾. Nevertheless, evidence from acute metabolic studies is equivocal with regard to the efficacy of combining EAA supplementation with resistance exercise to stimulate MPS in older adults. Early research demonstrated a similar, albeit delayed, acute MPS response in older compared with younger adults following the ingestion of 20 g EAA 1 h after cessation of bilateral leg extension exercise (eight sets of ten repetitions; 70% of one repetition maximum strength)⁽⁵⁹⁾. Moreover, the phosphorylation status of key intramyocellular signalling targets in the mTORC1 pathway (i.e. mTOR, S6K1, 4E-BP1 and eEF2) did not differ between age groups. In contrast, other studies have observed a blunting of the mTORC1 pathway⁽¹²⁴⁾, when EAA were ingested following resistance exercise. A comparable profile of aminoacidaemia. mTORC1 signalling. and/or MPS has also been documented, irrespective of age^(39,125–127). Therefore, it has been suggested that optimising both anabolic stimuli (i.e. EAA/dietary protein and resistance exercise) may effectively counter anabolic resistance in older adults⁽¹²⁸⁻¹³¹⁾ and augment anabolic responses and adaptations.

Longitudinal studies of combined EAA and resistance exercise interventions in older adults demonstrate inconsistent findings regarding skeletal muscle mass and function. A systematic review of sixteen trials reported beneficial effects of combined EAA or protein supplementation and resistance training on body composition $(n \ 3 \ studies)$, strength $(n \ 3 \ studies)$ and physical performance $(n \ 2 \ \text{studies})$ in older adults⁽¹³²⁾. This review concluded that individuals who suffer from sarcopenia and/or frailty could benefit more from combined resistance exercise and EAA or protein supplementation strategies than healthy older adults. However, it should be highlighted that only one study⁽¹⁰⁹⁾ administered free form EAA, variable methodologies were employed between studies, and no study reported habitual protein intakes which together precludes a definitive conclusion regarding the efficacy of EAA ingestion on muscle mass and function in healthy older adults. Kim $et \ al.^{(109)}$ reported increased knee extension strength (9.3%) in community-dwelling older Japanese women with sarcopenia following twice-daily ingestion of a leucine-enriched EAA mixture (6 g daily; 42 % leucine) alongside twice-weekly comprehensive physical training (incorporating resistance-type activities) for 3 months. Leg muscle mass increased in the EAA and exercise group as well as the exercise only group; however, it is noteworthy that EAA supplementation was required to facilitate improvements in knee extension strength. An important population that could benefit from resistance training combined with EAA supplementation is older adults with sarcopenia recovering from injury. In a randomised controlled trial which recruited patients with osteoporotic hip fractures, 2 months of daily EAA (8 g

total daily; about 30 % leucine) supplementation in combination with a physical exercise programme including resistance-type activities significantly improved handgrip strength, timed up and go performance, and Iowa Level of Assistance score only in individuals with sarcopenia⁽¹³³⁾. Therefore, existing evidence indicates that older adults recovering from potentially devastating injuries, such as hip fractures, may improve their functional strength by consuming EAA and engaging in resistancetype exercise activities.

Conclusions and future directions

NS Proceedings of the Nutrition Society

Ageing *per se* is associated with sarcopenia, which is exacerbated in older adults with reduced physical activity and malnutrition. The prevalence of sarcopenia is high in institutionalised settings (i.e. care homes), and in older adults with a range of clinical disorders (i.e. obesity, osteoporosis and cancer) or those recovering from disease and injury. Short-term inactivity due to injury or hospitalisation may result in anabolic resistance and further exacerbate rates of sarcopenia in ageing populations. Therefore, optimising energy and protein intakes in older adults is paramount to the successful management and prevention of sarcopenia. Nevertheless, older adults encounter challenges such as dysphagia and suppression of appetite, taste and smell. As such, addressing energy and protein deficiencies by consumption of highquality proteins on a daily and per meal basis offer evidence-based strategies for managing sarcopenia. However, the satiating properties of protein may create a 'protein paradox' whereby an increase in protein intake may be achieved at the expense of other vital nutrients due to partial energy redistribution. This in turn will likely affect the anabolic potential of the main meals (i. e. breakfast, lunch and dinner) and over the long-term compromise skeletal muscle health. We propose that EAA-based supplements enriched with L-leucine may be an alternative strategy to whole protein sources for assisting older adults to meet daily, and more importantly per meal, protein recommendations since the current evidence regarding the long-term impact on muscle mass and function is based primarily on underpowered observational studies. This strategy is particularly relevant during the periods of muscle disuse (i.e. before, during and after hospitalisation) when skeletal muscle becomes less sensitised to anabolic stimuli such as protein. Since resistance exercise enhances the anabolic potential of EAA, exercise interventions should complement, where appropriate, protein-based nutritional interventions to maximise MPS. We specifically emphasise the need to ingest EAA-based supplements alongside meals with suboptimum amounts of protein. However, supplements may be given between meals in cases where older adults receive ample protein with their main meals. Despite the emphasis of this review on maintenance of muscle mass and function, we should not underestimate the role of EAA on positively influencing other aspects of health, such as metabolic health and immunity. In this regard, future research is warranted

to explore the feasibility and efficacy of ingesting EAA on addressing per meal and daily protein recommendations in clinical (i.e. prehabilitation and rehabilitation) and institutionalised settings, where rates of sarcopenia, frailty, undernutrition and malnutrition are highly prevalent.

Acknowledgements

We wish to express our sincere thanks to the Nutrition Society for the invitation to contribute to the Proceedings of the Nutrition Society following the Nutrition Society Live 2020 event.

Financial Support

None.

Conflict of Interest

A published European Patent Application (No. EP14734541.7, status pending) by one of the authors (T. I.); the associated study was not financially supported by any external commercial organisation.

Authorship

The authors had joint responsibility for all aspects of preparation of this paper.

References

- 1. Wolfe RR (2006) The underappreciated role of muscle in health and disease. *Am J Clin Nutr* **84**, 475–482.
- 2. Cruz-Jentoft AJ, Bahat G, Bauer J *et al.* (2018) Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* **48**, 16–31.
- 3. Shafiee G, Keshtkar A, Soltani A *et al.* (2017) Prevalence of sarcopenia in the world: a systematic review and metaanalysis of general population studies. *J Diabetes Metab Disord* **16**, 21.
- 4. Mayhew AJ, Amog K, Phillips S *et al.* (2019) The prevalence of sarcopenia in community-dwelling older adults, an exploration of differences between studies and within definitions: a systematic review and meta-analyses. *Age Ageing* **48**, 48–56.
- Stump CS, Henriksen EJ, Wei Y et al. (2006) The metabolic syndrome: role of skeletal muscle metabolism. Ann Med 38, 389–402.
- 6. Beaudart C, Zaaria M, Pasleau F *et al.* (2017) Health outcomes of sarcopenia: a systematic review and meta-analysis. *PLoS ONE* **12**, e0169548.
- Kuh D, Karunananthan S, Bergman H et al. (2014) A life-course approach to healthy ageing: maintaining physical capability. Proc Nutr Soc 73, 237–248.
- 8. Morley JE (2012) Sarcopenia in the elderly. *Fam Pract* **29**, i44–i48.
- 9. Patel HP, White MC, Westbury L *et al.* (2015) Skeletal muscle morphology in sarcopenia defined using the EWGSOP criteria: findings from the Hertfordshire Sarcopenia Study (HSS). *BMC Geriatr* **15**, 171–171.

10

NS Proceedings of the Nutrition Society

- Piasecki M, Ireland A, Stashuk D *et al.* (2016) Age-related neuromuscular changes affecting human vastus lateralis. *J Physiol* **594**, 4525–4536.
- 11. Saoi M, Li A, McGlory C *et al.* (2019) Metabolic perturbations from step reduction in older persons at risk for sarcopenia: plasma biomarkers of abrupt changes in physical activity. *Metabolites* **9**, 134.
- Murphy CH & Roche HM (2018) Nutrition and physical activity countermeasures for sarcopenia: time to get personal? *Nutr Bull* 43, 374–387.
- 13. Lang T, Streeper T, Cawthon P *et al.* (2010) Sarcopenia: etiology, clinical consequences, intervention, and assessment. *Osteoporos Int* **21**, 543–559.
- 14. Lieffers J, Bathe O, Fassbender K *et al.* (2012) Sarcopenia is associated with postoperative infection and delayed recovery from colorectal cancer resection surgery. *Br J Cancer* **107**, 931.
- 15. Naseeb MA & Volpe SL (2017) Protein and exercise in the prevention of sarcopenia and aging. *Nutr Res* **40**, 1–20.
- 16. Thalacker-Mercer AE & Drummond MJ (2014) The importance of dietary protein for muscle health in inactive, hospitalized older adults. *Ann N Y Acad Sci* **1328**, 1–9.
- 17. Baum JI, Kim I-Y & Wolfe RR (2016) Protein consumption and the elderly: what is the optimal level of intake? *Nutrients* **8**, 359.
- Morley JE, Argiles JM, Evans WJ *et al.* (2010) Nutritional recommendations for the management of sarcopenia. *J Am Med Dir Assoc* 11, 391–396.
- 19. Johnson KO, Shannon OM, Matu J et al. (2020) Differences in circulating appetite-related hormone concentrations between younger and older adults: a systematic review and meta-analysis. Aging Clin Exp Res 32, 1233–1244.
- Elia M (2006) Nutrition and health economics. *Nutrition* 22, 576–578.
- 21. Elia M (2015) *The Cost of Malnutrition in England and Potential Cost Savings from Nutritional Interventions (full report)*. Malnutrition Action Group of BAPEN and National Institute for Health Research Southampton Biomedical Research Centre.
- Kaiser MJ, Bauer JM, Rämsch C et al. (2010) Frequency of malnutrition in older adults: a multinational perspective using the mini nutritional assessment. J Am Geriatr Soc 58, 1734–1738.
- Rist G, Miles G & Karimi L (2012) The presence of malnutrition in community-living older adults receiving home nursing services. *Nutr Diet* 69, 46–50.
- Geurden B, Franck E, Weyler J et al. (2015) The risk of malnutrition in community-living elderly on admission to hospital for major surgery. Acta Chir Belg 115, 341– 347.
- Sullivan DH, Bopp MM & Roberson PK (2002) Protein-energy undernutrition and life-threatening complications among the hospitalized elderly. *J Gen Intern Med* 17, 923–932.
- Sullivan DH & Walls RC (1995) The risk of lifethreatening complications in a select population of geriatric patients: the impact of nutritional status. J Am Coll Nutr 14, 29–36.
- Alam I, Almajwal AM, Alam W et al. (2019) The immune-nutrition interplay in aging – facts and controversies. Nutr Healthy Aging 5, 73–95.
- Murphy CH, Oikawa SY & Phillips SM (2016) Dietary protein to maintain muscle mass in aging: a case for per-meal protein recommendations. *J Frailty Aging* 5, 49–58.

- 29. Moore DR, Churchward-Venne TA, Witard O et al. (2015) Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. J Gerontol A Biol Sci Med Sci 70, 57–62.
- Millward DJ, Layman DK, Tomé D et al. (2008) Protein quality assessment: impact of expanding understanding of protein and amino acid needs for optimal health. Am J Clin Nutr 87, 1576s–1581s.
- 31. Phillips SM (2016) The impact of protein quality on the promotion of resistance exercise-induced changes in muscle mass. *Nutr Metab* **13**, 64.
- 32. Atherton PJ, Smith K, Etheridge T *et al.* (2010) Distinct anabolic signalling responses to amino acids in C2C12 skeletal muscle cells. *Amino Acids* **38**, 1533–1539.
- 33. Kimball SR, Shantz LM, Horetsky RL *et al.* (1999) Leucine regulates translation of specific mRNAs in L6 myoblasts through mTOR-mediated changes in availability of eIF4E and phosphorylation of ribosomal protein S6. *J Biol Chem* **274**, 11647–11652.
- 34. Moberg M, Apró W, Ekblom B et al. (2016) Activation of mTORC1 by leucine is potentiated by branched-chain amino acids and even more so by essential amino acids following resistance exercise. Am J Physiol Cell Physiol 310, C874–C884.
- Trommelen J, Betz MW & van Loon LJC (2019) The muscle protein synthetic response to meal ingestion following resistance-type exercise. *Sports Med* 49, 185–197.
- Murton AJ (2015) Muscle protein turnover in the elderly and its potential contribution to the development of sarcopenia. *Proc Nutr Soc* 74, 387–396.
- Cermak NM, Res PT, Groot LC et al. (2012) Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a metaanalysis. Am J Clin Nutr 96, 1454–1464.
- 38. Morton RW, Murphy KT, McKellar SR et al. (2018) A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. Br J Sports Med 52, 376–384.
- 39. Lees MJ, Wilson OJ, Webb EK et al. (2020) Novel essential amino acid supplements following resistance exercise induce aminoacidemia and enhance anabolic signaling irrespective of age: a proof-of-concept trial. Nutrients 12, 7.
- 40. Rasmussen BB, Tipton KD, Miller SL *et al.* (2000) An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* **88**, 386–392.
- 41. Poppitt SD, McCormack D & Buffenstein R (1998) Short-term effects of macronutrient preloads on appetite and energy intake in lean women. *Physiol Behav* 64, 279–285.
- 42. Hudson HM, Daubert CR & Mills RH (2000) The interdependency of protein-energy malnutrition, aging, and dysphagia. *Dysphagia* **15**, 31–38.
- 43. Landi F, Calvani R, Tosato M et al. (2016) Anorexia of aging: risk factors, consequences, and potential treatments. *Nutrients* **8**, 69.
- Boirie Y, Dangin M, Gachon P et al. (1997) Slow and fast dietary proteins differently modulate postprandial protein accretion. Proc Natl Acad Sci U S A 94, 14930–14935.
- 45. Tieland M, Borgonjen-Van den Berg KJ, Van Loon LJ et al. (2015) Dietary protein intake in Dutch elderly people: a focus on protein sources. *Nutrients* 7, 9697– 9706.
- 46. Bauer J, Biolo G, Cederholm T *et al.* (2013) Evidencebased recommendations for optimal dietary protein intake

in older people: a position paper from the PROT-AGE study group. J Am Med Dir Assoc 14, 542-559.

- 47. Campbell WW, Trappe TA, Wolfe RR *et al.* (2001) The recommended dietary allowance for protein may not be adequate for older people to maintain skeletal muscle. *J Gerontol A Biol Sci Med Sci* **56**, M373–M380.
- 48. Deutz NE, Bauer JM, Barazzoni R *et al.* (2014) Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN expert group. *Clin Nutr* **33**, 929–936.
- 49. Lonnie M, Hooker E, Brunstrom JM *et al.* (2018) Protein for life: review of optimal protein intake, sustainable dietary sources and the effect on appetite in ageing adults. *Nutrients* **10**, 360.
- 50. Phillips SM, Chevalier S & Leidy HJ (2016) Protein 'requirements' beyond the RDA: implications for optimizing health. *Appl Physiol Nutr Metab* **41**, 565–572.
- 51. Wolfe RR, Miller SL & Miller KB (2008) Optimal protein intake in the elderly. *Clin Nutr* **27**, 675–684.
- 52. Manore MM (2005) Exercise and the institute of medicine recommendations for nutrition. *Curr Sports Med Rep* **4**, 193–198.
- 53. Kim I-Y, Schutzler S, Schrader A et al. (2015) Quantity of dietary protein intake, but not pattern of intake, affects net protein balance primarily through differences in protein synthesis in older adults. Am J Physiol Endocrinol Metab 308, E21–E28.
- 54. Houston DK, Nicklas BJ, Ding J et al. (2008) Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the health, aging, and body composition (health ABC) study. Am J Clin Nutr 87, 150–155.
- 55. Mitchell CJ, Milan AM, Mitchell SM *et al.* (2017) The effects of dietary protein intake on appendicular lean mass and muscle function in elderly men: a 10-wk randomized controlled trial. *Am J Clin Nutr* **106**, 1375–1383.
- Rennie MJ (2009) Anabolic resistance: the effects of aging, sexual dimorphism, and immobilization on human muscle protein turnover. *Appl Physiol Nutr Metab* 34, 377–381.
- 57. Breen L & Phillips SM (2011) Skeletal muscle protein metabolism in the elderly: interventions to counteract the 'anabolic resistance' of ageing. *Nutr Metab* **8**, 68.
- Breen L, Stokes KA, Churchward-Venne TA *et al.* (2013) Two weeks of reduced activity decreases leg lean mass and induces 'anabolic resistance' of myofibrillar protein synthesis in healthy elderly. *J Clin Endocrinol Metab* 98, 2604–2612.
- 59. Drummond MJ, Dreyer HC, Pennings B et al. (2008) Skeletal muscle protein anabolic response to resistance exercise and essential amino acids is delayed with aging. J Appl Physiol 104, 1452–1461.
- 60. Wall BT, Gorissen SH, Pennings B *et al.* (2015) Aging is accompanied by a blunted muscle protein synthetic response to protein ingestion. *PLoS ONE* **10**, e0140903.
- Krok-Schoen JL, Archdeacon Price A, Luo M et al. (2019) Low dietary protein intakes and associated dietary patterns and functional limitations in an aging population: a NHANES analysis. J Nutr Health Aging 23, 338–347.
- 62. Butterworth M, Lees M, Harlow P *et al.* (2019) Acute effects of essential amino acid gel-based and whey protein supplements on appetite and energy intake in older women. *Appl Physiol Nutr Metab* **44**, 1141–1149.
- 63. Fiatarone Singh MA, Bernstein MA, Ryan AD et al. (2000) The effect of oral nutritional supplements on habitual dietary quality and quantity in frail elders. J Nutr Health Aging 4, 5–12.

- 64. Cramer JT, Cruz-Jentoft AJ, Landi F *et al.* (2016) Impacts of high-protein oral nutritional supplements among malnourished men and women with sarcopenia: a multicenter, randomized, double-blinded, controlled trial. J Am Med Dir Assoc 17, 1044–1055.
- 65. Veldhorst M, Smeets A, Soenen S *et al.* (2008) Proteininduced satiety: effects and mechanisms of different proteins. *Physiol Behav* **94**, 300–307.
- 66. Mollahosseini M, Shab-Bidar S, Rahimi MH et al. (2017) Effect of whey protein supplementation on long and short term appetite: a meta-analysis of randomized controlled trials. *Clin Nutr ESPEN* 20, 34–40.
- Veldhorst MA, Nieuwenhuizen AG, Hochstenbach-Waelen A et al. (2009) Dose-dependent satiating effect of whey relative to casein or soy. *Physiol Behav* 96, 675–682.
- 68. Paddon-Jones D & Leidy H (2014) Dietary protein and muscle in older persons. *Curr Opin Clin Nutr Metab Care* 17, 5–11.
- Yang Y, Breen L, Burd NA *et al.* (2012) Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 108, 1780–1788.
- 70. Tieland M, Franssen R, Dullemeijer C et al. (2017) The impact of dietary protein or amino acid supplementation on muscle mass and strength in elderly people: individual participant data and meta-analysis of RCT's. J Nutr Health Aging 21, 994–1001.
- Verdijk LB, Gleeson BG, Jonkers RAM *et al.* (2009) Skeletal muscle hypertrophy following resistance training is accompanied by a fiber type-specific increase in satellite cell content in elderly men. *J Gerontol A Biol Sci Med Sci* 64, 332–339.
- 72. Karlsen A, Bechshøft RL, Malmgaard-Clausen NM et al. (2019) Lack of muscle fibre hypertrophy, myonuclear addition, and satellite cell pool expansion with resistance training in 83–94-year-old men and women. Acta Physiol 227, e13271.
- 73. Yu F, Hedström M, Cristea A *et al.* (2007) Effects of ageing and gender on contractile properties in human skeletal muscle and single fibres. *Acta Physiol* **190**, 229–241.
- 74. Kosek DJ, Kim JS, Petrella JK *et al.* (2006) Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J Appl Physiol* 101, 531–544.
- 75. Sale DG (1987) Influence of exercise and training on motor unit activation. *Exerc Sport Sci Rev* **15**, 95–151.
- 76. Häkkinen K, Newton RU, Gordon SE et al. (1998) Changes in muscle morphology, electromyographic activity, and force production characteristics during progressive strength training in young and older men. J Gerontol A Biol Sci Med Sci 53, B415–B423.
- 77. Stamatakis E, Lee IM, Bennie J *et al.* (2018) Does strength-promoting exercise confer unique health benefits? A pooled analysis of data on 11 population cohorts with all-cause, cancer, and cardiovascular mortality endpoints. *Am J Epidemiol* **187**, 1102–1112.
- Liu CJ & Latham NK (2009) Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev* 3, 1–213.
- 79. Mayer F, Scharhag-Rosenberger F, Carlsohn A *et al.* (2011) The intensity and effects of strength training in the elderly. *Dtsch Arztebl Int* **108**, 359–364.
- 80. Krist L, Dimeo F & Keil T (2013) Can progressive resistance training twice a week improve mobility, muscle strength, and quality of life in very elderly nursing-home residents with impaired mobility? A pilot study. *Clin Interv Aging* 8, 443–448.

- 81. Franzke B, Neubauer O, Cameron-Smith D *et al.* (2018) Dietary protein, muscle and physical function in the very Old. *Nutrients* **10**, 935.
- 82. Traylor DA, Gorissen SH & Phillips SM (2018) Perspective: protein requirements and optimal intakes in aging: are we ready to recommend more than the recommended daily allowance? *Adv Nutr* 9, 171–182.
- mended daily allowance? Adv Nutr 9, 171–182.
 83. Devries MC, McGlory C, Bolster DR et al. (2018) Leucine, not total protein, content of a supplement is the primary determinant of muscle protein anabolic responses in healthy older women. J Nutr 148, 1088–1095.
- 84. Hamarsland H, Laahne JAL, Paulsen G et al. (2017) Native whey induces higher and faster leucinemia than other whey protein supplements and milk: a randomized controlled trial. *BMC Nutr* **3**, 10.
- 85. Park S, Church DD, Azhar G et al. (2020) Anabolic response to essential amino acid plus whey protein composition is greater than whey protein alone in young healthy adults. J Int Soc Sports Nutr 17, 9.
- 86. Atherton PJ, Etheridge T, Watt PW et al. (2010) Muscle full effect after oral protein: time-dependent concordance and discordance between human muscle protein synthesis and mTORC1 signaling. Am J Clin Nutr 92, 1080–1088.
- Mitchell WK, Phillips BE, Hill I *et al.* (2017) Human skeletal muscle is refractory to the anabolic effects of leucine during the postprandial muscle-full period in older men. *Clin Sci (Lond)* 131, 2643–2653.
- Ferrando AA, Paddon-Jones D, Hays NP *et al.* (2010) EAA supplementation to increase nitrogen intake improves muscle function during bed rest in the elderly. *Clin Nutr* 29, 18–23.
- Dillon EL, Sheffield-Moore M, Paddon-Jones D et al. (2009) Amino acid supplementation increases lean body mass, basal muscle protein synthesis, and insulin-like growth factor-I expression in older women. J Clin Endocrinol Metab 94, 1630–1637.
- Paddon-Jones D, Sheffield-Moore M, Aarsland A et al. (2005) Exogenous amino acids stimulate human muscle anabolism without interfering with the response to mixed meal ingestion. Am J Physiol Endocrinol Metab 288, E761–E767.
- Rondanelli M, Opizzi A, Antoniello N et al. (2011) Effect of essential amino acid supplementation on quality of life, amino acid profile and strength in institutionalized elderly patients. *Clin Nutr* **30**, 571–577.
- 92. Bukhari SS, Phillips BE, Wilkinson DJ et al. (2015) Intake of low-dose leucine-rich essential amino acids stimulates muscle anabolism equivalently to bolus whey protein in older women at rest and after exercise. Am J Physiol Endocrinol Metab 308, E1056–E1065.
- 93. Ispoglou T, Deighton K, King RF et al. (2017) Novel essential amino acid supplements enriched with L-leucine facilitate increased protein and energy intakes in older women: a randomised controlled trial. Nutr J 16, 75.
- 94. Ispoglou T, White H, Preston T *et al.* (2016) Doubleblind, placebo-controlled pilot trial of L-leucine-enriched amino-acid mixtures on body composition and physical performance in men and women aged 65–75 years. *Eur J Clin Nutr* **70**, 182–188.
- 95. Katsanos CS, Kobayashi H, Sheffield-Moore M et al. (2006) A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. Am J Physiol Endocrinol Metab 291, E381–E387.
- Moran TH & Dailey MJ (2011) Intestinal feedback signaling and satiety. *Physiol Behav* 105, 77–81.

- 97. Yeomans MR, Lee MD, Gray RW et al. (2001) Effects of test-meal palatability on compensatory eating following disguised fat and carbohydrate preloads. Int J Obes Relat Metab Disord 25, 1215–1224.
- McCrickerd K & Forde CG (2016) Sensory influences on food intake control: moving beyond palatability. *Obes Rev* 17, 18–29.
- 99. Johnson F & Wardle J (2014) Variety, palatability, and obesity. *Adv Nutr* 5, 851–859.
- 100. Mariotti F, Tomé D & Mirand PP (2008) Converting nitrogen into protein--beyond 6.25 and Jones' factors. *Crit Rev Food Sci Nutr* 48, 177–184.
- 101. Hulmi JJ, Lockwood CM & Stout JR (2010) Effect of protein/essential amino acids and resistance training on skeletal muscle hypertrophy: a case for whey protein. *Nutr Metab* **7**, 51.
- 102. Volpi E, Kobayashi H, Sheffield-Moore M *et al.* (2003) Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr* **78**, 250–258.
- 103. Tipton KD, Gurkin BE, Matin S et al. (1999) Nonessential amino acids are not necessary to stimulate net muscle protein synthesis in healthy volunteers. J Nutr Biochem 10, 89–95.
- 104. Cuthbertson D, Smith K, Babraj J *et al.* (2005) Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. *FASEB J* **19**, 422–424.
- Walker DK, Dickinson JM, Timmerman KL et al. (2011) Exercise, amino acids, and aging in the control of human muscle protein synthesis. *Med Sci Sports Exerc* 43, 2249– 2258.
- 106. Wilkinson DJ, Bukhari SSI, Phillips BE et al. (2018) Effects of leucine-enriched essential amino acid and whey protein bolus dosing upon skeletal muscle protein synthesis at rest and after exercise in older women. Clin Nutr 37, 2011–2021.
- 107. Beaudart C, Rabenda V, Simmons M et al. (2018) Effects of protein, essential amino acids, B-hydroxy B-methylbutyrate, creatine, dehydroepiandrosterone and fatty acid supplementation on muscle mass, muscle strength and physical performance in older people aged 60 years and over. A systematic review of the literature. J Nutr Health Aging 22, 117–130.
- Scognamiglio R, Piccolotto R, Negut C *et al.* (2005) Oral amino acids in elderly subjects: effect on myocardial function and walking capacity. *Gerontology* 51, 302–308.
- 109. Kim HK, Suzuki T, Saito K *et al.* (2012) Effects of exercise and amino acid supplementation on body composition and physical function in community-dwelling elderly Japanese sarcopenic women: a randomized controlled trial. *J Am Geriatr Soc* **60**, 16–23.
- 110. Verhoeven S, Vanschoonbeek K, Verdijk LB et al. (2009) Long-term leucine supplementation does not increase muscle mass or strength in healthy elderly men. Am J Clin Nutr 89, 1468–1475.
- 111. Leenders M, Verdijk LB, van der Hoeven L et al. (2011) Prolonged leucine supplementation does not augment muscle mass or affect glycemic control in elderly type 2 diabetic men. J Nutr 141, 1070–1076.
- 112. Suryawan A, Orellana RA, Fiorotto ML *et al.* (2011) Triennial growth symposium: leucine acts as a nutrient signal to stimulate protein synthesis in neonatal pigs. *J Anim Sci* **89**, 2004–2016.
- 113. Solerte SB, Gazzaruso C, Bonacasa R *et al.* (2008) Nutritional supplements with oral amino acid mixtures increases whole-body lean mass and insulin sensitivity in elderly subjects with sarcopenia. *Am J Cardiol* **101**, 69E–77E.

- 114. Zhou X, Xing B, He G *et al.* (2018) The effects of electrical acupuncture and essential amino acid supplementation on sarcopenic obesity in male older adults: a randomized control study. *Obes Facts* **11**, 327–334.
- Borsheim E, Bui QU, Tissier S et al. (2008) Effect of amino acid supplementation on muscle mass, strength and physical function in elderly. *Clin Nutr* 27, 189–195.
 Markofski MM, Jennings K, Timmerman KL et al.
- 116. Markofski MM, Jennings K, Timmerman KL et al. (2018) Effect of aerobic exercise training and essential amino acid supplementation for 24 weeks on physical function, body composition, and muscle metabolism in healthy, independent older adults: a randomized clinical trial. J Gerontol A 74, 1598–1604.
- 117. Luiking YC, Deutz NE, Memelink RG *et al.* (2014) Postprandial muscle protein synthesis is higher after a high whey protein, leucine-enriched supplement than after a dairy-like product in healthy older people: a randomized controlled trial. *Nutr J* **13**, 9.
- 118. Burd NA, Wall BT & van Loon LJ (2012) The curious case of anabolic resistance: old wives' tales or new fables? *J Appl Physiol* **112**, 1233–1235.
- 119. Kirk B, Mooney K, Amirabdollahian F *et al.* (2019) Exercise and dietary-protein as a countermeasure to skeletal muscle weakness: Liverpool hope universitysarcopenia aging trial (LHU-SAT). *Front Physiol* **10**, 1–11.
- 120. Gwin JA, Church DD, Wolfe RR *et al.* (2020) Muscle protein synthesis and whole-body protein turnover responses to ingesting essential amino acids, intact protein, and protein-containing mixed meals with considerations for energy deficit. *Nutrients* **12**, 1–15.
- 121. Sepulveda PV, Bush ED, Baar KJC et al. (2015) Pharmacology of manipulating lean body mass. Clin Exp Phramacol Physiol **42**, 1–13.
- 122. Wolfe RR (2002) Regulation of muscle protein by amino acids. J Nutr 132, 3219S–3224S.
- 123. Dickinson JM, Fry CS, Drummond MJ et al. (2011) Mammalian target of rapamycin complex 1 activation is required for the stimulation of human skeletal muscle protein synthesis by essential amino acids. J Nutr 141, 856– 862.
- 124. Francaux M, Demeulder B, Naslain D *et al.* (2016) Aging reduces the activation of the mTORC1 pathway after resistance exercise and protein intake in human skeletal

muscle: potential role of REDD1 and impaired anabolic sensitivity. *Nutrients* 8, 47.

- 125. Farnfield MM, Breen L, Carey KA *et al.* (2012) Activation of mTOR signalling in young and old human skeletal muscle in response to combined resistance exercise and whey protein ingestion. *Appl Physiol Nutr Metab* **37**, 21–30.
- 126. Pennings B, Koopman R, Beelen M et al. (2011) Exercising before protein intake allows for greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in both young and elderly men. Am J Clin Nutr 93, 322–331.
- 127. Symons TB, Sheffield-Moore M, Mamerow MM et al. (2011) The anabolic response to resistance exercise and a protein-rich meal is not diminished by age. J Nutr Health Aging 15, 376–381.
- 128. Fry CS & Rasmussen BB (2011) Skeletal muscle protein balance and metabolism in the elderly. *Curr Aging Sci* **4**, 260–268.
- 129. Shad BJ, Thompson JL & Breen L (2016) Does the muscle protein synthetic response to exercise and amino acidbased nutrition diminish with advancing age? A systematic review. Am J Physiol Endocrinol Metab 311, E803–E817.
- 130. Moro T, Brightwell CR, Deer RR *et al.* (2018) Muscle protein anabolic resistance to essential amino acids does not occur in healthy older adults before or after resistance exercise training. *J Nutr* **148**, 900–909.
- 131. Marshall RN, Smeuninx B, Morgan PT *et al.* (2020) Nutritional strategies to offset disuse-induced skeletal muscle atrophy and anabolic resistance in older adults: from whole-foods to isolated ingredients. *Nutrients* **12**, 1533.
- 132. Gade J, Pedersen RJ & Beck AM (2018) Effect of protein or essential amino acid supplementation during prolonged resistance exercise training in older adults on body composition, muscle strength, and physical performance parameters: a systematic review. *Rehabil Process Outcome* 7, 1179572718765760.
- 133. Invernizzi M, de Sire A, D'Andrea F, *et al.* (2019) Effects of essential amino acid supplementation and rehabilitation on functioning in hip fracture patients: a pilot randomized controlled trial. *Aging Clin Exp Res* **31**, 1517–1524.