



LEEDS
BECKETT
UNIVERSITY

Citation:

Sawicki, AJ and Żemojtel-Piotrowska, M and Balcerowska, JM and Sawicka, MJ and Piotrowski, J and Sedikides, C and Jonason, PK and Maltby, J and Adamovic, M and Agada, AMD and Ahmed, O and Al-Shawaf, L and Appiah, SCY and Ardi, R and Babakr, ZH and Bălăţescu, S and Bonato, M and Cowden, RG and Chobthamkit, P and De Pretto, L and Gouveia, VV and Haretche, C and Iliško, D and Aruta, JJB and Jia, F and Jovanović, V and Jukić, T and Kamble, SV and Khachatryan, N and Klicperova-Baker, M and Koralov, M and Kovacs, M and Kretchner, M and Fernandez, AL and Liik, K and Malik, NI and Malysheva, K and Moon, C and Muehlbacher, S and Nartova-Bochaver, S and Torres-Marín, J and Özsoy, E and Park, J and Piccinelli, E and Ramos-Diaz, J and Ridić, O and Samekin, A and Starc, A and Kiêu, TTT and Tomsik, R and Umeh, CS and Wills-Herrera, E and Wlodarczyk, A and Vally, Z and Zand, S (2022) The fear of COVID-19 scale: Its structure and measurement invariance across 48 countries. *Psychological Assessment*. ISSN 1040-3590 DOI: <https://doi.org/10.1037/pas0001102>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/8376/>

Document Version:

Article (Accepted Version)

This article was originally published in *Psychological Assessment* and is copyright of the American Psychological Association (APA).

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

The Fear of COVID-19 Scale:
Its Structure and Measurement Invariance Across 48 Countries

Artur J. Sawicki, University of Gdansk, Poland

Magdalena Żemojtel-Piotrowska, Cardinal Stefan Wyszyński University in Warsaw, Poland

Julia M. Balcerowska, University of Gdansk, Poland

Monika J. Sawicka, University of Warmia and Mazury, Poland

Jarosław Piotrowski, Cardinal Stefan Wyszyński University in Warsaw, Poland

Constantine Sedikides, University of Southampton, United Kingdom

Peter K. Jonason, University of Padua, Italy; Cardinal Stefan Wyszyński University in Warsaw,
Poland

John Maltby, University of Leicester, United Kingdom

Mladen Adamovic, Auckland Business School, University of Auckland, New Zealand; Monash
Business School, Monash University, Australia

Attiso M.G. Agada, Clinique Biasa, Togo

Oli Ahmed, University of Chittagong, Bangladesh

Laith Al-Shawaf, Department of Psychology, University of Colorado – Colorado Springs, USA

Seth Christopher Yaw Appiah, Kwame Nkrumah University of Science and Technology, Ghana

Rahkman Ardi, Universitas Airlangga, Indonesia

Zana Hasan Babakr, Soran University, Iraq

Sergiu Bălăţescu, University of Oradea, Romania

Mario Bonato, University of Padova, Italy

Richard G. Cowden, Harvard University, United States

Phatthanakit Chobthamkit, Thammasat University, Thailand

Laura De Pretto, Leeds Trinity University, United Kingdom

Valdiney V. Gouveia, Federal University of Paraiba, Brazil

Carmen Haretche, Universidad de la República, Uruguay

Dzintra Iliško, Daugavpils University, Latvia

John Jamir Benzon Aruta, De La Salle University, Philippines
Fanli Jia, Seton Hall University, USA
Veljko Jovanović, University of Novi Sad, Serbia
Tomislav Jukić, Josip Juraj Strossmayer University, Croatia
Shanmukh V. Kamble, Karnatak University, India
Narine Khachatryan, Yerevan State University, Armenia
Martina Klicperova-Baker, Czech Academy of Sciences, Czech Republic
Metodi Koralov, New Bulgarian University, Bulgaria
Monika Kovacs, ELTE Eötvös Loránd University, Hungary
Mabelle Kretchner, Reichman University, Israel
Aitor Larzabal Fernandez, Pontificia Universidad Católica del Ecuador-Ambato, Ecuador
Kadi Liik, Tallinn University, Estonia
Najma Iqbal Malik, University of Sargodha, Pakistan
Karine Malysheva, Taras Shevchenko National University of Kyiv, Ukraine
Chanki Moon, Leeds Beckett University, United Kingdom
Stephan Muehlbacher, Karl Landsteiner University of Health Sciences, Austria
Sofya Nartova-Bochaver, HSE University, Russia
Jorge Torres-Marín, University of Granada, Spain
Emrah Özsoy, Sakarya University, Turkey
Joonha Park, NUCB Business School, Japan
Elena Piccinelli, Instituto Universitário de Lisboa (ISCTE-IUL), CIS-IUL, Portugal
Jano Ramos-Diaz, Universidad de Ciencias y Humanidades, Peru
Ognjen Ridić, International University of Sarajevo, Bosnia & Herzegovina
Adil Samekin, M. Narikbayev KAZGUU University, Nur-Sultan, Kazakhstan
Andrej Starc, University of Ljubljana, Slovenia
Kiều Thị Thanh Trà, Ho Chi Minh City University of Education, Vietnam
Robert Tomsik, Research Institute for Child Psychology and Pathopsychology, Slovakia
Charles S. Umeh, College of Medicine, University of Lagos, Nigeria
Eduardo Wills-Herrera, Los Andes University, Colombia

Anna Włodarczyk, Universidad Catolica del Norte, Chile

Zahir Vally, United Arab Emirates University, United Arab Emirates

Somayeh Zand, Islamic Azad University, Rasht Branch, Rasht, Iran

The work of Artur Sawicki, Magdalena Żemojtel-Piotrowska, and Jarosław Piotrowski was supported by grant number 2017/26/E/HS6/00282 from the National Science Centre, Poland. Peter Jonason's work was partially funded by the Polish National Agency for Academic Exchange (PPN/U LM/2019/1/00019/U/00001) and a grant from the National Science Centre of Poland (2019/35/B/HS6/00682). The authors declare that there was no direct collaboration between researchers from conflicting countries in accord with many national laws.

Data, study materials, and all analyses codes are available at <https://osf.io/hxmzb>

The study was a part of a preregistered project, which details are available at <https://osf.io/hpwbj>

Contribution of each co-author in is pointed out in the brackets: Artur Sawicki

(Conceptualization: Lead; Data curation: Lead; Formal analysis: Lead; Methodology: Supporting;

Visualization: Lead; Writing – original draft: Lead; Writing – review & editing: Lead),

Magdalena Żemojtel-Piotrowska (Conceptualization: Lead; Funding acquisition: Equal;

Methodology: Lead; Project administration: Lead; Supervision: Equal; Writing – original draft:

Lead; Writing – review & editing: Lead), Julia M. Balcerowska (Conceptualization: Supporting;

Writing – original draft: Equal; Writing – review & editing: Supporting), Monika J. Sawicka

(Writing – original draft: Equal), Jarosław Piotrowski (Methodology: Lead; Project

administration: Lead), Constantine Sedikides (Supervision: Lead; Writing – original draft: Lead;

Writing –review & editing: Supporting), Peter K. Jonason (Writing – review & editing:

Supporting), John Maltby (Investigation: Equal), Mladen Adamovic (Investigation: Equal),

Attiso M.G. Agada (Investigation: Equal), Oli Ahmed (Investigation: Equal; Writing – review &

editing: Supporting), Laith Al Shawaf (Investigation: Equal; Writing – review & editing:

Supporting), Seth Christopher Yaw Appiah (Investigation: Equal), Rahkman Ardi (Investigation: Equal), Zana Hasan Babakr (Investigation: Equal), Sergiu Bălătescu (Investigation: Equal), Mario Bonato (Investigation: Equal), Richard G. Cowden (Funding acquisition: Supporting; Investigation: Equal; Writing – review & editing: Supporting), Phatthanakit Chobthamkit (Investigation: Equal), Laura De Pretto (Investigation: Equal), Valdiney V. Gouveia (Investigation: Equal), Carmen Haretche (Investigation: Equal), Dzintra Iliško (Investigation: Equal), John Jamir Benzon Aruta (Investigation: Equal), Fanli Jia (Investigation: Equal; Writing – review & editing: Supporting), Veljko Jovanović (Investigation: Equal; Writing – review & editing: Supporting), Tomislav Jukić (Investigation: Equal), Shanmukh V. Kamble (Investigation: Equal), Narine Khachatryan (Investigation: Equal), Martina Klicperova-Baker (Investigation: Equal), Metodi Koralov (Investigation: Equal), Monika Kovacs (Investigation: Equal), Mabelle Kretchner (Investigation: Equal), Aitor Larzabal Fernandez (Investigation: Equal), Kadi Liik (Investigation: Equal), Najma Iqbal Malik (Investigation: Equal), Karine Malysheva (Investigation: Equal), Chanki Moon (Funding acquisition: Supporting; Investigation: Equal), Stephan Muehlbacher (Investigation: Equal), Sofya Nartova-Bochaver (Investigation: Equal), Jorge Torres-Marín (Investigation: Equal), Emrah Özsoy (Investigation: Equal), Joonha Park (Investigation: Equal), Elena Piccinelli (Investigation: Equal; Writing – review & editing: Supporting), Jano Ramos-Diaz (Investigation: Equal), Ognjen Ridić (Investigation: Equal), Adil Samekin (Investigation: Equal), Andrej Starc (Investigation: Equal), Trà Thị Thanh Kiêu (Investigation: Equal), Robert Tomsik (Investigation: Equal), Charles S. Umeh (Investigation: Equal), Eduardo Wills-Herrera (Investigation: Equal), Anna Włodarczyk (Investigation: Equal), Zahir Vally (Investigation: Equal), Somayeh Zand (Investigation: Equal).

Corresponding author: Artur Sawicki, Institute of Psychology, University of Gdańsk, Bażyńskiego 4 Street, 80-309 Gdańsk, Poland, Tel.: +48 58 523 43 24; artur.sawicki@ug.edu.pl

Abstract

COVID-19 has been a source of fear around the world. We asked whether the measurement of this fear is trustworthy and comparable across countries. In particular, we explored the measurement invariance and cross-cultural replicability of the widely-used Fear of COVID-19 scale (FCV-19S), testing community samples from 48 countries ($N = 14,558$). The findings indicate that the FCV-19S has a somewhat problematic structure, yet the one-factor solution is replicable across cultural contexts and could be used in studies that compare people who vary on gender and educational level. The validity of the scale is supported by a consistent pattern of positive correlations with perceived stress and general anxiety. However, given the unclear structure of the FCV-19S, we recommend using latent factor scores, instead of raw scores, especially in cross-cultural comparisons.

Keywords: fear of COVID; measurement invariance; cross-cultural studies; coronavirus

Public Significance Statement: The study suggests that the widely-used Fear of COVID-19 scale (FCV-19S) could be used in cross-cultural research. However, given the scale's unclear structure, we recommend implementing the latent factor approach instead of relying on raw scores.

The Fear of COVID-19 Scale:

Its Structure and Measurement Invariance Across 48 Countries

Confirmed cases and deaths linked to the COVID-19 pandemic continue to occur daily worldwide (World Health Organization, 2021). The pandemic has also afflicted the public health systems and global economy. More relevant to this article, the pandemic has impacted mental health (Pfefferbaum & North, 2020; Torales et al., 2020), including increases in psychological distress (Xiong et al., 2020), loneliness (Zhou et al., 2021), depression, and anxiety (Fofana et al., 2020). The pandemic remains a source of unpredictability and threat (Counted et al., 2020; Fofana et al., 2020; Govender et al., 2020) that are likely to culminate in fear (Lazarus, 2006). Critically, the pandemic constitutes a direct source of fear (Enea et al., 2021; Fitzpatrick et al., 2020; Fofana et al., 2020), which negatively influences functioning in occupational (Fu et al., 2021; Hu et al., 2020) and family (Trougakos et al., 2020) life domains. As such, a measure of fear of COVID-19 can facilitate both basic and interventional research. We aimed to evaluate the psychometric properties of the most popular relevant instrument, the Fear of Covid-19 Scale (FCV-19S; Ahorsu et al., 2020). The pertinent article has been cited over 1700 times (Google Scholar; November 2, 2021). We assessed the FCV-19S's measurement invariance across 48 countries and tested its validity by examining its relations with gender, educational level, anxiety, and stress. The cross-cultural applicability of the scale has both empirical and practical implications. The scale is useful in identifying predictors and consequences of fear of COVID-19 globally and in shaping national and international policies to prevent the deleterious psychological effects of the pandemic.

Measuring Fear of COVID-19 Around the World

The FCV-19S (Ahorsu et al., 2020) was developed in response to the need to investigate the implications of the pandemic for physical and mental health. The item pool was chosen from 30 scales measuring fear (e.g., of dental examinations, chronic pain, or cancer) in diverse samples. This pool was then reduced by a panel of experts and researchers. The pool was further reduced based on psychometric analyses, resulting in a 7-item scale. We list these items in the original order (Ahorsu et al.), and we refer to them in the same order in our article: 1. "I am most afraid of coronavirus-19," "2. It makes me uncomfortable to think about coronavirus-19," "3. My

hands become clammy when I think about coronavirus-19,” “4. I am afraid of losing my life because of coronavirus-19, “5. When watching news and stories about coronavirus-19 on social media, I become nervous or anxious,” “6. I cannot sleep because I’m worrying about getting coronavirus-19, “7. My heart races or palpitates when I think about getting coronavirus-19” (see also Table 7). The FCV-19S, developed in Iranian samples, has a unifactorial structure with good internal consistency and validity, as demonstrated by its positive relationships with anxiety, depression, perceived infectability, and germ aversion (Ahorsu et al.).

The scale has been used widely across cultures. In relation to its psychometric properties, the FCV-19S has been validated chiefly through exploratory factor analyses (EFA) and confirmatory factor analyses (CFA). Most of these analyses produced a one-factor solution, although some yielded correlations of error variances between particular items (Alyami et al., 2020; Cavaleiro & Sticca, 2020; Mailliez et al., 2020; Wakashima et al., 2020). The rest of the analyses, though, produced a two-factor solution, with one factor reflecting psychological (items 1, 2, 4, 5) and another physiological (items 3, 6, 7) aspects of fear of COVID-19 (Barrios et al., 2021; Bitan et al., 2020; Iversen et al., 2021; Midorikawa et al., 2021; Reznik et al., 2020). Further, analyses in a few countries (i.e., Iran, Japan, Peru) suggested a modified adoption of the bi-factor model involving a general factor and two specific factors (i.e., physiological and psychological; Caycho-Rodríguez, Tomás et al., 2021; Huarcaya-Victoria et al., 2020; Masuyama et al., 2020).

In summary, the factorial structure of FCV-19S is unclear. Although most analyses pointed to a unifactorial solution, other analyses indicated that the corresponding model fitted the data poorly, requiring additional modifications such as correlating the error variances of the items or proposing two independent factors. Therefore, the validity of the FCV-19S across cultures is yet to be established.

Measurement Invariance

Given the popularity of the FCV-19S in many countries, we zeroed in on the comparability of results across cultural contexts both in terms of level of fear of COVID-19 and the scale’s predictors, correlates, and consequences. Measurement invariance allows a researcher to examine the extent to which scale-relevant results obtained in different cultures or groups (e.g.,

gender, educational level, religious denomination) are due to actual differences between cultures or groups (Cieciuch et al., 2019; Milfont & Fischer, 2010). Three levels of measurement invariance are typically assessed in cross-cultural studies: configural, metric, and scalar. Configural measurement invariance informs whether the scale factor structure (i.e., number of factors) is the same across compared groups. Metric invariance informs whether factor loadings are similar across groups, and allows for comparisons of correlates and regression weights across groups. Scalar measurement invariance informs whether the residuals are equivalent across compared groups, permitting comparisons of the latent factor scores. Multi-group confirmatory factor analysis provides evidence of whether parameters in the model are equal. Thus, the measurement invariance approach might be considered demanding, especially if a large number of groups is analyzed and scalar invariance is needed to test the hypotheses (Cieciuch et al., 2019). The alignment procedure, based on the Bayesian framework, addresses this problem (Muthen & Asparouhov, 2013). The procedure tests approximate measurement invariance, which extracts means as trustworthy as possible (and to some extent invariant), letting researchers to compare group levels of the studied phenomena.

Measurement invariance, albeit crucial for reliable comparisons between groups, has been under-investigated in the context of the FCV-19S. We were able to locate only a handful of studies examining the measurement invariance of the FCV-19S. One study on the scale's cross-cultural measurement invariance among Latin American countries found that the general population in Argentina, Colombia, Ecuador, El Salvador, Mexico, Paraguay, and Uruguay conceptualizes fear of COVID-19 in a bi-factorial structure (Caycho-Rodríguez, Valencia et al., 2021). However, this study identified only partial scalar invariance, indicating that one item (4. "I am afraid of losing my life because of coronavirus-19") is not invariant. One cross-cultural study did not report the scale's factorial structure and psychometric properties (Ali et al., 2021), and another illustrated invariance between the Dominican Republic and Spain (Piqueras et al., 2020). Lastly, a recent study (Lin et al., 2021) across 11 countries (i.e., Bangladesh, Brazil, Cuba, France, Iran, Italy, Japan, New Zealand, Pakistan, Taiwan, United Kingdom), primarily based on datasets from previous studies on the psychometric properties of the FCV-19S, showed that a one-factor solution of the FCV-19S indicated partial scalar invariance across countries and scalar

invariance across gender and age groups (i.e., child, young to middle-aged adult, older people). Other studies also demonstrated invariance between genders (Piqueras et al., 2020; Sakib et al., 2020) and healthcare workers versus non-healthcare workers (Huarcaya-Victoria et al., 2020). Further, two studies showed invariance between age groups (younger vs. older, with the specific age threshold varying across studies; Huarcaya-Victoria et al., 2020; Piqueras et al., 2020; Sakib et al., 2020). Taken together, although former studies have been concerned with FCV-19S's equivalency across a limited number of cultures (and some groups), no study has examined the scale's equivalency globally via a standardized procedure of data collection in culturally diversified context (i.e., most world regions).

Validity Criteria

Given the enormous popularity of the FCV-19S, some evidence has been accumulated pertaining to the relationship between fear of COVID-19 and demographic variables, like gender and education, and the relationship between fear of COVID-19 and general anxiety or stress. Fear of COVID-19 is higher among women (Ahmed et al., 2020; Gao et al., 2020; Lei et al., 2020), with a meta-analysis indicating that women are generally more responsive to negative emotional stimuli (Stevens & Hamann, 2012). Also, fear of COVID-19 is higher among individuals with lower education (Gao et al., 2020; Mazza et al., 2020; Olagoke et al., 2020; Wang et al., 2020), as these individuals have reduced access to health care (Oliver & Massialos, 2004) and are thus more likely to suffer severe consequences of COVID-19 (i.e., hospitalization; Niedzwiedz et al., 2020). Fear is an adaptive response to threat (Ramikie & Ressler, 2018), usually followed by avoidance. If the threat is difficult to avoid, as is the case with the pandemic, it may cascade into general anxiety and stress (Gross & Jazaieri, 2014). Indeed, the literature points to positive relationships between fear of COVID-19 on the one hand, and general anxiety as well as stress on the other (Ahorsu et al., 2020; Bitan et al., 2020; Mertens et al., 2021; Satici et al., 2020; Tsipropoulou et al., 2020). It is important, though, to examine the magnitude of overlap between measures of the FCV-19S and anxiety, as well as measures of the FCV-19S and stress. High correlations would indicate a considerable overlap between measured constructs (Vatcheva et al., 2016). As such, we considered both anxiety and stress as external criteria of fear of COVID-19.

Current Study

We examined the factorial structure and measurement invariance of the FCV-19S across 48 cultures, and across gender and education. To this end, we analyzed data originating from a single project, with a standardized way of data collection, using identical online surveys across all countries. We expected to find metric levels of measurement invariance, allowing for a reliable comparison of the scale's predictors and correlates. We also examined the scale's validity, expecting higher levels of fear of COVID-19 among women and less educated individuals, and moderate positive correlations with anxiety and stress. We measured anxiety and stress with well-established and widely used—including cross-culturally—scales (Leung et al., 2010; Marteau et al., 1992; Vallejo et al., 2018). These were the State-Trait Anxiety Inventory 6 (STAI-6; Tluczek et al., 2009) and the Perceived Stress Scale 4 (PSS-4; Cohen et al., 1983).

Method

Participants

We collected the data online between 24th April and 20th November 2020 as a part of the international project “COVID-19, Personality and Quality of Life: Self-Enhancement in the Time of Pandemic”, spanning 60 countries.¹ Participants were invited to engage in the study via email or an announcement on Facebook forums devoted to COVID-related topics, which included a link to the project's website. There, participants reported their nationality and country of residence, and selected their preferred language version out of 35 languages². We did not offer remuneration, except for the Republic of South Africa and the United Kingdom (2GBP, or approximately 2.5USD, per participant in each case)³. We set a minimum $N = 100$ for each country (Supplementary Material, Table S7). Samples from 48 countries reached that threshold (Table 1). We included only participants over the age of 18 years who answered all measures⁴. The sample comprised 14,558 participants (65.64% women, 33.76% men, 0.60% “other” or

¹ See: <https://osf.io/hpwbj> (preregistered OSF project).

² Most participants in each country ($M = 96.17\%$, $SD = 6.46\%$) selected the country's official language (e.g., Italians selected Italian). Latvians were the exception: Only 64.90% of them selected Latvian, with most of the rest (> 30%) selecting Russian.

³ We decided on no participant remuneration due to lack of funding. However, in these two countries, we encountered insurmountable difficulties with data collection, probably due to many competing survey demands. We managed to carry out the surveys after the local collaborators secured funding from their home institutions.

⁴ We presented measures in a separate random order for each participant. We checked data quality via three (randomly displayed across the survey) attention-checks (e.g., “This question is for checking your attention. Please mark number 2”). Wrong responses to any of the three items resulted in exclusion.

unreported), aged between 18 and 98 years ($M = 31.70$, $SD = 12.34$). Of participants, 1.15% had a Primary Education degree, 4.32% a Lower Secondary Education degree, 28.75% an Upper Secondary Education degree, (37.52%) a Bachelor's (or equivalent) degree, 22.94% a Master's [or equivalent] degree, and 5.33% a PhD degree. Lastly, participants' responses to a socioeconomic status question ("How would you describe the economic status of your family on a scale from 1 to 7?"; 1 = *much lower than average*, 4 = *average*, 7 = *much higher than average*) hovered around the scale mean ($M = 4.26$, $SD = 1.67$).

Procedure and Measures

Scales were presented in a separate random order for each participant. Where a translation to a local language was unavailable, local team members translated it using a back-translation procedure (Brislin, 1970). In the present study, we only analyzed the data about fear of COVID-19, anxiety, and stress. Fear of COVID-19 was measured with the FCV-19S (1 = *strongly disagree*, 7 = *strongly agree*). Anxiety was measured with the 6-item STAI-6 (Tluczek et al., 2009). Participants indicated how frequently (0 = *never*, 4 = *very often*) in the last month they experienced each anxiety-relevant state (e.g., "I felt tense," "I felt worried"). Stress was measured with the 4-item PSS-4 (Cohen et al., 1983). Participants indicated how frequently (0 = *never*, 4 = *very often*) in the last month they experienced each relevant state (e.g., "... unable to control the important things in your life?", "... felt that things were going your way?" [reverse-scored]). We report descriptive statistics and estimates of internal consistency for all scales in Table 6. The project was reviewed and approved by the Ethics Committee of Cardinal Stefan Wyszyński University in Warsaw.

Data-Analytic Strategy

We began by conducting a CFA of the FCV-19S across all (48) studied countries. We tested several factor structure variations of the FCV-19S, with a single-factor structure as the starting point. We followed up with exploratory testing of the two-factor model and the modified single-factor model in which we estimated freely (6) error covariances. In the CFA, where country-level samples had a poor fit, we used modification indices to identify areas where model fit could be improved by relaxing parameter constraints iteratively and retesting the model after each modification. Subsequently, in all groups with an acceptable model fit, we carried out multi-

group confirmatory factor analysis (MGCFA) to test for configural, metric, as well as scalar invariance among countries, between men and women, and among educational levels. In the MGCFA, we tested for metric invariance by relaxing factor loading constraints in non-invariant countries, one at a time, based on modification indices. We followed the MGCFA with the alignment procedure to identify the most non-invariant parameters (Cieciuch et al., 2019). Subsequently, we repeated this procedure for anxiety and stress in countries in which we found metric or partial metric invariance of the FCV-19S.⁵ Finally, in all countries, we examined the correlations among latent (invariant) factors of the three scales measuring fear of Covid-19, anxiety, and stress.

In all analyses, we used IBM SPSS 26 and Mplus 7.2 (Muthen & Muthen, 1998-2014). In factor analyses, we used the Robust Maximum Likelihood (MLR) estimator due to: (1) relatively small sample sizes, (2) 7-category response scale (Rhemtulla et al., 2012), and (3) potential deviations from normality (Yuan & Bentler, 2000). Examining model fit, we relied on the most common fit indices: Chi-squared, CFI, RMSEA, and SRMR, with the following thresholds for acceptable fit: $CFI > .95$, $RMSEA < .10$, $SRMR < .08$ (Brown, 2015; Hu & Bentler, 1999). We implemented a more liberal RMSEA threshold due to its inflated rejection rates of models with small *df* (Kenny et al., 2015), which is congruent with MacCallum et al.'s (1996) suggestion that RMSEA between .08 and .10 indicates a mediocre fit. In the MGCFA comparing gender and educational levels, we relied on the following thresholds for a meaningful difference between the models: $\Delta CFI = -.002$ (Meade et al., 2008), $\Delta RMSEA = .015$ (Chen, 2007). However, given the large number of compared groups, we relied in the cross-cultural MGCFA on more liberal thresholds to test metric invariance ($\Delta CFI = -.02$, $\Delta RMSEA = .03$; Rutkowski & Svetina, 2014). To test scalar invariance, we implemented Meade et al.'s (2008) criteria. In the alignment optimization, we used 25% of non-invariant parameters as a threshold for trustworthy mean estimations (Cieciuch et al., 2019). In all analyses, we used MGCFA estimations wherever we found

⁵ Using anxiety and stress as criteria requires correlational analysis. Correlation coefficients are comparable between groups only when both correlated variables are invariant on the metric level. Therefore, we analyzed the structure of STAI-6 and PSS-4 only in countries (43 out of 48) in which we found metric invariance for the FCV-19S. We obtained partial metric invariance in 40 countries for STAI-6 and in 33 countries for PSS-4. We present results of the MGCFA analyses for STAI-6 and PSS-4 Supplemental Material. Detailed results of single-group CFAs are available at the OSF page: <https://osf.io/hxmzb>

invariance with this method. We used scores estimated via the alignment method only when we obtained scalar invariance with MGCFA. Data, study materials, and all analyses codes are available at <https://osf.io/hxmzb>.

Results

The FCV-19S Structure Across Countries

We present descriptive statistics for all FCV-19S items in Table 8, indicating that items 3, 6, and 7 were, to a varying degree, right-skewed. The initial single-factor model fitted the data poorly in all countries, except Russia, where it was acceptable (Table S1, Supplemental Material). Therefore, we modified models for each country, relaxing error covariances, based on modification indices⁶. The suggested modification indices repeatedly revealed several covariances which, estimated freely, improved the model fit (between errors of items 1 and 2; of items 1 and 4; of items 2 and 5 [psychological aspects of fear], and of items 3, 6, and 7 [physiological aspects of fear]). These results appear to suggest a two-factor structure, with one factor pertinent to psychological aspects of fear and the other factor pertinent to physiological aspects of fear (as in prior research; Iversen et al., 2021). We explored this potential two-factor structure (see Table S2, Supplemental Material for CFAs results), which showed improved global fit. However, we failed to replicate this solution in every country: in most countries, the same covariances as in the single-factor model needed to be relaxed to improve fit, and the two factors were highly correlated (average $\rho = .78$, $SD_{\rho} = .09$), which might result in multicollinearity issues in future studies, thus limiting the utility of the scale (Vatcheva et al., 2016).⁷ We considered this a cue that the scale is unifactorial but that there may be other (than fear of COVID-19) sources of intercorrelations between items. Although testing the invariance of different baseline models across groups is a valid approach, the purpose of this research was to arrive at a broad, cross-culturally comparable solution of FCV-19S. Therefore, to achieve a model applicable to all

⁶ Given that relaxing error covariances is regarded by some (Sellbom & Tellegen, 2019) as a somewhat controversial practice, we also applied the alternative strategy of dropping items that we considered redundant based on modification indices. We tested both single-factor and two-factor solutions, iteratively dropping items 1, 2, and 6. Out of the alternative solutions that we tested, a five-item, two-factor model (with items 4 and 5 measuring psychological aspects of fear and items 3, 6, and 7 physiological aspects of fear, respectively) fitted the best. Although it was well-fitted in 34 out of 48 countries, the two factors were highly correlated (average $\rho = .85$, $SD_{\rho} = .08$). The results are available on the project's OSF page: <https://osf.io/hxmzb>.

⁷ Noticeable exceptions were Iran ($r = .59$), Japan ($r = .59$), and United Arab Emirates ($r = .49$).

studied groups (Figure 1), we relaxed the abovementioned six covariances most commonly indicated by modification indices. Given this exploratory approach, we opted not to relax any more covariances⁸. This modified single-factor model (Table 2) was well-fitted to the data in 43 out of 48 countries (exceptions: Chile, Israel, Italy, Slovakia, Slovenia).⁹

Group-Level Measurement Invariance

After establishing an acceptable model fit (i.e., our modified single-factor model), we tested, using an MGCFA, whether it is cross-culturally invariant in 43 of the 48 countries (exceptions, as above, were: Chile, Israel, Italy, Slovakia, Slovenia). We obtained configural invariance between those countries. Moreover, we found partial metric invariance after relaxing several factor loadings (17 [5.65%]) and excluding Serbia, but we obtained no scalar invariance (Table 3). Thus, we followed MCFA with the alignment procedure in which we examined the non-invariance on two levels, looking for items that were non-invariant most often and for countries in which most items were non-invariant. The basic assumption of the alignment procedure is the configural invariance of the model; therefore, we tested it on 43 out of 48 countries again (including Serbia). It showed that, in total, 7.31% of factor loadings were non-invariant (mostly in Latvia [4], Serbia [4], and the Republic of South Africa [3]), supporting the partial metric invariance obtained in the MGCFA. Moreover, 24.92% of intercepts were non-invariant¹⁰ (mostly items 1 and 4—see Table 7), which shows that means can be estimated and compared with trustworthiness (keeping 25% of the non-invariant intercepts as the threshold of acceptability). We present latent mean levels of fear of COVID-19 across 43 countries in Table 6. We display in Figure 2 distributions of approximate scores estimated with the alignment procedure across countries.

In addition, we compared the fit of the three tested models in both genders and at different educational levels. First, we removed the 85 participants who reported “other” for gender. Again,

⁸ We present estimated correlation coefficients between listed error terms in Table S3, Supplemental Material.

⁹ In those countries, single-factor, two-factor, and modified single-factor models were not well-fitted to the data. Thus, we exploratorily searched for the best-fitted model in each country. A two-factor model, with several error covariances relaxed, fit best in Chile (covariances: 1+4, 3+6, 2+4, 2+7), Slovakia (1+2 and 2+5), and Slovenia (1+2, 2+5, 5+7). In Israel, a 1-factor with correlations (3+6+7 and 4+5) fit best. We found no acceptably fitted model in Italy.

¹⁰ In Bulgaria, Czech Republic, Kazakhstan, Poland, Portugal, and Turkey, 4 out of 7 intercepts were non-invariant. We suggest interpreting means for those countries with caution.

our 1-factor modified model was the best fit in both genders (Table 4). We found partial scalar invariance after relaxing items' 2, 3, and 5 intercepts among women. Second, we compared model fit across educational levels and found that the 1-factor modified model fitted the best, with slightly worse fit in “primary education” and “doctoral” groups. We found partial scalar invariance across educational levels, after relaxing intercepts of items 2 and 5 in the Bachelor's and Master's degree groups, items 2 and 4 in the doctoral group, and item 1 in the upper secondary degree group (Table 5).

Relations with External Criteria

Lastly, we used our modified measurement model to examine individual differences in fear of COVID-19. We examined gender differences (independent samples *t*-test, using partially scalar invariant latent factor scores of FCV-19S), educational level differences (one-way Analysis of Variance with Bonferroni correction, using partially scalar invariant latent factor scores of FCV-19S), and correlations (Pearson's *r*, using partially metric invariant latent factor scores of FCV-19S) with anxiety and stress. Women feared COVID-19 more than men, $t_{14467} = 11.92, p < .001$, Cohen's $d = 0.21$. Participants with the least education feared COVID-19 the most ($F_{5,14550} = 68.81, p < .001, \eta^2 = .023$), and this relation was linear (Supplementary Material, Figure S1). Also, fear of COVID-19 was positively related to anxiety and stress in most countries (except Japan and Uruguay), both in men and women and at different educational levels (Tables 4 and 5).

Discussion

Given the extensive, cross-cultural use of the FCV-19S (Ahorsu et al., 2020) in different populations, we set to find out if the results obtained are comparable across cultures, genders, and educational levels by inspecting the scale's measurement invariance. In particular, we tested FCV-19S's cross-cultural, cross-gender, and cross-educational replicability in 48 countries. We supplemented our analyses by using external criteria, and specifically anxiety and stress, as important correlates of fear of COVID-19 (Ahorsu et al., 2020; Satici et al., 2020; Tsipropoulou et al., 2020). Based on prior research, we expected that women (vs. men) and persons with lower (vs. higher) educational levels would experience more fear.

The FCV-19S Structure Across Countries, Genders, and Educational Levels

We did not replicate the unifactorial structure of the scale, proposed in the original study (Ahorsu et al., 2020), in any of our samples. The literature on the FCV-19S has reported both a one-factor and a two-factor solution. As such, we pitted a one-factor against a two-factor model. The one-factor uncorrelated solution did not fit the data well, as the items formed two groups: one related to psychological aspects (items 1, 2, 4, 5) and one to physiological aspects (items 3, 6, 7) of fear. Further analyses did not support the two-factor solution either, indicating non-optimal (Vatcheva et al., 2016) high correlation between the two factors. Thus, the final model that we propose is unifactorial, yet includes additional correlations among or between items (items 3, 6, 7; items 1, 2; items 1, 4; items 2, 5). This modified single-factor solution indicates that fear of the COVID-19, as measured by the FCV-19S, is a rather homogenous, or at least a general, phenomenon. Finally, the FCV-19S had good reliability (Tables 4-6).

Additional analyses offered insights into cross-cultural differences in the measurement of fear of COVID-19, revealing, in some countries, deviations from the single-factor modified model that we propose. This model did not fit the data well in Chile, Israel, Italy, Slovakia, and Slovenia. Different models fit best in these countries, with relatively similar 2-factor solutions fitting in Slovakia and Slovenia, but a different 2-factor solution fitting in Chile. In Israel, a 1-factor solution fitted best, although still not meeting the criteria for acceptable fit. Moreover, no factorial solution fit in Italy, despite exploring several variations of 1-factor and 2-factor models.

Comparability of FCV-19S Results Across Countries, Genders, and Educational Levels

The modified single-factor solution showed a considerable level of equivalence across countries (partial metric invariance— in 43 of 48 countries), genders (partial scalar invariance), and educational levels (scalar invariance). The single-factor modified solution, then, assures reliable comparisons among countries or groups. In particular, metric invariance allows for reliable comparisons of predictors and consequences of fear of COVID-19 across different countries. Although not scalar-invariant, the results of the alignment procedure suggest that comparable mean levels of fear of COVID-19 can be trustworthily estimated cross-culturally. Partial scalar invariance allows for comparing levels of fear across genders and educational groups.

Our findings partially replicated Lin et al.'s (2021) results, as we obtained partial scalar invariance for gender. However, unlike Lin et al., we obtained only metric levels of cross-country invariance using the MGCFA approach. We collected data from more countries than Lin et al., and the number of countries or group tested diminishes the likelihood of uncovering scalar invariance (Cieciuch et al., 2019). Yet, we were able to compare indirectly levels of latent scores across countries using the alignment approach. Cross-country ranking of fear of COVID-19 differed somewhat between the studies. We provided an illustration in reference to Bangladesh. In our study, level of fear of COVID-19 in Iran was in the middle of the country rankings and lower than in Bangladesh (Figure 2); however, in Lin et al., level of fear of COVID-19 was at the top of the country rankings and higher than in Bangladesh.

Lack of scalar invariance could stem from different sources, from translation issues to referring to different experiences (Oshio, 2010). The most non-invariant items were 1 (“I am most afraid of the coronavirus-19”) and 4 (“I am afraid of losing my life because of coronavirus-19”), which referred to psychological aspects of fear of COVID-19. Therefore, these two items were mostly responsible for the lack of scalar invariance, implying that participants from different countries answered these questions in an incomparable manner. Items referring to physiological aspects of fear were less problematic, which means that they were understood similarly in different countries. Moreover, in all samples, we observed distinct distributions of scoring on the two aspects of fear: scoring on items measuring psychological aspects of fear was normally distributed, whereas scoring on items measuring physiological aspects of fear was right-skewed.

Relationship with External Criteria

Finally, the study confirmed the validity of the FCV-19S. Replicating prior research (Sakib et al., 2020), women (vs. men) experienced more fear of COVID-19. We also found that less (vs. more) educated participants experienced increased fear of COVID-19. This may be due to less-educated participants having reduced access to health care and to customized (i.e., adapted to their medical knowledge) health promotion media messages (Oliver & Massialos, 2004), as well as their weaker tendency to respond to such messages (Iversen & Kraft, 2006) or engage in self-management in illnesses (Adams, 2010). Similarly, a positive relation of general anxiety and

stress with fear of COVID-19 is consistent with prior findings regarding fear of COVID-19 measured both with the FCV-19S (Ahorsu et al., 2020; Satici et al., 2020) and a single-item scale (Fitzpatrick et al., 2020). Notably, the strength of those relationships distinguishes fear of Covid-19 from general anxiety and stress: mean r was .32 for the relationship between fear of COVID-19 and anxiety, and .26 for the relationship between fear of COVID-19 and stress. These result patterns confirm the divergent validity of FCV-19S.

Limitations and Future Directions

Although we collected data in many countries, our samples were convenient rather than representative, and relatively small. Also, the gender balance was not optimal, as most participants were women. Further, most participants were well-educated, as less educated participants likely had reduced access to the internet. (The online character of our study was necessitated by the pandemic.) Sample characteristics were related to model fit (Supplementary Material, Table S8), showing that it was negatively related to age (i.e., better fit in younger samples) and gender (i.e., better fit in samples consisting of women). Examining the utility of the FCV-19S in specific populations (such as the elderly or persons with chronic illnesses) is a promising line of inquiry. As we did not control for response-style bias, we could not rule out that the results might have been, to some extent, distorted. Future research with this scale could adjust the scores accordingly (e.g., controlling for acquiescence bias with anchoring vignettes; He et al., 2017). Moreover, given that the pandemic did not spread evenly across the world, there were country differences in infection rates. Infection rates were on the rise in some countries (e.g., Brazil, Ecuador), but had stabilized in others (e.g., Austria, Slovenia), during our data collection period. Therefore, future validation work would also do well to compare the structure of the FCV-19S in the same country, but in different time periods.

Our results indicate that the FCV-19S items do not form one clear, uncorrelated factor, suggesting some heterogeneity in the measurement model. We obtained the proposed solution with a data-driven strategy of correlating errors based on modification indices, which might be somewhat controversial (Sellbom & Tellegen, 2019), possibly impairing its replicability. Given the exploratory character of our strategy, we interpreted those correlations (i.e., psychological and physiological aspects of fear) in a post-hoc manner. We obtained equivocal support for cross-

cultural comparisons in levels of fear of COVID-19, and this heterogeneity was further visible in invariance tests. Items measuring psychological aspects of fear of COVID-19 were non-invariant more often than those measuring the physiological aspects of such fear. Further, their distribution differed, indicating that physiological aspects of fear were less pronounced. Therefore, follow-up research could examine independently psychological from physiological aspects of fear of COVID-19 (Mertens et al., 2021). Regardless, more conclusive evidence about an underlying factor structure is only obtainable with longitudinal data, as cross-sectional data can obscure it (VanderWeele & Batty, 2020).

Concluding Remarks

How did the FCV-19S (Ahorsu et al., 2020) fare? We examined the scale in 48 countries and found that it is unifactorial. However, we also found that the scale is in need of measurement modifications, which currently limit its utility. Factor analyses indicated that the scale assesses two aspects of fear of COVID-19, psychological and physiological, but the issue needs more thorough investigation. Although the scale is suitable for cross-cultural research, it is limited when it comes to examining correlates and predictors of fear of COVID-19 (as indicated by metric levels of invariance). We suggest that the FCV-19S might be cautiously used in studies focused on cross-cultural comparisons of level of fear of COVID-19 (as per the findings on alignment optimization). However, given the number of adjustments needed in the model, we recommend the latent variable approach, relying on latent scores. Finally, we suggest that the scale be used on comparing level of fear of COVID-19 between genders and educational groups (as per the finding on scalar-level of invariance).

References

- Adams, R. J. (2010). Improving health outcomes with better patient understanding and education. *Risk Management and Healthcare Policy*, 3, 61-72. <https://doi.org/10.2147/RMHP.S7500>
- Ahmed, M. Z., Ahmed, O., Aibao, Z., Hanbin, S., Siyu, L., & Ahmad, A. (2020). Epidemic of COVID-19 in China and associated psychological problems. *Asian Journal of Psychiatry*, 51, 102092. <https://doi.org/10.1016/j.ajp.2020.102092>
- Ahorsu, D. K., Lin, C. Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2020). The fear of COVID-19 scale: Development and initial validation. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00270-8>
- Ali, M., Uddin, Z., Banik, P. C., Hegazy, F. A., Zaman, S., Ambia, A. S. M., ... & Ahsan, G. U. (2021). Knowledge, attitude, practice and fear of COVID-19: A cross-cultural study. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-021-00638-4>
- Alyami, M., Henning, M., Krägeloh, C. U., & Alyami, H. (2020). Psychometric evaluation of the Arabic version of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00316-x>
- Barrios, I., Ríos-González, C., O'Higgins, M., González, I., García, O., Díaz, N. R., ... & Torales, J. (2021). Psychometric properties of the Spanish version of the Fear of COVID-19 scale in Paraguayan population. *Irish Journal of Psychological Medicine*, 1-6. doi:10.1017/ipm.2021.5
- Bitan, D. T., Grossman-Giron, A., Bloch, Y., Mayer, Y., Shiffman, N., & Mendlovic, S. (2020). Fear of COVID-19 scale: Psychometric characteristics, reliability and validity in the Israeli population. *Psychiatry Research*, 289, 113100. <https://doi.org/10.1016/j.psychres.2020.113100>
- Brislin, R.W. (1970). Back-translation for cross-cultural research. *Journal of Cross-Cultural Psychology*, 1(3), 185-216. <https://doi.org/10.1177/135910457000100301>
- Brown, T. A. (2015). *Confirmatory factor analysis for applied research*. Guilford Press.
- Cavalheiro, F. R. S., & Sticca, M. G. (2020). Adaptation and validation of the Brazilian version of the fear of COVID-19 scale. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00415-9>
- Caycho-Rodríguez, T., Tomás, J. M., Barboza-Palomino, M., Ventura-León, J., Gallegos, M., Reyes-Bossio, M., & Vilca, L. W. (2021). Assessment of fear of COVID-19 in older adults: Validation

- of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00438-2>
- Caycho-Rodríguez, T., Valencia, P. D., Vilca, L. W., Cervigni, M., Gallegos, M., Martino, P., ... & Burgos Videla, C. (2021). Cross-cultural measurement invariance of the fear of COVID-19 scale in seven Latin American countries. *Death Studies*. Advance online publication. <https://doi.org/10.1080/07481187.2021.1879318>
- Chen, F. F. (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, 14(3), 464-504. <https://doi.org/10.1080/10705510701301834>
- Cieciuch, J., Davidov, E., Schmidt, P., & Algesheimer, R. (2019). How to obtain comparable measures for cross-national comparisons. *KZfSS Kölner Zeitschrift für Soziologie und Sozialpsychologie (The Cologne Journal of Sociology and Social Psychology)*, 71(1), 157-186. <https://doi.org/10.1007/s11577-019-00598-7>
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385-396. <https://doi.org/10.2307/2136404>
- Counted, V., Pargament, K. I., Bachera, A. O., Joynt, S., & Cowden, R. G. (2020). Hope and well-being in vulnerable contexts during the COVID-19 pandemic: Does religious coping matter? *Journal of Positive Psychology*. Advance online publication. <https://doi.org/10.1080/17439760.2020.1832247>
- Enea, V., Eisenbeck, N., Petrescu, T. C., & Carreno, D. F. (2021). Perceived impact of quarantine on loneliness, death obsession, and preoccupation with God: Predictors of increased fear of COVID-19. *Frontiers in Psychology*, 12, 643977. <https://doi.org/10.3389/fpsyg.2021.643977>
- Fitzpatrick, K. M., Harris, C., & Drawve, G. (2020). Fear of COVID-19 and the mental health consequences in America. *Psychological Trauma: Theory, Research, Practice, and Policy*, 12(S1), S17–S21. <https://doi.org/10.1037/tra0000924>
- Fofana, N. K., Latif, F., Sarfraz, S., Bashir, M. F., & Komal, B. (2020). Fear and agony of the pandemic leading to stress and mental illness: an emerging crisis in the novel coronavirus (COVID-19) outbreak. *Psychiatry Research*, 291, 113230. <https://doi.org/10.1016/j.psychres.2020.113230>

- Fu, S. (Q.), Greco, L. M., Lennard, A. C., & Dimotakis, N. (2021). Anxiety responses to the unfolding COVID-19 crisis: Patterns of change in the experience of prolonged exposure to stressors. *Journal of Applied Psychology, 106*(1), 48–61. <https://doi.org/10.1037/apl0000855>
- Gao, J., Zheng, P., Jia, Y., Chen, H., Mao, Y., Chen, S., ... & Dai, J. (2020). Mental health problems and social media exposure during COVID-19 outbreak. *PLoS One, 15*(4), e0231924. <https://doi.org/10.1371/journal.pone.0231924>
- Govender, K., Cowden, R. G., Nyamaruze, P., Armstrong, R. M., & Hatane, L. (2020). Beyond the disease: Contextualized implications of the COVID-19 pandemic for children and young people living in Eastern and Southern Africa. *Frontiers in Public Health, 8*(504), <https://doi.org/10.3389/fpubh.2020.00504>
- Gross, J. J., & Jazaieri, H. (2014). Emotion, emotion regulation, and psychopathology: An affective science perspective. *Clinical Psychological Science, 2*(4), 387-401. <https://doi.org/10.1177/2167702614536164>
- He, J., Van de Vijver, F. J., Fetvadjev, V. H., de Carmen Dominguez Espinosa, A., Adams, B., Alonso-Arbiol, I., ... & Möttus, R. (2017). On enhancing the cross-cultural comparability of Likert-scale personality and value measures: A comparison of common procedures. *European Journal of Personality, 31*(6), 642-657. <https://doi.org/10.1002/per.2132>
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural equation modeling: a multidisciplinary journal, 6*(1), 1-55. <https://doi.org/10.1080/10705519909540118>
- Huarcaya-Victoria, J., Villarreal-Zegarra, D., Podestà, A., & Luna-Cuadros, M. A. (2020). Psychometric properties of a Spanish version of the Fear of COVID-19 Scale in general population of Lima, Peru. *International Journal of Mental Health and Addiction*. Advance Online Publication. <https://doi.org/10.1007/s11469-020-00354-5>
- Iversen, A. C., & Kraft, P. (2006). Does socioeconomic status and health consciousness influence how women respond to health related messages in media? *Health Education Research, 21*(5), 601-610. <https://doi.org/10.1093/her/cyl014>
- Iversen, M. M., Norekvål, T. M., Oterhals, K., Fadnes, L. T., Mæland, S., Pakpour, A. H., & Breivik, K. (2021). Psychometric properties of the Norwegian version of the Fear of COVID-19

Scale. *International Journal of Mental Health and Addiction*. Advance online publication.

<https://doi.org/10.1007/s11469-020-00454-2>

Kenny, D. A., Kaniskan, B., & McCoach, D. B. (2015). The performance of RMSEA in models with small degrees of freedom. *Sociological methods & research*, 44(3), 486-507.

<https://doi.org/10.1177/0049124114543236>

Lazarus, R. S. (2006). *Stress and emotion: A new synthesis*. Springer Publishing Company

Lei, L., Huang, X., Zhang, S., Yang, J., Yang, L., & Xu, M. (2020). Comparison of prevalence and associated factors of anxiety and depression among people affected by versus people unaffected by quarantine during the COVID-19 epidemic in southwestern China. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*, 26: e924609-1–e924609-12. <https://doi.org/10.12659/MSM.924609>

Leung, D. Y., Lam, T. H., & Chan, S. S. (2010). Three versions of Perceived Stress Scale: validation in a sample of Chinese cardiac patients who smoke. *BMC Public Health*, 10(1), 1-7.

<https://doi.org/10.1186/1471-2458-10-513>

Lin, C.-Y., Hou, W. L., Mamun, M. A., Aparecido da Silva, J., Broche-Pérez, Y., Ullah, I., ... & Pakpour, A. H. (2021). Fear of COVID-19 Scale (FCV-19S) across countries: Measurement invariance issues. *Nursing Open*. <https://doi.org/10.1002/nop2.855>

Mailliez, M., Griffiths, M. D., & Carre, A. (2020). Validation of the French version of the Fear of COVID-19 Scale and its associations with depression, anxiety and differential emotions.

International Journal of Mental Health and Addiction, 1–15. Advance online publication.

<https://doi.org/10.1007/s11469-021-00499-x>

Marteau, T. M., & Bekker, H. (1992). The development of a six-item short-form of the state scale of the Spielberger State—Trait Anxiety Inventory (STAI). *British Journal of Clinical Psychology*,

31(3), 301-306. <https://doi.org/10.1111/j.2044-8260.1992.tb00997.x>

Masuyama, A., Shinkawa, H., & Kubo, T. (2020). Validation and psychometric properties of the Japanese version of the Fear of COVID-19 Scale among adolescents. *International Journal of*

Mental Health and Addiction. Advance online publication. [https://doi.org/10.1007/s11469-020-](https://doi.org/10.1007/s11469-020-00368-z)

[00368-z](https://doi.org/10.1007/s11469-020-00368-z)

- Mazza, C., Ricci, E., Biondi, S., Colasanti, M., Ferracuti, S., Napoli, C., & Roma, P. (2020). A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: Immediate psychological responses and associated factors. *International Journal of Environmental Research and Public Health*, *17*(9), 3165. <https://doi.org/10.3390/ijerph17093165>
- Milfont, T. L., & Fischer, R. (2010). Testing measurement invariance across groups: Applications in cross-cultural research. *International Journal of Psychological Research*, *3*(1), 111-130. <https://doi.org/10.21500/20112084.857>
- Meade, A. W., Johnson, E. C., & Braddy, P. W. (2008). Power and sensitivity of alternative fit indices in tests of measurement invariance. *Journal of Applied Psychology*, *93*(3), 568-592. <https://doi.org/10.1037/0021-9010.93.3.568>
- Mertens, G., Duijndam, S., Smeets, T., & Lodder, P. (2021). The latent and item structure of COVID-19 fear: A comparison of four COVID-19 fear questionnaires using SEM and network analyses. *Journal of Anxiety Disorders*, *81*, 102415. <https://doi.org/10.1016/j.janxdis.2021.102415>
- Midorikawa, H., Aiba, M., Lebowitz, A., Taguchi, T., Shiratori, Y., Ogawa, T., ... & Tachikawa, H. (2021). Confirming validity of The Fear of COVID-19 Scale in Japanese with a nationwide large-scale sample. *PLoS One*, *16*(2), e0246840. <https://doi.org/10.1371/journal.pone.0246840>
- Muthén, B., & Asparouhov, T. (2013). BSEM measurement invariance analysis. *Mplus Web Notes*, *17*, 1-48.
- Muthén, L. K., & Muthén, B. O. (1998–2014). *Mplus user's guide*. Muthén & Muthén.
- Niedzwiedz, C. L., O'Donnell, C. A., Jani, B. D., Demou, E., Ho, F. K., Celis-Morales, C., ... & Katikireddi, S. V. (2020). Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank. *BMC medicine*, *18*(160), 1-14. <https://doi.org/10.1186/s12916-020-01640-8>
- Olagoke, A. A., Olagoke, O. O., & Hughes, A. M. (2020). Exposure to coronavirus news on mainstream media: The role of risk perceptions and depression. *British Journal of Health Psychology*, *25*, 865-874, e12427. <https://doi.org/10.1111/bjhp.12427>
- Oliver, A., & Mossialos, E. (2004). Equity of access to health care: outlining the foundations for action. *Journal of Epidemiology & Community Health*, *58*(8), 655-658. <https://doi.org/10.1136/jech.2003.017731>

- Oshio, A. (2010). Culture and well-being. In E. Diener, J. E. Helliwell, & D. Kahneman (Eds.), *International Differences in Well-being* (pp. 34-69). Oxford University Press.
- Pfefferbaum, B., & North, C. S. (2020). Mental health and the Covid-19 pandemic. *New England Journal of Medicine*, 383, 510-512. <https://doi.org/10.1056/NEJMp2008017>
- Piqueras, J. A., Gomez-Gomez, M., Marzo, J. C., Gomez-Mir, P., Falco, R., Valenzuela, B., & COVID, P. S. (2020). Validation of the Spanish version of Fear of COVID-19 Scale: Its association with acute stress and coping. *International journal of mental health and addiction*, 1–18. Advance online publication. <https://doi.org/10.1007/s11469-021-00615-x>
- Ramikie, T. S., & Ressler, K. J. (2018). Mechanisms of sex differences in fear and posttraumatic stress disorder. *Biological psychiatry*, 83(10), 876-885. <https://doi.org/10.1016/j.biopsych.2017.11.016>
- Reznik, A., Gritsenko, V., Konstantinov, V., Khamenka, N., & Isralowitz, R. (2020). COVID-19 fear in Eastern Europe: Validation of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00283-3>
- Rhemtulla, M., Brosseau-Liard, P. É., & Savalei, V. (2012). When can categorical variables be treated as continuous? A comparison of robust continuous and categorical SEM estimation methods under suboptimal conditions. *Psychological methods*, 17(3), 354. <https://doi.org/10.1037/a0029315>
- Rutkowski, L., & Svetina, D. (2004). Assessing the hypothesis of measurement invariance in the context of large-scale international surveys. *Educational and Psychological Measurement*, 74, 31–57. <https://doi.org/10.1177/0013164413498257>
- Sakib, N., Bhuiyan, A. I., Hossain, S., Al Mamun, F., Hosen, I., Abdullah, A. H., ... & Mamun, M. A. (2020). Psychometric validation of the Bangla Fear of COVID-19 Scale: Confirmatory factor analysis and Rasch analysis. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00289-x>
- Satici, B., Gocet-Tekin, E., Deniz, M. E., & Satici, S. A. (2020). Adaptation of the Fear of COVID-19 Scale: Its association with psychological distress and life satisfaction in Turkey. *International Journal of Mental Health and Addiction*. Advance online publication. <https://doi.org/10.1007/s11469-020-00294-0>

- Sellbom, M., & Tellegen, A. (2019). Factor analysis in psychological assessment research: Common pitfalls and recommendations. *Psychological assessment, 31*(12), 1428.
<https://doi.org/10.1037/pas0000623>
- Stevens, J. S., & Hamann, S. (2012). Sex differences in brain activation to emotional stimuli: a meta-analysis of neuroimaging studies. *Neuropsychologia, 50*(7), 1578-1593.
<https://doi.org/10.1016/j.neuropsychologia.2012.03.011>
- Thuczek A., Henriques, J.B., & Brown R.L. (2009). Support for the reliability and validity of a six-item state anxiety scale derived from the State-Trait Anxiety Inventory. *Journal of Nursing Measure; 17*(1), 19–28. <https://doi.org/10.1891/1061-3749.17.1.19>
- Torales, J., O'Higgins, M., Castaldelli-Maia, J. M., & Ventriglio, A. (2020). The outbreak of COVID-19 coronavirus and its impact on global mental health. *International Journal of Social Psychiatry, 66*(4), 317–320. <https://doi.org/10.1177/0020764020915212>
- Trougakos, J. P., Chawla, N., & McCarthy, J. M. (2020). Working in a pandemic: Exploring the impact of COVID-19 health anxiety on work, family, and health outcomes. *Journal of Applied Psychology, 105*(11), 1234–1245. <https://doi.org/10.1037/ap10000739>
- Tsipropoulou, V., Nikopoulou, V. A., Holeva, V., Nasika, Z., Diakogiannis, I., Sakka, S., ... & Parlapani, E. (2020). Psychometric Properties of the Greek Version of FCV-19S. *International Journal of Mental Health and Addiction*. Advance online publication.
<https://doi.org/10.1007/s11469-020-00319-8>
- Wakashima, K., Asai, K., Kobayashi, D., Koiwa, K., Kamoshida, S., & Sakuraba, M. (2020). The Japanese version of the Fear of COVID-19 scale: Reliability, validity, and relation to coping behavior. *PLoS one, 15*(11), e0241958. <https://doi.org/10.1371/journal.pone.0241958>
- Vatcheva, K. P., Lee, M., McCormick, J. B., & Rahbar, M. H. (2016). Multicollinearity in regression analyses conducted in epidemiologic studies. *Epidemiology, 6*(2), 227.
<https://doi.org/10.4172/2161-1165.1000227>
- Vallejo, M. A., Vallejo-Slocker, L., Fernández-Abascal, E. G., & Mañanes, G. (2018). Determining factors for stress perception assessed with the Perceived Stress Scale (PSS-4) in Spanish and other European samples. *Frontiers in Psychology, 9*, 37.
<https://doi.org/10.3389/fpsyg.2018.00037>

- VanderWeele, T. J., & Batty, C. J. (2020). On the dimensional indeterminacy of one-wave factor analysis under causal effects. *arXiv preprint arXiv:2001.10352*.
- Wang, C., Pan, R., Wan, X., Tan, Y., Xu, L., Ho, C. S., & Ho, R. C. (2020). Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *International Journal of Environmental Research and Public Health*, 17(5), 1729. <https://doi.org/10.3390/ijerph17051729>
- World Health Organization (2021). *WHO coronavirus disease (COVID-19) dashboard*. From https://covid19.who.int/?gclid=CjwKCAiAv4n9BRA9EiwA30WND9FhCIrWc2SWZG-fZrUMN-cY-2yyYc9lyWc3tyMq1Q2d--Bioxy4iRoCrP0QAvD_BwE accessed at 26.02.2021
- Xiong, J., Lipsitz, O., Nasri, F., Lui, L. M., Gill, H., Phan, L., ... & McIntyre, R. S. (2020). Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *Journal of Affective Disorders*, 277, 55-64. <https://doi.org/10.1016/j.jad.2020.08.001>
- Yuan, K. H., & Bentler, P. M. (2000). 5. Three likelihood-based methods for mean and covariance structure analysis with nonnormal missing data. *Sociological methodology*, 30(1), 165-200. <https://doi.org/10.1111/0081-1750.00078>
- Zhou, X., Sedikides, C., Mo. T., Li, W., Hong, E., & Wildschut, T. (2021). The restorative power of nostalgia: Thwarting loneliness by raising happiness during the COVID-19 pandemic. *Social Psychological and Personality Science*. Advance online publication. <https://doi.org/10.17605/OSF.IO/U5RJB>

Table 1
Sample Characteristics and Language Selected Across Countries

Country	<i>N</i>	% men	<i>M</i> _{age}	<i>SD</i> _{age}	<i>M</i> _{SES}	<i>SD</i> _{SES}	Dominant Language
Total	14557	33.7 ^a	31.30	12.12	4.35	1.07	Various
Armenia	182	35.7	29.63	10.52	4.42	1.12	Armenian
Australia	142	44.4	47.15	17.90	3.95	1.27	English
Austria	310	47.1	40.21	12.05	4.55	1.30	German
Bangladesh	403	56.8	24.29	5.87	3.88	0.70	Bangla
Bosnia & Hercegovina	330	34.8	28.32	11.18	5.01	1.10	Bosnian
Brazil	375	23.5	35.14	12.75	4.34	1.14	Brazilian Portuguese
Bulgaria	282	29.1	36.45	9.97	4.62	0.99	Bulgarian
Chile	220	33.2	30.65	9.98	4.39	1.08	Spanish
China	225	33.3	29.57	11.56	3.93	1.04	Chinese
Colombia	127	41.7	21.80	10.30	5.77	0.91	Spanish
Croatia	239	39.3	36.94	14.70	5.02	1.15	Croatian
Czech	445	18	35.33	13.11	4.60	1.04	Czech
Ecuador	667	36.1	25.73	8.49	4.17	0.89	Spanish
Estonia	261	26.1	40.31	12.69	4.60	1.17	Estonian
Ghana	121	79.3	32.31	6.53	4.07	1.26	English
Hungary	118	10.2	39.59	13.53	4.58	1.02	Hungarian
India	328	49.1	29.78	8.97	4.27	1.08	English
Indonesia	350	22	23.49	8.13	4.53	0.99	Indonesian
Iran	177	46.9	26.50	6.28	4.07	1.13	Farsi
Iraq	146	60.3	29.06	7.29	4.09	1.03	Kurdish
Israel	162	38.3	40.86	16.12	5.02	1.17	Hebrew
Italy	106	23.6	40.08	17.33	4.23	0.89	Italian
Japan	269	79.6	19.82	2.92	4.14	1.11	Japanese
Kazakhstan	242	42.6	26.66	9.27	4.32	1.14	Russian
Latvia	163	30.1	37.80	12.17	4.41	0.95	Latvian
Lebanon	103	20.4	28.55	9.58	4.41	1.06	English
Malaysia	124	18.5	25.65	8.05	4.20	1.02	English
Nigeria	180	45	34.36	9.86	4.37	1.06	English
Pakistan	191	36.6	24.30	5.89	4.69	1.12	Urdu
Peru	151	29.8	29.29	11.42	3.25	1.19	Spanish
Philippines	188	33.5	36.46	13.88	4.78	0.82	Filipino
Poland	279	34.1	31.21	11.69	4.70	1.11	Polish
Portugal	1285	1.7	31.55	8.27	4.18	0.91	Portuguese
Romania	283	24.4	32.55	11.82	4.59	0.97	Romanian
Russia	332	16.6	33.83	12.23	3.81	1.18	Russian
Serbia	872	59.6	24.78	9.72	4.53	1.14	Serbian
Slovakia	227	11	42.66	12.25	4.53	0.91	Slovakian
Slovenia	422	17.3	35.90	11.12	4.16	1.03	Slovenian
Republic of South Africa	992	44	32.86	11.92	3.15	1.40	English
Spain	564	14.5	38.04	14.14	4.06	1.06	Spanish
Thailand	212	38.2	27.50	8.21	4.18	1.06	Thai
Togo	154	55.2	32.10	6.33	3.95	1.14	French
Turkey	399	30.1	25.10	7.27	4.34	0.97	Turkish
Ukraine	340	26.5	27.68	11.44	4.20	1.15	Ukrainian
United Arab Emirates	165	29.7	29.42	7.14	4.44	1.27	English
United Kingdom	259	29.3	34.96	12.14	4.15	1.16	English
Uruguay	160	43.8	43.61	15.28	4.89	0.87	Spanish
Vietnam	279	70.6	21.85	6.40	4.05	1.03	Vietnamese

Note. ^a 0.60% "other" or unreported.

Table 2

Separate Confirmatory Factor Analyses in All Studied Countries: Single-Factor Model With Modifications

Country	χ^2	CFI	RMSEA	90% CI	SRMR	Factor loadings
Armenia	13.02	.989	.059	.000-.114	.021	.58-.81
Australia	16.27	.978	.085	.019-.145	.027	.67-.87
Austria	7.43	1.00	.000	.000-.064	.013	.58-.78
Bangladesh	28.5	.984	.080	.049-.112	.020	.66-.83
Bosnia & Hercegovina	14.89	.989	.051	.000-.091	.017	.55-.84
Brazil	9.44	.999	.022	.000-.067	.014	.63-.77
Bulgaria	15.35	.990	.057	.000-.100	.014	.36-.83
Chile	38.94	.955	.133	.093-.176	.031	.57-.82
China	10.05	.997	.034	.000-.089	.019	.59-.77
Colombia	7.59	1.00	.000	.000-.101	.020	.53-.90
Croatia	20.33	.978	.080	.037-.124	.028	.61-.86
Czech	39.64	.959	.094	.066-.124	.026	.62-.81
Ecuador	21.58	.994	.050	.025-.077	.015	.62-.85
Estonia	27.41	.968	.096	.058-.137	.026	.61-.80
Ghana	8.39	.999	.020	.000-.110	.021	.59-.83
Hungary	8.37	.999	.020	.000-.111	.021	.58-.74
India	31.09	.966	.094	.060-.130	.032	.62-.82
Indonesia	11.83	.995	.037	.000-.078	.020	.49-.82
Iran	17.1	.979	.080	.024-.133	.032	.51-.86
Iraq	15.52	.979	.080	.002-.140	.033	.56-.76
Israel	33.54	.932	.140	.093-.191	.041	.53-.76
Italy	31.19	.929	.165	.107-.228	.041	.62-.79
Japan	4.77	1.00	.000	.000-.048	.020	.33-.96
Kazakhstan	15.09	.987	.061	.000-.107	.029	.45-.77
Latvia	8.75	.998	.024	.000-.097	.021	.21-.93
Lebanon	15.21	.970	.094	.000-.165	.028	.58-.86
Malaysia	17.42	.970	.097	.032-.160	.030	.60-.90
Nigeria	12.95	.992	.059	.000-.115	.018	.69-.84
Pakistan	11.64	.992	.049	.000-.115	.018	.67-.90
Peru	10.33	.993	.044	.000-.111	.028	.51-.84
Philippines	17.97	.984	.081	.030-.132	.027	.51-.88
Poland	24.52	.970	.086	.049-.126	.022	.44-.81
Portugal	28.36	.994	.045	.027-.063	.012	.61-.76
Romania	13.97	.992	.051	.000-.095	.016	.65-.76
Russia	9.41	.998	.023	.000-.071	.014	.40-.72
Serbia	24.94	.986	.049	.028-.072	.020	.54-.79
Slovakia	26.35	.966	.101	.060-.144	.028	.47-.81
Slovenia	48.15	.961	.109	.081-.140	.027	.67-.81
South Africa	23.67	.994	.044	.024-.066	.013	.60-.91
Spain	18.76	.994	.049	.020-.078	.013	.61-.86
Thailand	11.99	.994	.049	.000-.101	.014	.66-.89
Togo	11.68	.992	.055	.000-.117	.019	.70-.82
Turkey	20.66	.991	.063	.030-.097	.013	.62-.85
Ukraine	16.25	.987	.055	.012-.094	.022	.53-.72
United Arab Emirates	18.85	.979	.091	.037-.144	.028	.50-.87
United Kingdom	15.52	.990	.060	.002-.105	.021	.71-.84
Uruguay	8.62	.998	.022	.000-.097	.022	.61-.77
Vietnam	6.01	1.00	.000	.000-.058	.009	.54-.79

Note. Covariances estimated freely: 3+6, 3+7, 6+7, 1+2, 1+4, 2+5; $df = 8$.

Table 3*Multigroup Confirmatory Factor Analysis of Fear of COVID-19 Scale in 43 Countries*

Multigroup Confirmatory Factor Analysis					
Invariance level	χ^2	CFI	RMSEA	90% CI	SRMR
Configural ($df = 344$)	624.23	.989	.057	.051-.063	.020
Metric ($df = 596$)	1470.38	.966	.076	.072-.081	.084
Partial Metric ($df = 581$)	1169.98	.976	.065	.061-.070	.062
Scalar ($df = 848$)	5477.74	.823	.146	.143-.150	.136
Levels of Invariance comparison					
	$\Delta \chi^2$	Δ CFI	Δ RMSEA	Δ SRMR	Δdf
Configural vs. Metric	986.90	-.023	.019	.064	252
Configural vs. Partial Metric	666.10	-.013	.008	.042	237
Partial Metric vs. Scalar	4642.97	-.137	.076	.065	252

Note. $N = 13419$. All χ^2 difference tests were significant at $p < .001$. Samples not used in MGCFA: Chile, Israel, Italy, Slovakia, Slovenia, Serbia.

Table 4

Confirmatory Factor Analysis and Multi-Group Confirmatory Factor Analysis in Men and Women: Single-Factor Model with Modifications

Separate Confirmatory Factor Analyses												
Gender	χ^2	CFI	RMSEA	90% CI	SRMR	Factor loadings						
Men	65.40	.996	.038	.030-.047	.009	.72-.82						
Women	135.29	.995	.041	.035-.047	.009	.69-.80						
Multigroup Confirmatory Factor Analysis												
Invariance level	χ^2	CFI	RMSEA	90% CI	SRMR	Factor loadings						
Configural ($df = 16$)	191.79	.995	.039	.034-.044	.009	-						
Metric ($df = 22$)	262.94	.994	.039	.035-.043	.016	-						
Scalar ($df = 28$)	682.00	.983	.057	.053-.061	.028	-						
Partial scalar ($df = 25$)	289.87	.993	.038	.034-.042	.017	-						
Levels of Invariance comparison												
	$\Delta \chi^2$	Δ CFI	Δ RMSEA	Δ SRMR	Δ df							
Configural vs. Metric	71.15	-.001	.000	.007	6							
Metric vs. Scalar	419.06	-.011	.018	.012	6							
Metric vs. Partial Scalar	26.93	-.001	-.001	.001	3							
Descriptive statistics												
	Fear of Covid			Anxiety			Stress			Correlations		
	M	SD	α	M	SD	α	M	SD	α	r_{f-a}	r_{f-s}	r_{a-s}
Men	2.90	1.50	.92	2.64	0.68	.81	2.70	0.68	.63	.34**	.30**	.53**
Women	3.11	1.38	.90	2.89	0.69	.86	2.84	0.72	.73	.34**	.29**	.59**

Note. $N = 14470$; covariances estimated freely: 3+6, 3+7, 6+7, 1+2, 1+4, 2+5; $df = 8$.

We excluded 85 participants from the analysis for not reporting sex or reporting "other". χ^2 difference tests were both significant, $p < .001$. We calculated correlation coefficients based on latent factor scores estimated with MGCFAs. r_{f-a} = correlation between the FCV-19S and STAI-6, r_{f-s} = correlation between the FCV-19S and PSS-4, r_{a-s} = correlation between STAI-6 and PSS-4.

Table 5

Confirmatory Factor Analysis and Multi-Group Confirmatory Factor Analysis in Different Education Levels: Single-Factor Model with Modifications (Covariances Estimated Freely: 3+6, 3+7, 6+7, 1+2, 1+4, 2+5; $df = 8$)

Separate Confirmatory Factor Analyses												
Gender	χ^2	CFI	RMSEA	90% CI	SRMR	Factor loadings						
Primary	11.43	.993	.051	.000-.112	.026	.62-.86						
Lower Secondary	6.36	1.00	.000	.000-.040	.008	.73-.86						
Upper Secondary	62.45	.996	.040	.031-.050	.009	.73-.83						
Bachelor or equivalent	68.04	.996	.037	.029-.045	.009	.67-.80						
Master or equivalent	41.46	.996	.036	.025-.047	.009	.64-.77						
PhD	38.33	.985	.070	.049-.093	.018	.65-.76						
Multigroup Confirmatory Factor Analysis												
Invariance level	χ^2	CFI	RMSEA	90% CI	SRMR	Factor loadings						
Configural ($df = 48$)	226.08	.995	.039	.034-.044	.010	-						
Metric ($df = 78$)	362.66	.993	.039	.034-.043	.023	-						
Scalar ($df = 108$)	652.68	.986	.046	.042-.049	.025	-						
Partial Scalar ($df = 101$)	463.10	.991	.038	.035-.042	.022	-						
Levels of Invariance comparison												
	$\Delta \chi^2$	Δ CFI	Δ RMSEA	Δ SRMR	Δ df							
Configural vs. Metric	136.58	-.002	.000	.013	30							
Metric vs. Scalar	290.02	-.007	.007	.002	30							
Metric vs. Partial Scalar	100.44	-.002	-.001	.001	23							
Fear of Covid-19 descriptive statistics												
	Fear of Covid			Anxiety			Stress			Correlations		
	<i>M</i>	<i>SD</i>	α	<i>M</i>	<i>SD</i>	α	<i>M</i>	<i>SD</i>	α	r_{f-a}	r_{f-s}	r_{a-s}
Primary	4.08	1.62	.91	2.82	0.68	.68	2.96	0.60	.08	-	.30**	-
Lower Secondary	3.39	1.68	.93	2.78	0.76	.84	2.85	0.77	.68	.31**	.35**	.59**
Upper Secondary	3.13	1.55	.92	2.81	0.72	.84	2.86	0.71	.67	.26**	.33**	.58**
Bachelor	3.05	1.36	.90	2.84	0.69	.85	2.81	0.70	.72	.28**	.34**	.57**
Master	2.84	1.24	.89	2.79	0.67	.86	2.70	0.69	.72	.33**	.39**	.58**
PhD	2.80	1.26	.90	2.66	0.67	.86	2.59	0.68	.72	.38**	.43**	.55**

Note. $N = 14557$; covariances estimated freely: 3+6, 3+7, 6+7, 1+2, 1+4, 2+5; $df = 8$.

χ^2 difference tests were both significant, $p < .001$. We calculated correlation coefficients based on latent factor scores estimated with MGCFA. r_{f-a} = correlation between the FCV-19S and STAI-6, r_{f-s} = correlation between the FCV-19S and PSS-4, r_{a-s} = correlation between STAI-6 and PSS-4.

** $p < .01$.

Table 6

Descriptive Statistics, Reliability, and Correlation Coefficients Between Studied Variables in All Countries

Country	Fear of Covid			Anxiety			Stress			Correlations		
	<i>M</i>	<i>SD</i>	α	<i>M</i>	<i>SD</i>	α	<i>M</i>	<i>SD</i>	α	r_{f-a}	r_{f-s}	r_{a-s}
Armenia	2.47	1.20	0.89	2.91	0.68	0.83	2.83	0.69	0.62	.39**	.27**	.42**
Australia	3.02	1.29	0.91	2.65	0.77	0.91	2.65	0.73	0.78	.23**	.22**	.60**
Austria	1.91	0.87	0.84	2.67	0.74	0.88	2.47	0.80	0.82	.46**	.38**	.70**
Bangladesh	3.49	1.39	0.90	2.90	0.78	0.86	2.84	0.75	0.69	.28**	.20**	.67**
Bosnia & Hercegovina	2.23	1.07	0.87	2.64	0.63	0.87	2.75	0.60	0.62	.20**	.09	.50**
Brazil	3.95	1.20	0.88	3.14	0.64	0.85	3.04	0.71	0.75	.50**	.30**	.64**
Bulgaria	2.56	1.04	0.86	2.69	0.66	0.88	2.6	0.63	0.77	.44**	.36**	.62**
Chile	3.32	1.36	0.89	3.04	0.73	0.88	2.84	0.80	0.82	.34**	.18**	.69**
China	3.32	1.20	0.89	2.71	0.52	0.79	2.71	0.59	0.71	.22**	.35**	.60**
Colombia	3.09	1.23	0.86	2.86	0.66	0.85	2.69	0.74	0.74	.31**	.21**	.56**
Croatia	2.19	1.01	0.88	2.63	0.65	0.87	2.68	0.63	0.61	.27**	.24**	.54**
Czech	2.11	0.91	0.85	2.74	0.72	0.92	2.68	0.76	0.81	.40**	.35**	.70**
Ecuador	3.69	1.38	0.90	2.84	0.66	0.83	2.80	0.65	0.67	.34**	.27**	.56**
Estonia	2.03	0.90	0.88	2.53	0.65	0.86	2.49	0.71	0.83	.34**	.27**	.71**
Ghana	3.45	1.51	0.90	2.43	0.63	0.77	2.65	0.60	0.46	.52**	.31**	.47**
Hungary	2.37	1.07	0.87	2.99	0.77	0.87	2.78	0.75	0.81	.45**	.40**	.63**
India	3.17	1.32	0.89	2.73	0.67	0.81	2.76	0.64	0.54	.41**	.32**	.56**
Indonesia	3.21	1.07	0.84	2.86	0.59	0.81	2.83	0.62	0.73	.29**	.25**	.61**
Iran	3.01	1.29	0.88	2.94	0.72	0.88	2.96	0.66	0.69	.22**	.16*	.60**
Iraq	2.86	1.17	0.85	2.91	0.80	0.87	2.70	0.61	0.54	.15	.23**	.42**
Israel	2.37	1.05	0.87	2.85	0.74	0.91	2.53	0.68	0.68	.48**	.36**	.71**
Italy	2.70	1.07	0.89	2.89	0.67	0.87	2.78	0.75	0.82	.43**	.31**	.69**
Japan	4.11	1.06	0.83	2.88	0.53	0.57	2.99	0.47	0.27	.19**	.08	.41**
Kazakhstan	2.61	1.11	0.85	2.58	0.65	0.84	2.65	0.73	0.75	.28**	.17*	.50**
Latvia	3.81	1.30	0.92	2.99	0.49	0.77	2.92	0.51	0.57	.41**	.28**	.68**
Lebanon	2.98	1.20	0.85	3.40	0.67	0.88	3.31	0.74	0.80	.09	.06	.68**
Malaysia	3.47	1.17	0.87	2.92	0.54	0.82	2.90	0.65	0.77	.18	.12	.53**
Nigeria	2.86	1.42	0.92	2.42	0.63	0.79	2.54	0.67	0.65	.29	.24	.54**
Pakistan	3.19	1.35	0.92	2.72	0.56	0.64	2.78	0.53	0.16	.13	.25**	.21**
Peru	3.81	1.35	0.88	2.86	0.66	0.85	2.77	0.66	0.65	.31**	.32**	.60**
Philippines	3.84	1.29	0.89	2.78	0.58	0.82	2.81	0.62	0.60	.38**	.24**	.57**
Poland	2.13	0.90	0.82	2.94	0.75	0.88	2.87	0.82	0.81	.39**	.27**	.68**
Portugal	3.52	1.20	0.86	3.07	0.64	0.87	2.86	0.73	0.81	.38**	.31**	.64**
Republic of South Africa	4.47	1.60	0.91	2.84	0.63	0.65	3.04	0.61	0.29	.37**	.31**	.61**
Romania	2.68	1.16	0.90	2.65	0.59	0.86	2.67	0.64	0.74	.41**	.22**	.55**
Russia	2.52	0.99	0.83	2.81	0.67	0.88	2.89	0.75	0.79	.31**	.21**	.57**
Serbia	1.84	0.89	0.84	2.44	0.72	0.85	2.70	0.71	0.65	.35**	.37**	.59**
Slovakia	2.57	0.94	0.87	2.52	0.56	0.83	2.58	0.60	0.67	.38**	.30**	.63**
Slovenia	2.36	1.15	0.89	2.51	0.76	0.90	2.62	0.77	0.81	.23**	.31**	.74**
Spain	2.96	1.31	0.90	3.04	0.79	0.89	2.89	0.93	0.86	.37**	.29**	.70**
Thailand	3.27	1.28	0.90	2.92	0.59	0.80	2.94	0.63	0.60	.23**	.32**	.49**
Togo	3.56	1.65	0.91	2.74	0.57	0.76	2.74	0.60	0.55	.20*	.20*	.60**
Turkey	3.05	1.35	0.90	2.99	0.67	0.84	3.09	0.62	0.62	.31**	.16**	.46**
Ukraine	2.42	1.00	0.83	2.71	0.66	0.86	2.75	0.66	0.71	.27**	.22**	.60**

United Arab Emirates	4.27	1.03	0.86	3.32	0.60	0.77	2.80	0.61	0.76	.40**	.36**	.47**
United Kingdom	2.94	1.27	0.91	2.80	0.76	0.91	2.85	0.75	0.76	.44**	.38**	.79**
Uruguay	2.39	0.97	0.87	2.60	0.60	0.84	2.28	0.67	0.74	.14	.05	.62**
Vietnam	3.78	1.34	0.88	2.87	0.54	0.82	2.82	0.57	0.65	.17**	.26**	.46**

Note. We calculated correlation coefficients based on latent factor scores estimated via MGCFA. We used italics to designate correlation coefficients calculated with observed means due to the non-invariance of one of the measures. r_{f-a} = correlation between the FCV-19S and STAI-6, r_{f-s} = correlation between the FCV-19S and PSS-4, r_{a-s} = correlation between STAI-6 and PSS-4. * $p < .05$, ** $p < .01$.

Table 8*Item-level Descriptive Statistics of Fear of Covid-19 Scale*

Item	<i>M</i> [range]	<i>SD</i> [range]	Skewness [range]	Kurtosis [range]
1. I am most afraid of coronavirus-19	3.71 [1.83, 5.60]	1.60 [1.01, 2.06]	0.07 [-1.06, 1.58]	-0.42 [-1.29, 3.25]
2. It makes me uncomfortable to think about coronavirus-19	3.96 [2.45, 5.80]	1.69 [1.04, 2.10]	-0.13 [-0.89, 0.98]	-0.81 [-1.42, 1.37]
3. My hands become clammy when I think about coronavirus-19	2.20 [1.27, 4.01]	1.29 [0.69, 2.04]	1.33 [-0.22, 2.77]	2.16 [-1.44, 11.39]
4. I am afraid of losing my life because of coronavirus-19	3.17 [1.43, 5.13]	1.69 [0.86, 2.25]	0.56 [-0.87, 2.81]	0.06 [1.49, 9.53]
5. When I watch news and stories about coronavirus-19 on social media, I become nervous or anxious	3.55 [2.25, 5.70]	1.73 [1.27, 2.11]	0.11 [-1.22, 1.02]	-0.87 [-1.45, 1.51]
6. I cannot sleep because I'm worrying about getting coronavirus-19	2.07 [1.21, 3.41]	1.27 [0.67, 1.98]	1.56 [-0.08, 4.57]	3.16 [-1.47, 25.82]
7. My heart races or palpitates when I think about getting coronavirus-19	2.35 [1.25, 3.92]	1.45 [0.76, 2.05]	1.26 [-0.25, 3.58]	1.64 [-1.37, 13.49]

Note. Detailed descriptive statistics for all items across studied countries can be found in the project's OSF page: <https://osf.io/hxmzb>

Figure 1. *The Modified Model of the Fear of COVID-19 Scale Tested in Multigroup Confirmatory Factor Analyses*




Figure 2

Distributions of Fear of COVID-19 Scores Estimated with the Alignment Procedure Across 43 Countries. Vertical Lines Represent Medians

