



LEEDS  
BECKETT  
UNIVERSITY

---

Citation:

Arumi-Prat, I and Cirera, E and McKenna, J and Puig-Ribera, AM (2022) From Secondary School to University: Associations Between Domain-Specific Sedentary Behaviors and Lifestyle Risk Behaviors. *Inquiry*, 59. p. 469580221118843. ISSN 1945-7243 DOI: <https://doi.org/10.1177/00469580221118843>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/8984/>

Document Version:

Article (Published Version)

---

Creative Commons: Attribution-Noncommercial 4.0

© The Author(s) 2022

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

# From Secondary School to University: Associations Between Domain-Specific Sedentary Behaviors and Lifestyle Risk Behaviors

INQUIRY: The Journal of Health Care  
Organization, Provision, and Financing  
Volume 59: 1–12  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/00469580221118843  
journals.sagepub.com/home/inq



Ignasi Arumi-Prat, PhD<sup>1</sup> , Eva Cirera, PhD<sup>1</sup>,  
Jim McKenna, PhD<sup>2</sup>, and Anna Maria Puig-Ribera, PhD<sup>1</sup>

## Abstract

This study examined associations between changes in domain-specific sedentary behaviors and changes in health-related lifestyles of Spanish secondary school students ( $n=113$ ) to their first year of university. During the transitions from the end of high school to the beginning of university, engagement in sedentary behaviors have emerged as potential additional behavioral risk factors. Understanding how sedentary behaviors interconnect with other (un)healthy behaviors will inform interventions on multiple risk behaviors across this critical life period. A 3-year longitudinal survey assessed associations between domain-specific sedentary behaviors and leisure time physical activity (IPAQ), alcohol and tobacco consumption, and fruit and vegetable intake (24-h dietary recall), using Generalized Estimating Equations. Spending time on sedentary transportation was associated with a greater likelihood of smoking, whereas sedentary weekend homework was associated with a reduced likelihood of consuming alcohol. The lowest and highest tertiles for sedentary screen use and leisure-time PA were also less likely not to meet the recommendations for fruit and vegetable consumption. For specific sedentary behaviors, associations were gender-based or affected by leisure time physical activity. From secondary school to university, specific sedentary behaviors are linked to lifestyle risk factors. Over this transitional period, public health interventions targeting reduced sedentary behaviors may bring multiple benefits by also preventing other harmful behaviors.

## Keywords

behavior, sedentary, unhealthy, adolescents, physical activity

### What is already known on this topic?

The adolescence-university transition is a critical life period to understand how lifestyle behaviors may cluster, to target disease prevention and promote sustainable health-enhancing behavior into adulthood and for the next generations. While time spent sitting and reclining while expending little energy—named sedentary behavior—and engagement in specific sedentary activities have emerged as potential additional behavioral risk factors, observational and interventional evidence investigating how to effectively tackle sedentary behavior in older adolescents and young adults going through this major life transition is limited, with evidence dominated by research in younger adolescents (10-14 years) and physical inactivity. Even less is known about how specific SB domains influence different lifestyle behaviors, a key issue for developing public health interventions that can effectively modify SBs while also influencing other problematic behaviors.

### What this study adds?

From secondary school to university, specific SBs are associated to lifestyle risk factors some of which are gender-based and/or related to leisure-time PA. Time sitting doing weekend homework was protective against tobacco use in adolescents who spent little time doing leisure PA. In girls, doing sedentary weekend homework also protected against alcohol consumption. Sedentary weekend transport time was associated with higher tobacco consumption while the lowest and highest tertiles for sedentary screen use and leisure-time PA were less likely not to meet the recommendations for fruit and vegetable consumption.

### What are your research's implications toward theory, practice, or policy?

Over this transitional period, implementing public health interventions targeting reduced SBs may bring multiple benefits by also preventing other harmful behaviors.



## Introduction

Non-communicable diseases (NCDs) are the biggest cause of premature deaths worldwide.<sup>1</sup> While many NCDs manifest later in adulthood, these are partly the result of behavioral risk factors established over time, which typically begin in adolescence.<sup>2,3</sup> In 2019, NCDs in adolescents aged 10 to 24 years accounted for 86.4% of all years lived with disability and 38.8% of total deaths.<sup>4</sup> In this context, it is critical to develop a better understanding of how interventions could effectively be implemented for tackling those behavioral risk factors that drive NCDs in adolescence and continuing into adulthood.<sup>2,3,5</sup>

In adolescents, NCDs can be prevented by tackling common behavioral risk behaviors, mainly tobacco use, low fruit and vegetable intake, poor diet, physical inactivity, drug use, and harmful use of alcohol.<sup>5-7</sup> Additionally, time spent sitting and reclining while expending little energy—named sedentary behavior (SB)—and engagement in specific sedentary activities such as screen-based behaviors have emerged as potential additional behavioral risk factors for adolescents' health and well-being.<sup>3</sup> However, observational and interventional evidence investigating how to effectively tackle sedentary behavior in older adolescents and young adults going through this major life transition is limited, with evidence dominated by research in younger adolescents (10-14 years) and physical inactivity.<sup>2,3</sup>

During this life transition, physical activity (PA) diminishes, driven in part by reductions in active transportation<sup>8</sup> and active commuting to school.<sup>9</sup> Involvement in sport also drops while engagement in sedentary behaviors (SBs) rises, especially due to increased sitting time while socializing, for transport<sup>10</sup> and in recreational screen-time.<sup>11,12</sup> Indeed, at universities many young adults report higher sedentary behavior after the COVID-19 pandemic,<sup>13</sup> with prolonged sitting being associated with worse physical and mental health, behavioral conduct, and reduced sleep duration.<sup>14</sup> Just as SBs increase from secondary school to university,<sup>10</sup> so too does alcohol consumption, average daily cigarette consumption<sup>15</sup> and fruit and vegetable intake declines.<sup>16</sup> However, evidence on the influence SB changes have on other modifiable unhealthy behaviors over this life period is scarce.

In this context, monitoring changes of SB patterns over the adolescent-university transition period in association with changes in other behavioral risk behaviors is important for

developing public health interventions that can effectively modify SBs while also influencing other problematic behaviors. Given that not all SBs are equally harmful, and that most studies have focused on cross-sectional associations between recreational screen-based sedentary behavior, alcohol use,<sup>17</sup> earlier initiation of cannabis and tobacco consumption,<sup>18</sup> poor diet,<sup>19</sup> and individual sport participation,<sup>10</sup> it is important to understand how specific SB domains influence different life-style behaviors.<sup>14</sup>

To understand whether SBs displace and/or activate (un) healthy behaviors in the adolescence-university transition, this study undertook formative research to inform public health interventions on how to effectively tackle the behavioral risk factors that jeopardize adolescents into adulthood' health. The present study examined associations between changes in domain-specific sedentary behaviors and changes in health-related lifestyles of Spanish secondary school students (16-17 years old) to their first year of university (18-19 years old).

## Methods

### *Study Design and Sample Recruitment*

A 3-year longitudinal study was designed to assess associations between changes in domain-specific SBs and lifestyle risk behaviors in Spanish adolescents (n=113) from the county of Osona (Barcelona). Adolescents were followed from secondary school to university (16, 17, and 18 years of age; Year 1, 2, and 3 respectively). From an initial potential sample of 695 teenagers, 662 responded in Year 1 (95% response rate), 480 in Year 2 (69% response rate), and 180 in Year 3 (26% response rate). Only the university undergraduate students (n=113, 16% of the initial potential sample and 17% of Year 1 respondents) who completed the survey in all 3 years were included. The Ethics Committee of University of Vic-Central University of Catalonia approved the study (2011), and all participants signed a written informed consent every year before completing the survey. Recruitment procedures have been described in detail elsewhere.<sup>10</sup>

### *Data Collection and Variables*

Data were collected using a 42-item survey that gathered data on (i) sociodemographic variables (age, gender, height, weight, and place of residence), (ii) domain-specific SBs,

<sup>1</sup>University of Vic-Central University of Catalonia, Barcelona, Spain

<sup>2</sup>Leeds Beckett University, Leeds, UK

Received 28 March 2022; revised 6 July 2022; revised manuscript accepted 20 July 2022

### **Corresponding Author:**

Eva Cirera, Sport and Physical Activity Research Group, Centre for Health and Social Care Research, Department of Experimental Science and Methodology, Faculty of Health Sciences and Welfare, University of Vic-Central University of Catalonia, Vic, Barcelona 08500, Spain.

Email: [eva.cirera@uvic.cat](mailto:eva.cirera@uvic.cat)

and (iii) lifestyle risk behaviors (tobacco and alcohol consumption; fruit and vegetable consumption; and leisure-time physical activity).

### *Tobacco and Alcohol Consumption*

Alcohol and tobacco consumption were recorded using the FRESQ questionnaire, which has good reliability for both alcohol ( $r = .66-.72$ ) and tobacco consumption ( $r = .79-.82$ ).<sup>20</sup> Alcohol measures included alcohol consumption over the past 12 months (yes/no), and frequency of consumption (daily, weekly, monthly, <monthly, never). Tobacco consumption variables included current cigarette consumption (yes/no), and frequency of use (<1 cigarette a day, 1 cigarette a day, between 2 and 5 cigarettes a day, more than 5 cigarettes a day). Both alcohol and tobacco variables were categorized as weekly alcohol consumption (yes/no) and daily tobacco consumption (yes/no).

### *Fruit and Vegetable Consumption*

Fruit and vegetable consumption was assessed with 2 specific questions: “How many servings of fruit do you eat on a typical day?” and “How many servings of vegetables do you eat on a typical day?” using the 24-h dietary recall data as the gold standard.<sup>21</sup> Cronbach alpha for this measure was .74 among university students.<sup>22</sup> Under-consumption was defined by yes/no responses to an item about eating 5 servings of fruits and/or vegetables a day.<sup>21</sup>

### *Leisure-Time Physical Activity*

Leisure-time PA was measured using the Spanish version of the International Physical Activity Questionnaire (IPAQ) long form.<sup>23</sup> The IPAQ assessed min/week of light-intensity PA (LPA), moderate-intensity PA (MPA), and vigorous-intensity PA (VPA) during the last 7 days. The IPAQ has shown good validity for assessing different intensities of PA domains in healthy European adolescents aged 15 to 17 years ( $R_s = .17-.30$ ).<sup>24</sup> Leisure-time PA was categorized as the sum of time spent in PA at all intensities (min/week) in the following tertiles: (i) less than 180 min/week, (ii) between 180 and 259 min/week, and (iii) more than 360 min/week.

### *Domain-Specific Sedentary Behavior*

The sedentary behavior questionnaire (Active Where? survey—Section R)<sup>25</sup> assessed sitting time (min/day) during weekdays and weekends and across domains<sup>26</sup>: (1) television viewing (television + video); (2) computer use (computer games + internet use); (3) socializing behaviors (sitting with friends); (4) school (school attendance + homework); (5) transport (private + public transport); and (6) sedentary hobbies (reading, playing music, and doing handicrafts). Responses were categorized into 15-min blocks, 30-min blocks, and 1-h blocks, concluding with  $\geq 5$  h. The Active Where? survey was

designed specifically for youth and has shown good reliability in most sitting domains, with a percentage agreement ranging from 27.1% to 76%.<sup>25</sup> Time spent (min/day) on each SB-domain during weekdays and weekends were categorized into tertiles and “no use” to describe 0 min/day of time spent in that specific SB-domain. A new variable on “screen use time” was described as the sum of SBs spent watching TV and using computers (computer games + internet use) and categorized into 2 groups according to the 24-h movement guidelines for children and teenagers from 5 to 17 years old<sup>27</sup>: less than 120 min/day, or 120 or more minutes/day.

### *Statistical Analysis*

A descriptive analysis of the subjects' characteristics by year (Years 1, 2, and 3) was performed using proportions according to data type ( $n = 113$ ). The temporal variation of each lifestyle risk behavior and SB-specific domains across years from secondary school to university was described using proportions by year.

Generalized Estimating Equations (GEE) assessed associations between lifestyle risk behaviors (dependent variables) and each domain-specific SBs (independent variables).<sup>28</sup> This methodology is useful for analyzing repeated measures of the same individual over time, assuming independence between individuals but not within observations of the same individual. Associations between variables were modeled separately; weekdays and weekends. A binomial distribution of the dependent variables (lifestyle risk factors) was assumed and logit was used as the link function. The starting models included each SB-specific domain, adjusted by all possible measured confounders (gender, year, and leisure-time PA, as well as interactions between SB-specific domain and gender and, between SB-specific domain and leisure-time PA).

The final models included only the SB-specific domain and the adjusted variables identified as confounders (ie, those variables that could change SB-specific domain coefficients for more than 10%).<sup>29</sup> Odds Ratios (OR) and 95% CI are shown graphically for SB-specific domains, indicating, in each case, the adjusted variables and the interactions included in the model. The analysis was performed using STATA software 12.

## **Results**

Participants' baseline characteristics ( $n = 113$ ) are summarized in Table 1 (year 1).

### *Temporal Variations in Lifestyle Risk Behaviors, Domain-Specific SBs and Leisure-Time PA*

From secondary school to university, the proportion of habitual smokers and weekly consumers of alcohol increased by +2.7% and +8.8% respectively. Similarly, the proportion of adolescents not meeting the recommendations for daily fruit and vegetable consumption increased by +8.3% (Table 1).

**Table 1.** Patterns of Lifestyle Behaviors and Different Sedentary Domains Across Years.

	Year 1	Year 2	Year 3	Total
	n (%)	n (%)	n (%)	n (%)
Boys	48 (42.5)	48 (42.5)	48 (42.5)	144 (42.5)
Girls	65 (57.5)	65 (57.5)	65 (57.5)	195 (57.5)
Habitual smokers	30 (26.5)	31 (27.4)	33 (29.2)	94 (27.7)
Weekly alcohol consumption	15 (13.3)	21 (18.6)	25 (22.1)	61 (18.0)
<5 Fruit and veg per day	79 (70.5)	80 (70.8)	89 (78.8)	248 (73.4)
Weekly leisure time PA				
<180'	30 (26.5)	27 (23.9)	44 (38.9)	101 (29.8)
180'-359'	32 (28.3)	38 (33.6)	37 (32.7)	107 (31.6)
360+	51 (45.1)	48 (42.5)	32 (28.3)	131 (38.6)
Computer use				
Weekdays				
No use	5 (4.4)	3 (2.7)	1 (0.9)	9 (2.7)
≤60	36 (31.9)	56 (49.6)	43 (38.1)	135 (39.8)
61-120	38 (33.6)	35 (31.0)	39 (34.5)	112 (33.0)
121+	34 (30.1)	19 (16.8)	30 (26.5)	83 (24.5)
Weekends				
No use	1 (0.9)	4 (3.5)	8 (7.1)	13 (3.8)
≤60	25 (22.1)	39 (34.5)	46 (40.7)	110 (32.4)
61-180	49 (43.4)	47 (41.6)	39 (34.5)	135 (39.8)
181+	38 (33.6)	23 (20.4)	20 (17.7)	81 (23.9)
TV viewing				
Weekdays				
No use	15 (13.3)	10 (8.8)	10 (8.8)	35 (10.3)
≤30	34 (30.1)	37 (32.7)	35 (31.0)	106 (31.3)
31-90	32 (28.3)	32 (28.3)	34 (30.1)	98 (28.9)
91+	32 (28.3)	34 (30.1)	34 (30.1)	100 (29.5)
Weekends				
No use	12 (10.6)	8 (7.1)	19 (16.8)	39 (11.5)
≤60	44 (38.9)	44 (38.9)	36 (31.9)	124 (36.6)
61-120	28 (24.8)	35 (31.0)	32 (28.3)	95 (28.0)
121+	29 (25.7)	26 (23.0)	26 (23.0)	81 (23.9)
Screen use				
Weekdays				
≤120'	68 (39.8)	54 (47.80)	39 (34.5)	138 (40.7)
>120'	45 (60.2)	59 (52.2)	74 (65.5)	201 (59.3)
Weekends				
≤120'	21 (18.6)	22 (19.50)	41 (36.3)	84 (24.8)
>120'	92 (81.4)	91 (80.5)	72 (63.7)	255 (75.2)
Homework/school				
Weekdays				
≤390	22 (19.5)	21 (18.6)	67 (59.3)	110 (32.4)
391-480	64 (56.6)	67 (59.3)	28 (24.8)	159 (46.9)
481+	27 (23.9)	25 (22.1)	18 (15.9)	70 (20.7)
Weekends				
No use	6 (5.3)	9 (8.0)	20 (17.7)	35 (10.3)
≤60	52 (46.0)	53 (46.9)	27 (23.9)	132 (38.9)
61-120	25 (22.1)	22 (19.5)	34 (30.1)	81 (23.9)
121+	30 (26.6)	29 (25.7)	32 (28.3)	91 (26.8)

(continued)

**Table 1. (continued)**

	Year 1	Year 2	Year 3	Total
	n (%)	n (%)	n (%)	n (%)
<b>Sedentary socialization</b>				
Weekdays				
No use	12 (10.6)	11 (9.7)	5 (4.4)	28 (8.3)
≤30	56 (49.6)	50 (44.2)	38 (33.6)	144 (42.5)
31-60	33 (29.2)	32 (28.3)	30 (26.5)	95 (28.0)
61+	12 (10.6)	20 (17.7)	40 (35.4)	72 (21.2)
Weekends				
No use	9 (8.0)	5 (4.4)	2 (1.8)	16 (4.7)
≤60	55 (48.7)	47 (41.6)	21 (18.6)	123 (36.3)
61-150	19 (16.8)	28 (24.8)	46 (40.7)	93 (27.4)
151+	30 (26.5)	33 (29.2)	44 (38.9)	107 (31.6)
<b>Sedentary transport</b>				
Weekdays				
No use	26 (23.0)	26 (23.0)	15 (13.3)	67 (19.8)
≤15	40 (35.4)	39 (34.5)	18 (15.9)	97 (28.6)
16-30	36 (31.9)	36 (31.9)	23 (20.4)	95 (28.0)
31+	11 (9.7)	12 (10.6)	57 (50.4)	80 (23.6)
Weekends				
No use	24 (21.2)	23 (20.4)	22 (19.5)	69 (20.4)
≤15	16 (14.2)	15 (13.3)	14 (12.4)	45 (13.3)
16-60	57 (50.4)	60 (53.1)	57 (50.4)	174 (51.3)
61+	16 (14.2)	15 (13.3)	20 (17.7)	51 (15.0)
<b>Sedentary hobbies</b>				
Weekdays				
No use	29 (25.7)	49 (43.4)	30 (26.5)	108 (31.9)
≤15	13 (11.5)	14 (12.4)	15 (13.3)	42 (12.4)
16-60	31 (27.4)	24 (21.2)	38 (33.6)	93 (27.4)
61+	40 (35.4)	26 (23.0)	30 (26.5)	96 (28.3)
Weekends				
No use	29 (25.7)	39 (34.5)	28 (24.8)	96 (28.3)
≤15	13 (11.5)	7 (6.2)	6 (5.3)	26 (7.7)
16-75	34 (30.1)	33 (29.2)	37 (32.7)	104 (30.7)
76+	37 (32.7)	34 (30.1)	42 (37.2)	113 (33.3)

The percentage of adolescents doing  $\geq 360$  min of leisure-time PA/week decreased by  $-16.8\%$  from year 1 to year 3 (Table 1). Regarding SB, sitting time at school was high across secondary school years (80.2% spent  $\geq 390$  min/day sitting at school and doing homework), with a sharp decrease during the first year of university (40.7% spending  $\geq 390$  min/weekday). While adolescents spending  $> 120$  min/day screen use during weekends reduced by 17.7%, sitting while socializing and for sedentary weekday transport increased during the adolescence-university transition (Table 1).

### **Transitional Associations Between Domain-Specific Sedentary Behaviors and Tobacco Consumption**

*Domain-specific SBs and tobacco consumption during weekdays.* During weekdays, adolescents who sat more than 121 min per

day in front of a computer and more than 1 h a day socializing showed higher percentages of tobacco consumption than those who did not spend time on it (32.5% vs 22.2% computer, 40.3% vs 17.9% socializing). Higher percentages of tobacco consumption were moderately stable across years (Table 2). On the other hand, adolescents who spent more time on sedentary hobbies like music, arts, and crafts or sitting for school reasons had lower percentages of tobacco use compared to those who did not (22.9% vs 35.2% hobbies, 15.7% vs 31.8% school) (Table 2).

*Domain-specific SBs and tobacco consumption during weekends.* During weekends, adolescents with higher levels of SB spent on socialization (151+ min/day) had higher tobacco consumption compared to those who did not spend SB time socializing (38.3% vs 12.5%). Similar results were found with sedentary transport (41.2% vs 18.8%). On the other

**Table 2.** Prevalence of Lifestyle Risk Behaviors in the Specific-Domains of Sedentary Behavior.

	Habitual smokers, n (%)	Alcohol consumption, n (%)	<5 Fruit and vegetables, n (%)
Boys	40 (27.8)	36 (25.0)	105 (73.4)
Girls	54 (27.7)	25 (12.8)	143 (73.3)
Computer use			
Weekdays			
No use	2 (22.2)	0 (0.0)	5 (55.6)
≤60	38 (28.1)	20 (14.8)	92 (68.1)
61-120	27 (24.1)	22 (19.6)	85 (76.6)
121+	27 (32.5)	19 (22.9)	66 (79.5)
Weekends			
No use	6 (46.2)	1 (7.7)	9 (69.2)
≤60	31 (28.2)	23 (20.9)	74 (67.3)
61-180	35 (25.9)	23 (17.0)	100 (74.6)
181+	22 (27.2)	14 (17.3)	65 (80.2)
TV viewing terciles			
Weekdays			
No use	12 (34.3)	9 (25.7)	24 (68.6)
≤30	20 (18.9)	14 (13.2)	65 (61.9)
31-90	30 (30.6)	14 (14.3)	75 (76.5)
91+	32 (32.0)	24 (24.0)	84 (84.0)
Weekends			
No use	14 (35.9)	6 (15.4)	28 (71.8)
≤60	34 (27.4)	28 (22.6)	79 (63.7)
61-120	23 (24.2)	14 (14.7)	73 (77.7)
121+	23 (28.4)	13 (16.0)	68 (84.0)
Screen use			
Weekdays			
≤120'	59 (25.4)	22 (15.9)	87 (63.5)
>120'	35 (29.4)	39 (19.4)	161 (80.1)
Weekends			
≤120'	28 (33.3)	18 (21.4)	54 (64.3)
>120'	66 (25.9)	43 (16.9)	194 (76.4)
School/Homework terciles			
Weekdays			
≤390	35 (31.8)	29 (26.4)	87 (79.1)
391-480	48 (30.2)	25 (15.7)	112 (70.9)
481+	11 (15.7)	7 (10.0)	49 (70.0)
Weekends			
no use	12 (34.3)	12 (34.3)	30 (85.7)
≤60	42 (31.8)	32 (24.2)	95 (72.5)
61-120	19 (23.5)	9 (11.1)	58 (71.6)
121+	21 (23.1)	8 (8.8)	65 (71.4)
Socializing terciles			
Weekdays			
No use	5 (17.9)	5 (17.9)	21 (75.0)
≤30	36 (25.0)	21 (14.6)	104 (72.2)
31-60	24 (25.3)	19 (20.0)	68 (71.6)
61+	29 (40.3)	16 (22.2)	55 (77.5)
Weekends			
No use	2 (12.5)	3 (18.8)	11 (68.8)
≤60	27 (22.0)	17 (13.8)	82 (67.2)
61-150	24 (25.8)	15 (16.1)	71 (76.3)
151+	41 (38.3)	26 (24.3)	84 (78.5)

(continued)

**Table 2. (continued)**

	Habitual smokers, n (%)	Alcohol consumption, n (%)	<5 Fruit and vegetables, n (%)
Transport terciles			
Weekdays			
No use	19 (28.4)	11 (16.4)	48 (71.6)
≤15	20 (20.6)	13 (13.4)	72 (74.2)
16-30	29 (30.5)	19 (20.0)	69 (73.4)
31+	26 (32.5)	18 (22.5)	59 (73.8)
Weekends			
No use	13 (18.8)	13 (18.8)	49 (71.0)
≤15	13 (28.9)	9 (20.0)	30 (66.7)
16-60	47 (27.0)	29 (16.7)	128 (74.0)
61+	21 (41.2)	10 (19.6)	41 (80.4)
Hobbies terciles			
Weekdays			
No use	38 (35.2)	29 (26.9)	81 (75.0)
≤15	14 (33.3)	9 (21.4)	26 (63.4)
16-60	20 (21.5)	12 (12.9)	72 (77.4)
61+	22 (22.9)	11 (11.5)	69 (71.9)
Weekend			
No use	35 (36.5)	25 (26.0)	76 (79.2)
≤15	9 (34.6)	5 (19.2)	17 (68.0)
16-75	24 (23.1)	13 (12.5)	73 (70.2)
76+	26 (23.0)	18 (15.9)	82 (72.6)

Note. Cases and % of habitual smokers, weekly alcohol consumers and not meeting the 5 fruit and vegetable recommendations. All years.

hand, adolescents who spent more time sitting on the computer, doing homework and hobbies showed lower percentages of tobacco consumption than those who spent less time (27.2 vs 46.2 computer; 23.1% vs 34.3% homework; 23.0% vs 36.5% hobbies) (Table 2). After adjusting for gender, year and leisure-time PA, it was detected that girls who spent a greater amount of time watching TV were less likely to smoke than those who did not watch TV (OR=0.39, 95% CI 0.16-0.93) (Figure 1a).

In the most inactive group (<180 min/week of leisure time PA), adolescents spending no sitting time doing homework were more likely to smoke (OR=3.9, 95% CI 1.31-11.61) compared to those spending no sitting time doing homework and being in the most active group. In contrast, those who spent more time doing homework ( $\geq 480$  min/day) were less likely to smoke (OR=0.21, 95% CI 0.03-0.93) (Figure 1b).

Finally, adolescents spending 30 min or more per day sitting in motorized transport showed more likelihood of smoking compared to those who did not use motor vehicles (OR=1.69, 95% CI 1.01-2.83, 30-60 min; OR=2.2, 95% CI 1.2-4.12, 60+ min) (Figure 1c).

### Transitional Associations Between Domain-Specific SBs and Weekly Alcohol Consumption


*Domain-specific SBs and alcohol consumption during weekdays.* Adolescents sitting more than 2 h/day in front of a computer

and more than 1 h/day socializing had higher percentages of weekly alcohol consumption than those who did not (22.9% vs 0% computer, 22.2% vs 17.9% socializing). On the other hand, adolescents who had higher levels of sedentary hobbies ( $\geq 61$  min/day) and sitting for school reasons ( $\geq 481$  min/day), had lower percentages of alcohol consumption (11.5% and 10.0%) (Table 2).


Girls with higher sedentary time spent on TV viewing were less likely to consume alcohol weekly compared to girls who did not watch TV (OR=0.20, 95% CI 0.05-0.76, <30 min; OR=0.27, 95% CI 0.08-0.95, 30-60 min; and OR=0.27, 95% CI 0.07-0.99, 60+ min) (Figure 2a). And adolescents who spent at least 30 min/day doing sedentary hobbies were less likely to consume alcohol weekly compared to those who did not do any sedentary hobby (OR=0.38, 95% CI 0.17-0.81, 30-60 min; OR=0.39, 95% CI 0.17-0.86, 60+ min) (Figure 2b).

*Domain-specific SBs and alcohol consumption during weekends.* During weekends, adolescents with more time spent on sedentary socialization ( $\geq 151$  min/day) were more likely to have higher alcohol consumption compared to adolescents with less socialization time ( $\leq 60$  min/day) (24.3% vs 18.8%). On the other hand, those who spent more than 2 h/day doing homework and more than 75 min/day doing sedentary hobbies, presented lower percentages of alcohol than those spending no time on these activities (8.8% vs 34.3% homework; 15.9% vs 26.0% hobbies). Among those






**Figure 1.** Tobacco consumption risk related to SB-domains. OR (95% CI): Model 1a: tobacco = TV + gender + TV\*gender, Model 1b: tobacco = homework + year + leisure PA + homework\*leisure PA, and Model 1c: tobacco = transport + leisure PA.



**Figure 2.** Alcohol consumption risk related to SB-domains. Weekdays and weekends. OR (95% CI). Model 2a: alcohol = TV + gender + year + TV\*gender, Model 2b: alcohol = hobbies + gender + year + leisure PA + homework, and Model 1c: alcohol = homework + gender + leisure PA + homework\*gender.



**Figure 3.** Fruit and vegetable lack of consumption risk related to SB-domains. Weekdays. OR (95% CI). Model 3a, less than 5 fruits and vegetables = TV + leisure PA + TV\*leisure, Model 3b: less than 5 fruits and vegetables = screen use + year + leisure PA + screen use\*leisure PA.

spending time in sedentary hobbies, a regular reduction in alcohol consumption was detected (Table 2). After adjusting the model, we observed a protective effect of spending time doing homework in girls: those who spent more time doing homework had fewer opportunities to consume alcohol than those doing no homework (OR=0.23, 95% CI 0.06-0.88, <30 min; OR=0.22, 95% CI 0.06-0.85, 30-60 min; OR=0.13, 95% CI 0.03-0.53, >60 min) (Figure 2c).

### Transitional Associations Between Domain-Specific SBs and Fruit and Vegetable Consumption

**Domain-specific SBs and fruit and vegetable consumption during weekdays.** Adolescents who sat more than 2 h/day in front of the screen were more likely not to meet the fruit and vegetable recommendations compared to those who spent less than 2 h (80.1% vs 63.5%). On those spending more time in front of the computer, a sharp increase in not meeting the recommendations was seen across the years (Table 2).

After adjusting for gender, year and leisure-time PA, adolescents in the most active group ( $\geq 360$  min a week of leisure-time PA) who sat watching TV 60' or more per day and spent more than 2 h a day in front of a screen were less likely to meet the fruit and vegetable recommendations (OR=9.93, 95% CI 1.9-51.96; OR=3.8, 95% CI 1.72-8.45) respectively (Figure 3a).

In the most inactive group (<180 min a week of leisure-time PA) adolescents who spent less than 120 days of screen use were also less likely not to meet the fruit and vegetable recommendations (OR=4.48, 95% CI 1.54-13.03) (Figure 3b).

**Domain-specific SBs and fruit and vegetable consumption during weekends.** At weekends, adolescents who sat more than 2 h in front of a screen were more likely not to meet the fruit and vegetable recommendations (76.4%). Adolescents who spent no time doing homework and more time using sedentary transport or socializing, also showed higher percentages for not meeting the fruit and vegetable recommendations (85.7% doing homework; 78.5% socializing; and 80.4% sedentary transport) (Table 2). Among those spending more than 1 h/day using sedentary transport, the prevalence of not meeting the recommendations increased across the years (Table 2). After adjusting for gender, year, and leisure-time PA, no relation was observed between fruit and vegetable consumption and SB during weekends.

## Discussion

This study investigated the associations between changes in domain-specific SBs and changes in health-related lifestyles across the adolescence-university transition in a sample of Spanish adolescents. Adolescents and university students have more opportunities—and a greater requirement—to spend time sitting in several contexts.<sup>30</sup> This has added urgency, since post-pandemic; sedentary time increased by 52.7% in university students over this period.<sup>31</sup> Given its ubiquity, there is a need to understand whether any increases in SBs in specific contexts spills over to incrementally influence other lifestyle risk behaviors. With scarce longitudinal evidence associating SBs and lifestyle behaviors over this life period,<sup>30</sup> this study shows that SB patterns initiated *during* this time—rather than *because of* this time—affected further (un)healthy lifestyle behaviors.

### Main Findings of This Study

Three main findings were identified. The first finding highlighted that time spent on doing homework was associated with positive protective lifestyle behaviors. Time sitting doing weekend homework was protective against tobacco use in adolescents who spent little time doing leisure PA. In girls, doing sedentary weekend homework also protected against alcohol consumption. These are relevant findings since alcohol and tobacco consumption have been negatively related to academic performance and increased risk for skipping school among adolescents.<sup>32-35</sup> Similarly, students with higher grades have been reported to be less likely to engage in alcohol consumption behaviors, with males more likely to engage in alcohol consumption behaviors than females.<sup>36</sup> Taking into account previous evidence indicating that time spent in doing homework/

study without computer is positively associated with academic performance,<sup>37</sup> our results suggest that time spent sitting doing weekend homework is protective against tobacco and alcohol use. Our findings also suggest the key role leisure time PA plays in the consumption of alcohol and tobacco during this life period.<sup>37,38</sup>

The second finding highlighted that spending time doing specific SB-domains, like sedentary transportation was negatively associated with lifestyle risk factors. Specifically, sedentary weekend transport time was associated with higher tobacco consumption. Previous research has confirmed the relationship between tobacco consumption and sedentary time in adolescence<sup>39</sup> and also that tobacco use and sedentary transport are risk behaviors that have increased during the last years, especially among girls.<sup>40</sup> Our results confirm that sedentary transport need to be tackled during this transition period, particularly because of the potential role that active transportation might have for behavioral risk factor modification.<sup>41</sup>

The third finding highlighted that sedentary TV watching was protective to lifestyle behaviors -weekend tobacco consumption and weekday alcohol consumption—but only in girls. In addition, the lowest and highest tertiles for sedentary screen use and leisure-time PA were less likely not to meet the recommendations for fruit and vegetable consumption. Our results are similar to previous research in identifying associations between screen time and substance use (alcohol and/or cigarettes) among adolescents,<sup>42</sup> highlighting that overuse time spent in front of screens for leisure should be part of future interventions for tackling behavioral risk factors during this life transition period. However, the relationship between screen use and fruit and vegetable consumption followed the “Goldilocks effect”; both extremes (<180 and >360 min/week) may be at risk.

Evidence suggest that the type of screen media adolescents use while being sedentary may affect other lifestyle behaviors such as media that provide information about safe health and practices and behaviors.<sup>43-45</sup> However, our results suggested that screen time associations with lifestyle risk factors were also influenced by gender and leisure-time PA, indicating that gender and levels of leisure-time PA should be considered when designing preventive interventions targeting sedentary screen time. Public Health interventions on the transition from secondary to university should target reductions in SB-specific domains. In doing so they could influence other risk lifestyle behaviors that can begin to incubate their harm during this critical life period.<sup>13</sup>

### **Main Limitations and Strengths**

This study used self-report data which can lead to an overestimation of PA levels.<sup>46</sup> Although recall bias is common and would require validation against objective measures (ie, inclinometers or accelerometers), self-report tools allow the description of behavioral context and modes of SB.<sup>47</sup> In the future, self-report and objective methods should be

combined to accurately assess the patterns of both SBs and PA across this life period.

Although a larger sample size was preferable and no power calculation was conducted for this longitudinal study, this study presents data based on a medium-sized sample and is one of the first longitudinal studies of Spanish adolescents to address the emergence of SBs in relation to lifestyle risk behaviors. Given that SBs emerge with age, rather than at a given age, it is important to integrate a lifecourse perspective in SB-reduction interventions whenever possible.<sup>47</sup> Future studies could include a wider range of unhealthy behaviors, including fast food and/or sugary drinks.

### **Conclusions**

Given that NCDs often take years to develop, it is important to (i) establish when any unhealthy lifestyles emerge, (ii) identify any (un)healthy behavioral combinations, and (iii) intervene to offset potential long-term harm.<sup>48,49</sup> Adolescence signals the initiation of many risk behaviors<sup>50-52</sup> and therefore, the adolescence-university transition represents a critical life period to understand how lifestyle behaviors may cluster, to target disease prevention and promote sustainable health-enhancing behavior<sup>53</sup> into adulthood and for the next generations. Our findings indicate that from secondary school to university, specific SBs are associated to lifestyle risk factors some of which are gender-based and/or related to leisure-time PA. Over this transitional period, public health interventions targeting reduced SBs may bring multiple benefits by also preventing other harmful behaviors.

### **Acknowledgments**

Agència de Gestió d’Ajuts Universitaris, Universitat de Vic-Universitat Central de Catalunya, and Centres d’Estudis Sanitaris i Socials. Paul Marshall for his language help and Joan Carles Martori for his preliminary statistical analysis.

### **Author Contributions**

IAP carried out the longitudinal study, reviewed the current literature about our area of interest and drafted the manuscript; EC performed the statistical analysis, wrote the methods section and drafted the manuscript; APR participated in its design and coordination of the study, wrote the discussion part and helped to draft the manuscript. JMcK contributed to the preparation of the manuscript.


### **Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### **Funding**

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by funding from the Agència de Gestió d’Ajuts Universitaris i de Recerca (AGAUR) under grant number 2012 FI\_B 00506.

**ORCID iD**

Ignasi Arumi-Prat  <https://orcid.org/0000-0002-5839-2719>

**Supplemental Material**

Supplemental material for this article is available online.

**References**

1. Wagner KH, Brath H. A global view on the development of non communicable diseases. *Prev Med.* 2012;54 Suppl:S38-S41. doi:10.1016/j.ypmed.2011.11.012
2. Akseer N, Mehta S, Wigle J, et al. Non-communicable diseases among adolescents: current status, determinants, interventions and policies. *BMC Public Health.* 2020;20(1):1908. doi:10.1186/s12889-020-09988-5
3. van Sluijs EMF, Ekelund U, Crochemore-Silva I, et al. Physical activity behaviours in adolescence: current evidence and opportunities for intervention. *Lancet.* 2021;398(10298):429-442. doi:10.1016/S0140-6736(21)01259-9
4. Armocida B, Monasta L, Sawyer S, et al. Burden of non-communicable diseases among adolescents aged 10-24 years in the EU, 1990-2019: a systematic analysis of the Global Burden of Diseases Study 2019. *Lancet Child Adolesc Health.* 2022;6(6):367-383. doi:10.1016/S2352-4642(22)00073-6
5. Inchley J, Currie D, Budisavljevic S, Torsheim T, Jåstad A, Cosma A, et al. Spotlight on adolescent health and well-being. Findings from the 2017/2018 health behaviour in school-aged children (HBSC) survey in Europe and Canada. International report. Volume 1. Key findings. Copenhagen: WHO Regional Office for Europe; 2020. Licence: CC BY-NC-SA 3.0 IGO. Accessed July 2, 2022.
6. World Health Organization (WHO). Global action plan for the prevention and control of noncommunicable diseases 2013-2020. WHO. Published 2013. Accessed June 10, 2021. <https://apps.who.int/iris/bitstream/handle/10665/94384/?sequence=1>
7. Hardy LL, Mihrshahi S, Bellew W, Bauman A, Ding D. Children's adherence to health behavior recommendations associated with reducing risk of non-communicable disease. *Prev Med Rep.* 2017;8:279-285. doi:10.1016/j.pmedr.2017.10.006
8. Van Dyck D, De Bourdeaudhuij I, Deliens T, Deforche B. Can changes in psychosocial factors and residency explain the decrease in physical activity during the transition from high school to college or university? *Int J Behav Med.* 2015; 22(2):178-186. doi:10.1007/s12529-014-9424-4
9. Chillón P, Ortega FB, Ruiz JR, et al. Active commuting and physical activity in adolescents from Europe: results from the HELENA study. *Pediatr Exerc Sci.* 2011;23(2):207-217. doi:10.1123/pes.23.2.207
10. Prat IA, Viñolas EC, Cañas JCM, Wasley DA, Puig-Ribera A. From secondary school to university: associations between sport participation and total and domain-specific sedentary behaviours in Spanish students. *Eur J Pediatr.* 2020;179(10):1635-1645. doi:10.1007/s00431-020-03655-y
11. Grøntved A, Ried-Larsen M, Møller NC, et al. Youth screen-time behaviour is associated with cardiovascular risk in young adulthood: the European Youth Heart Study. *Eur J Prev Cardiol.* 2014;21(1):49-56. doi:10.1177/2047487312454760
12. de Almeida Silva FM, Menezes AS. Sedentary behavior, psychosocial stress indicators, and health-risk behaviors among adolescents in northeastern Brazil. *J Phys Act Health.* 2018;15(3):169-175. doi:10.1123/jpah.2015-0488
13. Bertrand L, Shaw KA, Ko J, Deprez D, Chilibeck PD, Zello GA. The impact of the coronavirus disease 2019 (COVID-19) pandemic on university students' dietary intake, physical activity, and sedentary behaviour. *Appl Physiol Nutr Metab.* 2021;46(3):265-272. doi:10.1139/apnm-2020-0990
14. Felez-Nobrega M, Bort-Roig J, Briones L, et al. Self-reported and activPALTM-monitored physical activity and sedentary behaviour in college students: not all sitting behaviours are linked to perceived stress and anxiety. *J Sports Sci.* 2020; 38(13):1566-1574. doi:10.1080/02640414.2020.1748359
15. Chen P, Jacobson KC. Developmental trajectories of substance use from early adolescence to young adulthood: gender and racial/ethnic differences. *J Adolesc Health.* 2012;50(2):154-163. doi:10.1016/j.jadohealth.2011.05.013
16. te Velde SJ, Twisk JWR, Brug J. Tracking of fruit and vegetable consumption from adolescence into adulthood and its longitudinal association with overweight [published correction appears in Br J Nutr. 2007 Oct;98(4):871]. *Br J Nutr.* 2007;98(2):431-438. doi:10.1017/S0007114507721451
17. Boers E, Afzali MH, Conrod P. A longitudinal study on the relationship between screen time and adolescent alcohol use: the mediating role of social norms. *Prev Med.* 2020;132:105992. doi:10.1016/j.ypmed.2020.105992
18. Williams GC, Battista K, Leatherdale ST. An examination of how age of onset for alcohol, cannabis, and tobacco are associated with physical activity, screen time and BMI as students are preparing to graduate from high school. *Prev Med Rep.* 2019;15:100956. doi:10.1016/j.pmedr.2019.100956
19. Mayne SL, Virudachalam S, Fiks AG. Clustering of unhealthy behaviors in a nationally representative sample of U.S. children and adolescents. *Prev Med.* 2020;130:105892. doi:10.1016/j.ypmed.2019.105892
20. Comín Bertrán E, Torrubia Beltri R, Mor Sancho J, Villalbí H Jr, Nebot Adell M. Fiabilidad de un cuestionario autoadministrado para investigar el nivel de ejercicio y el consumo de tabaco y de alcohol entre escolares. *Med Clin (Barc).* 1997;108(8):293-298.
21. Hall JN, Moore S, Harper SB, Lynch JW. Global variability in fruit and vegetable consumption. *Am J Prev Med.* 2009; 36(5):402-409.e5. doi:10.1016/j.amepre.2009.01.029
22. Peltzer K, Pengpid S, Samuels TA, et al. Prevalence of overweight/obesity and its associated factors among university students from 22 countries. *Int J Environ Res Public Health.* 2014;11(7):7425-7441. doi:10.3390/ijerph110707425
23. Roman-Viñas B, Serra-Majem L, Hagströmer M, Ribas-Barba L, Sjöström M, Segura-Cardona R. International Physical Activity Questionnaire: reliability and validity in a Spanish population. *Eur J Sport Sci.* 2010;10(5):297-304. doi:10.1080/17461390903426667
24. Hagströmer M, Bergman P, De Bourdeaudhuij I, et al. Concurrent validity of a modified version of the International Physical Activity Questionnaire (IPAQ-A) in European adolescents: the HELENA study. *Int J Obes.* 2008;32(Suppl 5):S42-S48. doi:10.1038/ijo.2008.182
25. Joe L, Carlson J. Active where? Individual item reliability report. *Active Living Research.* 2010: 2-3. [https://drjimsallis.org/Documents/Measures\\_documents/ActiveWhere\\_item\\_reliability\\_overview.pdf](https://drjimsallis.org/Documents/Measures_documents/ActiveWhere_item_reliability_overview.pdf)

26. Gorely T, Marshall SJ, Biddle SJ, Cameron N. Patterns of sedentary behaviour and physical activity among adolescents in the United Kingdom: project STIL. *J Behav Med.* 2007;30(6): 521-531. doi:10.1007/s10865-007-9126-3
27. Souza SD, Marques KC, Reuter CP. Screen time above recommendations in children and adolescents: analysis of the associated nutritional, behavioral and parental factors. *J Hum Growth Dev.* 2020;30(3):363-370. doi:10.7322/jhgd.v30.11067
28. Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics.* 1986;42(1):121-130.
29. Hosmer JDW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*, Vol. 398, 3rd ed. John Wiley & Sons; 2013.
30. Foley LS, Maddison R, Jiang Y, Olds T, Ridley K. It's not just the television: survey analysis of sedentary behaviour in New Zealand young people. *Int J Behav Nutr Phys Act.* 2011;8:132. doi:10.1186/1479-5868-8-132
31. Rodríguez-Larrad A, Mañas A, Labayen I, et al. Impact of COVID-19 confinement on physical activity and sedentary behaviour in Spanish university students: role of gender. *Int J Environ Res Public Health.* 2021;18(2):369. doi:10.3390/ijerph18020369
32. Joseph Onyebuchukwu I. The effect of alcohol consumption on the academic performance of undergraduate students. *Psychol Behav Sci.* 2015;4(4):147-153. doi:10.11648/j.pbs.20150404.12
33. Piazza-Gardner AK, Barry AE, Merianos AL. Assessing drinking and academic performance among a nationally representative sample of college students. *J Drug Issues.* 2016;46(4):347-353. doi:10.1177/0022042616659757
34. Robert PO, Kuipers MAG, Rathmann K, et al. Academic performance and adolescent smoking in 6 European cities: the role of friendship ties. *Int J Adolesc Youth.* 2019;24(1): 125-135. doi:10.1080/02673843.2018.1475288
35. Bugbee BA, Beck KH, Fryer CS, Arria AM. Substance use, academic performance, and academic engagement among high school seniors. *J Sch Health.* 2019;89(2):145-156. doi:10.1111/josh.12723
36. El Ansari W, Salam A, Suominen S. Is alcohol consumption associated with poor perceived academic performance? Survey of undergraduates in Finland. *Int J Environ Res Public Health.* 2020;17(4):1369. doi:10.3390/ijerph17041369
37. Sekulic D, Ostojic M, Vasilj M, Coric S, Zenic N. Gender-specific predictors of cigarette smoking in adolescents: an analysis of sport participation, parental factors and religiosity as protective/risk factors. *J Subst Use.* 2014;19(1-2):89-94.
38. Peltzer K. Leisure time physical activity and sedentary behavior and substance use among in-school adolescents in eight African countries. *Int J Behav Med.* 2010;17(4):271-278. doi:10.1007/s12529-009-9073-1
39. Guedes DP, Souza MV, Ferreirinha JE, Silva AJ. Physical activity and determinants of sedentary behavior in Brazilian adolescents from an underdeveloped region. *Percept Mot Skills.* 2012;114(2):542-552. doi:10.2466/06.13.17.PMS.114.2.542-552
40. Peltzer K, Pengpid S. Health risk behaviours among adolescents in Argentina: trends between 2007, 2012 and 2018 national cross-sectional school surveys. *BMC Pediatr.* 2021;21(1): 464. doi:10.1186/s12887-021-02929-0
41. Furie GL, Desai MM. Active transportation and cardiovascular disease risk factors in U.S. adults. *Am J Prev Med.* 2012;43(6):621-628. doi:10.1016/j.amepre.2012.06.034
42. Christodoulou G, Majmundar A, Chou CP, Pentz MA. Anhedonia, screen time, and substance use in early adolescents: a longitudinal mediation analysis. *J Adolesc.* 2020;78:24-32. doi:10.1016/j.adolescence.2019.11.007
43. Kenney EL, Gortmaker SL. United States adolescents' television, computer, videogame, smartphone, and tablet use: associations with sugary drinks, sleep, physical activity, and obesity. *J Pediatr.* 2017;182:144-149. doi:10.1016/j.jpeds.2016.11.015
44. Lesjak V, Stanojević-Jerković O. Physical activity, sedentary behavior and substance use among adolescents in Slovenian urban area. *Zdr Varst.* 2015;54(3):168-174. doi:10.1515/sjph-2015-0024
45. Grant JE, Lust K, Chamberlain SR. Problematic smartphone use associated with greater alcohol consumption, mental health issues, poorer academic performance, and impulsivity. *J Behav Addict.* 2019;8(2):335-342. doi:10.1556/2006.8.2019.32
46. Ekelund U, Tomkinson G, Armstrong N. What proportion of youth are physically active? Measurement issues, levels and recent time trends. *Br J Sports Med.* 2011;45(11):859-865. doi:10.1136/bjsports-2011-090190
47. Barnett TA, Kelly AS, Young DR, et al. Sedentary behaviors in today's youth: approaches to the prevention and management of childhood obesity: a scientific statement from the American Heart Association. *Circulation.* 2018;138(11):e142-e159. doi:10.1161/CIR.0000000000000591
48. Meader N, King K, Moe-Byrne T, et al. A systematic review on the clustering and co-occurrence of multiple risk behaviours. *BMC Public Health.* 2016;16:657. doi:10.1186/s12889-016-3373-6. Published 2016 Jul 29.
49. Noble N, Paul C, Turon H, Oldmeadow C. Which modifiable health risk behaviours are related? A systematic review of the clustering of smoking, nutrition, alcohol and physical activity ('SNAP') health risk factors. *Prev Med.* 2015;81:16-41. doi:10.1016/j.ypmed.2015.07.003
50. Hale DR, Viner RM. The correlates and course of multiple health risk behaviour in adolescence. *BMC Public Health.* 2016;16:458. doi:10.1186/s12889-016-3120-z
51. Hayes G, Dowd KP, MacDonncha C, Donnelly AE. Tracking of physical activity and sedentary behavior from adolescence to young adulthood: a systematic literature review. *J Adolesc Health.* 2019;65(4):446-454. doi:10.1016/j.jadohealth.2019.03.013
52. Molina-García J, Queralt A, Castillo I, Sallis JF. Changes in physical activity domains during the transition out of high school: psychosocial and environmental correlates. *J Phys Act Health.* 2015;12(10):1414-1420. doi:10.1123/jpah.2014-0412
53. Hills AP, Dengel DR, Lubans DR. Supporting public health priorities: recommendations for physical education and physical activity promotion in schools. *Prog Cardiovasc Dis.* 2015;57(4):368-374. doi:10.1016/j.pcad.2014.09.010