



LEEDS
BECKETT
UNIVERSITY

Citation:

Chantler, S (2018) The effect of an acute bout of rugby training on plasma concentrations of intestinal fatty acid binding protein. In: International Sport and Exercise Nutrition Conference 2018, 18 December 2018 - 20 December 2018, Newcastle, United Kingdom. (Unpublished)

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/9209/>

Document Version:

Conference or Workshop Item (Presentation)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.


The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.



The effect of an acute bout of rugby training on plasma concentrations of intestinal fatty acid binding protein



@sarahchantler
@23benjones
@K_Deighton

LEEDS BECKETT UNIVERSITY
CARNEGIE SCHOOL
OF SPORT

Sarah Chantler^{1,2}, Alex Griffiths¹, Greg Roe^{1,3}, Padraic Phibbs^{1,2}, Cameron Owen^{1,2}, Carlos Ramirez-Lopez^{1,2}, Glen Davison⁴, Ben Jones^{1,2} & Kevin Deighton¹



1
LEEDS BECKETT UNIVERSITY



2



3

4
University of Kent

Introduction

- Exercise appears to elicit damage to the gut endothelial cell lining as demonstrated through increases in concentration of plasma intestinal fatty acid binding protein (i-FABP).
- The majority of previous research has focussed on endurance exercise, with limited data on high intensity contact sports.
- The high intensity nature of rugby exercise, combined with trauma from collisions, suggests that this may cause considerable increases in gastrointestinal damage and permeability
- Therefore, the purpose of the study was to investigate the effects of a standardised rugby conditioning session on plasma i-FABP concentrations as a marker of gut endothelial damage.

Methods

- Nineteen academy level male rugby players completed a standardised high intensity rugby training session at the start of their preseason training period. Players were rotated through attacking and defending positions (Figure 1).
- The participants arrived after an overnight fast and abstaining from alcohol, spicy food, caffeine and exercise in the previous 24-hours.
- A venous blood sample and body weight were taken immediately before and after exercise.
- Skin fold measures were also taken by an ISAK accredited practitioner.




Figure 1: Schematic representation of the rugby training session

Results

Table 1: Anthropometric characteristics of academy level rugby participants

	All (n=19)	Forwards (n=9)	Backline (n=10)
Age (years)	20 (1.2)	20 (1.2)	20 (1.2)
Weight (kg)	100.0 (13.9)	111.8 (7.6)	89.3 (8.4)
Height (cm)	184.4 (7.2)	188.6 (7.1)	181.1 (5.6)
Σ SF (mm)	94.6 (32.5)	119.1 (26.8)	72.6 (18.5)

Data presented as mean (±SD) Σ SF, skin folds taken over 8 sites




Figure 2: Plasma concentrations of intestinal fatty acid binding protein (i-FABP) before and after an acute bout of rugby training. Data presented as mean (±SD) as well as individual changes for plasma i-FABP ; n=19, *p=0.003

Discussion

- There was a significant increase in plasma i-FABP concentrations in response to an acute bout of rugby training, which is indicative of damage to the gut endothelial lining.
- Further research is needed to determine if this damage is associated with increased gut permeability and how this is related to gastrointestinal symptoms experienced by rugby players around training.

This work is supported by the Carnegie School of Sport and the Yorkshire Carnegie Rugby club



INTERNATIONAL
SPORT + EXERCISE NUTRITION
CONFERENCE 2018

Sarah Chantler^{1,2}, Alex Griffiths¹, Greg Roe^{1,3}, Padraic Phibbs^{1,2}, Cameron Owen^{1,2}, Carlos Ramirez-Lopez^{1,2}, Glen Davison⁴, Ben Jones^{1,2} & Kevin Deighton¹



Introduction

- Exercise appears to elicit damage to the gut endothelial cell lining as demonstrated through increases in concentration of plasma intestinal fatty acid binding protein (i-FABP).
- The majority of previous research has focussed on endurance exercise, with limited data on high intensity contact sports.
- The high intensity nature of rugby exercise, combined with trauma from collisions, suggests that this may cause considerable increases in gastrointestinal damage and permeability
- Therefore, the purpose of the study was to investigate the effects of a standardised rugby conditioning session on plasma i-FABP concentrations as a marker of gut endothelial damage.

Methods

- Nineteen academy level male rugby players completed a standardised high intensity rugby training session at the start of their preseason training period. Players were rotated through attacking and defending positions (Figure 1).
- The participants arrived after an overnight fast and abstaining from alcohol, spicy food, caffeine and exercise in the previous 24-hours.
- A venous blood sample and body weight were taken immediately before and after exercise.
- Skin fold measures were also taken by an ISAK accredited practitioner.




Figure 1: Schematic representation of the rugby training session

Results

Table 1: Anthropometric characteristics of academy level rugby participants

	All (n=19)	Forwards (n=9)	Backline (n=10)
Age (years)	20 (1.2)	20 (1.2)	20 (1.2)
Weight (kg)	100.0 (13.9)	111.8 (7.6)	89.3 (8.4)
Height (cm)	184.4 (7.2)	188.6 (7.1)	181.1 (5.6)
Σ SF (mm)	94.6 (32.5)	119.1 (26.8)	72.6 (18.5)

Data presented as mean (±SD) Σ SF, skin folds taken over 8 sites




Figure 2: Plasma concentrations of intestinal fatty acid binding protein (i-FABP) before and after an acute bout of rugby training. Data presented as mean (±SD) as well as individual changes for plasma i-FABP ; n=19, *p=0.003


Discussion

- There was a significant increase in plasma i-FABP concentrations in response to an acute bout of rugby training, which is indicative of damage to the gut endothelial lining.
- Further research is needed to determine if this damage is associated with increased gut permeability and how this is related to gastrointestinal symptoms experienced by rugby players around training.

This work is supported by the Carnegie School of Sport and the Yorkshire Carnegie Rugby club



The effect of an acute bout of rugby training on plasma concentrations of intestinal fatty acid binding protein



@sarahchantler
@23benjones
@K_Deighton

LEEDS BECKETT UNIVERSITY
CARNEGIE SCHOOL
OF SPORT

Sarah Chantler^{1,2}, Ben Jones^{1,2}, Greg Roe^{1,3}, Padraic Phibbs^{1,2}, Cameron Owen^{1,2},
Alex Griffiths¹, Carlos Ramirez-Lopez^{1,2}, Glen Davison⁴, & Kevin Deighton¹



1
LEEDS
BECKETT
UNIVERSITY



2



3

4
University of
Kent

Introduction

- Exercise appears to elicit damage to the gut endothelial cell lining as demonstrated through increases in concentration of plasma intestinal fatty acid binding protein (i-FABP).
- The majority of previous research has focussed on endurance exercise, with limited data on high intensity contact sports.
- The high intensity nature of rugby exercise, combined with trauma from collisions, suggests that this may cause considerable increases in gastrointestinal damage and permeability.
- Therefore, the purpose of the study was to investigate the effect of a standardised rugby conditioning session on plasma i-FABP concentrations as a marker of gut endothelial damage.

Methods

- Nineteen academy level male rugby players completed a standardised high intensity rugby training session at the start of their preseason training period. Players were rotated through attacking and defending positions (Figure 1).




Figure 1: Schematic representation of the rugby training session A, attacker, D1, D2, D3, defender. The ball moves from position 1 to 2, and the players move~@

- The participants arrived after an overnight fast and abstaining from alcohol, spicy food, caffeine and exercise in the previous 24-hours.
- A venous blood sample were taken immediately before and after exercise.

Results

Table 1: Anthropometric characteristics of the participants

	All (n=19)	Forwards (n=9)	Backline (n=10)
Age (years)	20 (1)	20 (1.2)	20 (1.2)
Weight (kg)	100.0 (14)	111.8 (7.6)	89.3 (8.4)
Height (cm)	184.4 (7)	188.6 (7.1)	181.1 (5.6)
Σ SF (mm)	94.6 (33)	119.1 (26.8)	72.6 (18.5)

Data presented as mean (±SD) Σ SF, skin folds taken over 8 sites




Figure 2: Plasma concentrations of intestinal fatty acid binding protein (i-FABP) before and after an acute bout of rugby training. Data presented as mean (±SD) as well as individual changes for plasma i-FABP ; n=19, *p=0.003

Discussion

- There was a significant increase in plasma i-FABP concentrations in response to an acute bout of rugby training, which is indicative of damage to the gut endothelial lining.
- Further research is needed to determine if this damage is associated with increased gut permeability and how this is related to gastrointestinal symptoms experienced by rugby players around training.

This work is supported by the Carnegie School of Sport and the Yorkshire Carnegie Rugby club